

DEX-0291

363 596 CD

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 August 2002 (22.08.2002)

PCT

(10) International Publication Number
WO 02/064611 A1

(51) International Patent Classification⁷: C07H 21/02,
21/04

San Francisco, CA 94081 (US). SUN, Yongming [US/US];
869 S. Winchester Boulevard, Apartment 260, San Jose,
CA 95128 (US). LIU, Chenghua [CN/US]; 1125 Ranchero
Way #14, San Jose, CA 95117 (US).

(21) International Application Number: PCT/US02/04197

(22) International Filing Date: 12 February 2002 (12.02.2002)

(74) Agents: LICATA, Jane, Massey et al.; Licata & Tyrrell
P.C., 66 E. Main Street, Marlton, NJ 08053 (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/268,292 13 February 2001 (13.02.2001) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:

US 60/268,292 (CIP)
Filed on 13 February 2001 (13.02.2001)

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): DI-
ADEXUS, INC. [US/US]; 343 Oyster Point Boulevard,
South San Francisco, CA 94080 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): SALCEDA, Susana
[AR/US]; 4118 Crescendo Avenue, San Jose, CA 95136
(US). MACINA, Roberto, A. [AR/US]; 4118 Crescendo
Avenue, San Jose, CA 95136 (US). HU, Ping [US/US]; 108
Palmer Street, San Ramon, CA 94583 (US). RECIPON,
Herve [FR/US]; 85 Fortuna Avenue, San Francisco, CA
94115 (US). KARRA, Kalpana [IN/US]; 6511 Rainbow
Drive, San Jose, CA 95129 (US). CAFFERKEY, Robert
[IE/US]; 849 West Orange Avenue, Apartment 1030, South

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/064611 A1

(54) Title: COMPOSITIONS AND METHODS RELATING TO BREAST SPECIFIC GENES AND PROTEINS

(57) Abstract: The present invention relates to newly identified nucleic acids and polypeptides present in normal and neoplastic breast cells, including fragments, variants and derivatives of the nucleic acids and polypeptides. The present invention also relates to antibodies to the polypeptides of the invention, as well as agonists and antagonists of the polypeptides of the invention. The invention also relates to compositions comprising the nucleic acids, polypeptides, antibodies, variants, derivatives, agonists and antagonists of the invention and methods for the use of these compositions. These uses include identifying, diagnosing, monitoring, staging, imaging and treating breast cancer and non-cancerous disease states in breast tissue, identifying breast tissue, monitoring and identifying and/or designing agonists and antagonists of polypeptides of the invention. The uses also include gene therapy, production of transgenic animals and cells, and production of engineered breast tissue for treatment and research.

**COMPOSITIONS AND METHODS
RELATING TO BREAST SPECIFIC GENES AND PROTEINS**

This application claims the benefit of priority from U.S. Provisional Application
5 Serial No. 60/268,292 filed February 13, 2001, which is herein incorporated by reference
in its entirety.

FIELD OF THE INVENTION

The present invention relates to newly identified nucleic acid molecules and
polypeptides present in normal and neoplastic breast cells, including fragments, variants
10 and derivatives of the nucleic acids and polypeptides. The present invention also relates
to antibodies to the polypeptides of the invention, as well as agonists and antagonists of
the polypeptides of the invention. The invention also relates to compositions comprising
the nucleic acids, polypeptides, antibodies, variants, derivatives, agonists and antagonists
of the invention and methods for the use of these compositions. These uses include
15 identifying, diagnosing, monitoring, staging, imaging and treating breast cancer and non-
cancerous disease states in breast tissue, identifying breast tissue and monitoring and
identifying and/or designing agonists and antagonists of polypeptides of the invention.
The uses also include gene therapy, production of transgenic animals and cells, and
production of engineered breast tissue for treatment and research.

20 **BACKGROUND OF THE INVENTION**

Excluding skin cancer, breast cancer, also called mammary tumor, is the most
common cancer among women, accounting for a third of the cancers diagnosed in the
United States. One in nine women will develop breast cancer in her lifetime and about
192,000 new cases of breast cancer are diagnosed annually with about 42,000 deaths.
25 Bevers, *Primary Prevention of Breast Cancer*, in *BREAST CANCER*, 20-54 (Kelly K Hunt
et al., ed., 2001); Kochanek et al., 49 Nat'l. Vital Statistics Reports 1, 14 (2001).

In the treatment of breast cancer, there is considerable emphasis on detection and
risk assessment because early and accurate staging of breast cancer has a significant
impact on survival. For example, breast cancer detected at an early stage (stage T0,
30 discussed below) has a five-year survival rate of 92%. Conversely, if the cancer is not
detected until a late stage (i.e., stage T4), the five-year survival rate is reduced to 13%.
AJCC Cancer Staging Handbook pp. 164-65 (Irvin D. Fleming et al. eds., 5th ed. 1998).
Some detection techniques, such as mammography and biopsy, involve increased

-2-

discomfort, expense, and/or radiation, and are only prescribed only to patients with an increased risk of breast cancer.

Current methods for predicting or detecting breast cancer risk are not optimal. One method for predicting the relative risk of breast cancer is by examining a patient's risk factors and pursuing aggressive diagnostic and treatment regimens for high risk patients. A patient's risk of breast cancer has been positively associated with increasing age, nulliparity, family history of breast cancer, personal history of breast cancer, early menarche, late menopause, late age of first full term pregnancy, prior proliferative breast disease, irradiation of the breast at an early age and a personal history of malignancy. Lifestyle factors such as fat consumption, alcohol consumption, education, and socioeconomic status have also been associated with an increased incidence of breast cancer although a direct cause and effect relationship has not been established. While these risk factors are statistically significant, their weak association with breast cancer limited their usefulness. Most women who develop breast cancer have none of the risk factors listed above, other than the risk that comes with growing older. NIH Publication No. 00-1556 (2000).

Current screening methods for detecting cancer, such as breast self exam, ultrasound, and mammography have drawbacks that reduce their effectiveness or prevent their widespread adoption. Breast self exams, while useful, are unreliable for the detection of breast cancer in the initial stages where the tumor is small and difficult to detect by palpitation. Ultrasound measurements require skilled operators at an increased expense. Mammography, while sensitive, is subject to over diagnosis in the detection of lesions that have questionable malignant potential. There is also the fear of the radiation used in mammography because prior chest radiation is a factor associated with an increase incidence of breast cancer.

At this time, there are no adequate methods of breast cancer prevention. The current methods of breast cancer prevention involve prophylactic mastectomy (mastectomy performed before cancer diagnosis) and chemoprevention (chemotherapy before cancer diagnosis) which are drastic measures that limit their adoption even among women with increased risk of breast cancer. Bevers, *supra*.

A number of genetic markers have been associated with breast cancer. Examples of these markers include carcinoembryonic antigen (CEA) (Mughal et al., 249 JAMA 1881 (1983)) MUC-1 (Frische and Liu, 22 J. Clin. Ligand 320 (2000)), HER-2/neu (Haris et al., 15 Proc.Am.Soc.Clin.Oncology. A96 (1996)), uPA, PAI-1, LPA, LPC,

RAK and BRCA (Esteva and Fritsche, *Serum and Tissue Markers for Breast Cancer*, in BREAST CANCER, 286-308 (2001)). These markers have problems with limited sensitivity, low correlation, and false negatives which limit their use for initial diagnosis. For example, while the BRCA1 gene mutation is useful as an indicator of an increased risk for breast cancer, it has limited use in cancer diagnosis because only 6.2 % of breast cancers are BRCA1 positive. Malone et al., 279 JAMA 922 (1998). See also, Mewman et al., 279 JAMA 915 (1998) (correlation of only 3.3%).

Breast cancers are diagnosed into the appropriate stage categories recognizing that different treatments are more effective for different stages of cancer. Stage TX indicates that primary tumor cannot be assessed (i.e., tumor was removed or breast tissue was removed). Stage T0 is characterized by abnormalities such as hyperplasia but with no evidence of primary tumor. Stage Tis is characterized by carcinoma in situ, intraductal carcinoma, lobular carcinoma in situ, or Paget's disease of the nipple with no tumor. Stage T1 is characterized as having a tumor of 2 cm or less in the greatest dimension. Within stage T1, Tmic indicates microinvasion of 0.1 cm or less, T1a indicates a tumor of between 0.1 to 0.5 cm, T1b indicates a tumor of between 0.5 to 1 cm, and T1c indicates tumors of between 1 cm to 2 cm. Stage T2 is characterized by tumors from 2 cm to 5 cm in the greatest dimension. Tumors greater than 5 cm in size are classified as stage T4. Within stage T4, T4a indicates extension of the tumor to the chest wall, T4b indicates edema or ulceration of the skin of the breast or satellite skin nodules confined to the same breast, T4c indicates a combination of T4a and T4b, and T4d indicates inflammatory carcinoma. AJCC Cancer Staging Handbook pp. 159-70 (Irvin D. Fleming et al. eds., 5th ed. 1998). In addition to standard staging, breast tumors may be classified according to their estrogen receptor and progesterone receptor protein status. Fisher et al., 7 Breast Cancer Research and Treatment 147 (1986). Additional pathological status, such as HER2/neu status may also be useful. Thor et al., 90 J.Nat'l.Cancer Inst. 1346 (1998); Paik et al., 90 J.Nat'l.Cancer Inst. 1361 (1998); Hutchins et al., 17 Proc.Am.Soc.Clin.Oncology A2 (1998).; and Simpson et al., 18 J.Clin.Oncology 2059 (2000).

In addition to the staging of the primary tumor, breast cancer metastases to regional lymph nodes may be staged. Stage NX indicates that the lymph nodes cannot be assessed (e.g., previously removed). Stage N0 indicates no regional lymph node metastasis. Stage N1 indicates metastasis to movable ipsilateral axillary lymph nodes. Stage N2 indicates metastasis to ipsilateral axillary lymph nodes fixed to one another or

to other structures. Stage N3 indicates metastasis to ipsilateral internal mammary lymph nodes. Id.

Stage determination has potential prognostic value and provides criteria for designing optimal therapy. Simpson et al., 18 J. Clin. Oncology 2059 (2000). Generally, 5 pathological staging of breast cancer is preferable to clinical staging because the former gives a more accurate prognosis. However, clinical staging would be preferred if it were as accurate as pathological staging because it does not depend on an invasive procedure to obtain tissue for pathological evaluation. Staging of breast cancer would be improved 10 by detecting new markers in cells, tissues, or bodily fluids which could differentiate between different stages of invasion. Progress in this field will allow more rapid and reliable method for treating breast cancer patients.

Treatment of breast cancer is generally decided after an accurate staging of the primary tumor. Primary treatment options include breast conserving therapy (lumpectomy, breast irradiation, and surgical staging of the axilla), and modified radical 15 mastectomy. Additional treatments include chemotherapy, regional irradiation, and, in extreme cases, terminating estrogen production by ovarian ablation.

Until recently, the customary treatment for all breast cancer was mastectomy. Fonseca et al., 127 Annals of Internal Medicine 1013 (1997). However, recent data indicate that less radical procedures may be equally effective, in terms of survival, for 20 early stage breast cancer. Fisher et al., 16 J. of Clinical Oncology 441 (1998). The treatment options for a patient with early stage breast cancer (i.e., stage Tis) may be breast-sparing surgery followed by localized radiation therapy at the breast. Alternatively, mastectomy optionally coupled with radiation or breast reconstruction may be employed. These treatment methods are equally effective in the early stages of breast 25 cancer.

Patients with stage I and stage II breast cancer require surgery with chemotherapy and/or hormonal therapy. Surgery is of limited use in Stage III and stage IV patients. Thus, these patients are better candidates for chemotherapy and radiation therapy with surgery limited to biopsy to permit initial staging or subsequent restaging because cancer 30 is rarely curative at this stage of the disease. AJCC Cancer Staging Handbook 84, ¶. 164-65 (Irvin D. Fleming et al. eds., 5th ed. 1998).

In an effort to provide more treatment options to patients, efforts are underway to define an earlier stage of breast cancer with low recurrence which could be treated with lumpectomy without postoperative radiation treatment. While a number of attempts have

been made to classify early stage breast cancer, no consensus recommendation on postoperative radiation treatment has been obtained from these studies. Page et al., 75 Cancer 1219 (1995); Fisher et al., 75 Cancer 1223 (1995); Silverstein et al., 77 Cancer 2267 (1996).

5 As discussed above, each of the methods for diagnosing and staging breast cancer is limited by the technology employed. Accordingly, there is need for sensitive molecular and cellular markers for the detection of breast cancer. There is a need for molecular markers for the accurate staging, including clinical and pathological staging, of breast cancers to optimize treatment methods. Finally, there is a need for sensitive
10 molecular and cellular markers to monitor the progress of cancer treatments, including markers that can detect recurrence of breast cancers following remission.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art from the following description. It should be understood, however, that the following description and the specific examples, while
15 indicating preferred embodiments of the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

20 SUMMARY OF THE INVENTION

The present invention solves these and other needs in the art by providing nucleic acid molecules and polypeptides as well as antibodies, agonists and antagonists, thereto that may be used to identify, diagnose, monitor, stage, image and treat breast cancer and non-cancerous disease states in breast; identify and monitor breast tissue; and identify
25 and design agonists and antagonists of polypeptides of the invention. The invention also provides gene therapy, methods for producing transgenic animals and cells, and methods for producing engineered breast tissue for treatment and research.

Accordingly, one object of the invention is to provide nucleic acid molecules that are specific to breast cells and/or breast tissue. These breast specific nucleic acids
30 (BSNAs) may be a naturally-occurring cDNA, genomic DNA, RNA, or a fragment of one of these nucleic acids, or may be a non-naturally-occurring nucleic acid molecule. If the BSNA is genomic DNA, then the BSNA is a breast specific gene (BSG). In a preferred embodiment, the nucleic acid molecule encodes a polypeptide that is specific to breast. In a more preferred embodiment, the nucleic acid molecule encodes a

-6-

polypeptide that comprises an amino acid sequence of SEQ ID NO: 172 through 295. In another highly preferred embodiment, the nucleic acid molecule comprises a nucleic acid sequence of SEQ ID NO: 1 through 171. By nucleic acid molecule, it is also meant to be inclusive of sequences that selectively hybridize or exhibit substantial sequence

5 similarity to a nucleic acid molecule encoding a BSP, or that selectively hybridize or exhibit substantial sequence similarity to a BSNA, as well as allelic variants of a nucleic acid molecule encoding a BSP, and allelic variants of a BSNA. Nucleic acid molecules comprising a part of a nucleic acid sequence that encodes a BSP or that comprises a part of a nucleic acid sequence of a BSNA are also provided.

10 A related object of the present invention is to provide a nucleic acid molecule comprising one or more expression control sequences controlling the transcription and/or translation of all or a part of a BSNA. In a preferred embodiment, the nucleic acid molecule comprises one or more expression control sequences controlling the transcription and/or translation of a nucleic acid molecule that encodes all or a fragment
15 of a BSP.

Another object of the invention is to provide vectors and/or host cells comprising a nucleic acid molecule of the instant invention. In a preferred embodiment, the nucleic acid molecule encodes all or a fragment of a BSP. In another preferred embodiment, the nucleic acid molecule comprises all or a part of a BSNA.

20 Another object of the invention is to provided methods for using the vectors and host cells comprising a nucleic acid molecule of the instant invention to recombinantly produce polypeptides of the invention.

Another object of the invention is to provide a polypeptide encoded by a nucleic acid molecule of the invention. In a preferred embodiment, the polypeptide is a BSP.

25 The polypeptide may comprise either a fragment or a full-length protein as well as a mutant protein (mutein), fusion protein, homologous protein or a polypeptide encoded by an allelic variant of a BSP.

Another object of the invention is to provide an antibody that specifically binds to a polypeptide of the instant invention..

30 Another object of the invention is to provide agonists and antagonists of the nucleic acid molecules and polypeptides of the instant invention.

Another object of the invention is to provide methods for using the nucleic acid molecules to detect or amplify nucleic acid molecules that have similar or identical nucleic acid sequences compared to the nucleic acid molecules described herein. In a

-7-

preferred embodiment, the invention provides methods of using the nucleic acid molecules of the invention for identifying, diagnosing, monitoring, staging, imaging and treating breast cancer and non-cancerous disease states in breast. In another preferred embodiment, the invention provides methods of using the nucleic acid molecules of the invention for identifying and/or monitoring breast tissue. The nucleic acid molecules of the instant invention may also be used in gene therapy, for producing transgenic animals and cells, and for producing engineered breast tissue for treatment and research.

The polypeptides and/or antibodies of the instant invention may also be used to identify, diagnose, monitor, stage, image and treat breast cancer and non-cancerous disease states in breast. The invention provides methods of using the polypeptides of the invention to identify and/or monitor breast tissue, and to produce engineered breast tissue.

The agonists and antagonists of the instant invention may be used to treat breast cancer and non-cancerous disease states in breast and to produce engineered breast tissue.

Yet another object of the invention is to provide a computer readable means of storing the nucleic acid and amino acid sequences of the invention. The records of the computer readable means can be accessed for reading and displaying of sequences for comparison, alignment and ordering of the sequences of the invention to other sequences.

DETAILED DESCRIPTION OF THE INVENTION

Definitions and General Techniques

Unless otherwise defined herein, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclatures used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well-known and commonly used in the art. The methods and techniques of the present invention are generally performed according to conventional methods well-known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. *See, e.g., Sambrook et al., Molecular Cloning: A Laboratory Manual, 2d ed., Cold Spring Harbor Laboratory Press (1989) and Sambrook et al., Molecular Cloning: A Laboratory Manual, 3d ed., Cold Spring Harbor*

Press (2001); Ausubel *et al.*, Current Protocols in Molecular Biology, Greene Publishing Associates (1992, and Supplements to 2000); Ausubel *et al.*, Short Protocols in Molecular Biology: A Compendium of Methods from Current Protocols in Molecular Biology – 4th Ed., Wiley & Sons (1999); Harlow and Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press (1990); and Harlow and Lane, Using Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press (1999); each of which is incorporated herein by reference in its entirety.

Enzymatic reactions and purification techniques are performed according to manufacturer's specifications, as commonly accomplished in the art or as described herein. The nomenclatures used in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well-known and commonly used in the art. Standard techniques are used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

The following terms, unless otherwise indicated, shall be understood to have the following meanings:

A "nucleic acid molecule" of this invention refers to a polymeric form of nucleotides and includes both sense and antisense strands of RNA, cDNA, genomic DNA, and synthetic forms and mixed polymers of the above. A nucleotide refers to a ribonucleotide, deoxynucleotide or a modified form of either type of nucleotide. A "nucleic acid molecule" as used herein is synonymous with "nucleic acid" and "polynucleotide." The term "nucleic acid molecule" usually refers to a molecule of at least 10 bases in length, unless otherwise specified. The term includes single- and double-stranded forms of DNA. In addition, a polynucleotide may include either or both naturally-occurring and modified nucleotides linked together by naturally-occurring and/or non-naturally occurring nucleotide linkages.

The nucleic acid molecules may be modified chemically or biochemically or may contain non-natural or derivatized nucleotide bases, as will be readily appreciated by those of skill in the art. Such modifications include, for example, labels, methylation, substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as uncharged linkages (*e.g.*, methyl phosphonates, phosphotriesters, phosphoramidates, carbamates, etc.), charged linkages (*e.g.*, phosphorothioates, phosphorodithioates, etc.), pendent moieties (*e.g.*, polypeptides), intercalators (*e.g.*, acridine, psoralen, etc.), chelators, alkylators, and modified linkages

-9-

(*e.g.*, alpha anomeric nucleic acids, etc.) The term “nucleic acid molecule” also includes any topological conformation, including single-stranded, double-stranded, partially duplexed, triplexed, hairpinned, circular and padlocked conformations. Also included are synthetic molecules that mimic polynucleotides in their ability to bind to a designated
5 sequence via hydrogen bonding and other chemical interactions. Such molecules are known in the art and include, for example, those in which peptide linkages substitute for phosphate linkages in the backbone of the molecule.

A “gene” is defined as a nucleic acid molecule that comprises a nucleic acid sequence that encodes a polypeptide and the expression control sequences that surround
10 the nucleic acid sequence that encodes the polypeptide. For instance, a gene may comprise a promoter, one or more enhancers, a nucleic acid sequence that encodes a polypeptide, downstream regulatory sequences and, possibly, other nucleic acid sequences involved in regulation of the expression of an RNA. As is well-known in the art, eukaryotic genes usually contain both exons and introns. The term “exon” refers to a
15 nucleic acid sequence found in genomic DNA that is bioinformatically predicted and/or experimentally confirmed to contribute a contiguous sequence to a mature mRNA transcript. The term “intron” refers to a nucleic acid sequence found in genomic DNA that is predicted and/or confirmed to not contribute to a mature mRNA transcript, but rather to be “spliced out” during processing of the transcript.

20 A nucleic acid molecule or polypeptide is “derived” from a particular species if the nucleic acid molecule or polypeptide has been isolated from the particular species, or if the nucleic acid molecule or polypeptide is homologous to a nucleic acid molecule or polypeptide isolated from a particular species.

An “isolated” or “substantially pure” nucleic acid or polynucleotide (*e.g.*, an
25 RNA, DNA or a mixed polymer) is one which is substantially separated from other cellular components that naturally accompany the native polynucleotide in its natural host cell, *e.g.*, ribosomes, polymerases, or genomic sequences with which it is naturally associated. The term embraces a nucleic acid or polynucleotide that (1) has been removed from its naturally occurring environment, (2) is not associated with all or a
30 portion of a polynucleotide in which the “isolated polynucleotide” is found in nature, (3) is operatively linked to a polynucleotide which it is not linked to in nature, (4) does not occur in nature as part of a larger sequence or (5) includes nucleotides or internucleoside bonds that are not found in nature. The term “isolated” or “substantially pure” also can be used in reference to recombinant or cloned DNA isolates, chemically synthesized

-10-

polynucleotide analogs, or polynucleotide analogs that are biologically synthesized by heterologous systems. The term "isolated nucleic acid molecule" includes nucleic acid molecules that are integrated into a host cell chromosome at a heterologous site, recombinant fusions of a native fragment to a heterologous sequence, recombinant
5 vectors present as episomes or as integrated into a host cell chromosome.

A "part" of a nucleic acid molecule refers to a nucleic acid molecule that comprises a partial contiguous sequence of at least 10 bases of the reference nucleic acid molecule. Preferably, a part comprises at least 15 to 20 bases of a reference nucleic acid molecule. In theory, a nucleic acid sequence of 17 nucleotides is of sufficient length to
10 occur at random less frequently than once in the three gigabase human genome, and thus to provide a nucleic acid probe that can uniquely identify the reference sequence in a nucleic acid mixture of genomic complexity. A preferred part is one that comprises a nucleic acid sequence that can encode at least 6 contiguous amino acid sequences (fragments of at least 18 nucleotides) because they are useful in directing the expression
15 or synthesis of peptides that are useful in mapping the epitopes of the polypeptide encoded by the reference nucleic acid. *See, e.g., Geysen et al., Proc. Natl. Acad. Sci. USA* 81:3998-4002 (1984); and United States Patent Nos. 4,708,871 and 5,595,915, the disclosures of which are incorporated herein by reference in their entireties. A part may also comprise at least 25, 30, 35 or 40 nucleotides of a reference nucleic acid molecule,
20 or at least 50, 60, 70, 80, 90, 100, 150, 200, 250, 300, 350, 400 or 500 nucleotides of a reference nucleic acid molecule. A part of a nucleic acid molecule may comprise no other nucleic acid sequences. Alternatively, a part of a nucleic acid may comprise other nucleic acid sequences from other nucleic acid molecules.

The term "oligonucleotide" refers to a nucleic acid molecule generally
25 comprising a length of 200 bases or fewer. The term often refers to single-stranded deoxyribonucleotides, but it can refer as well to single- or double-stranded ribonucleotides, RNA:DNA hybrids and double-stranded DNAs, among others. Preferably, oligonucleotides are 10 to 60 bases in length and most preferably 12, 13, 14, 15, 16, 17, 18, 19 or 20 bases in length. Other preferred oligonucleotides are 25, 30, 35,
30 40, 45, 50, 55 or 60 bases in length. Oligonucleotides may be single-stranded, *e.g.* for use as probes or primers, or may be double-stranded, *e.g.* for use in the construction of a mutant gene. Oligonucleotides of the invention can be either sense or antisense oligonucleotides. An oligonucleotide can be derivatized or modified as discussed above for nucleic acid molecules.

Oligonucleotides, such as single-stranded DNA probe oligonucleotides, often are synthesized by chemical methods, such as those implemented on automated oligonucleotide synthesizers. However, oligonucleotides can be made by a variety of other methods, including *in vitro* recombinant DNA-mediated techniques and by
5 expression of DNAs in cells and organisms. Initially, chemically synthesized DNAs typically are obtained without a 5' phosphate. The 5' ends of such oligonucleotides are not substrates for phosphodiester bond formation by ligation reactions that employ DNA ligases typically used to form recombinant DNA molecules. Where ligation of such oligonucleotides is desired, a phosphate can be added by standard techniques, such as
10 those that employ a kinase and ATP. The 3' end of a chemically synthesized oligonucleotide generally has a free hydroxyl group and, in the presence of a ligase, such as T4 DNA ligase, readily will form a phosphodiester bond with a 5' phosphate of another polynucleotide, such as another oligonucleotide. As is well-known, this reaction can be prevented selectively, where desired, by removing the 5' phosphates of the other
15 polynucleotide(s) prior to ligation.

The term "naturally-occurring nucleotide" referred to herein includes naturally-occurring deoxyribonucleotides and ribonucleotides. The term "modified nucleotides" referred to herein includes nucleotides with modified or substituted sugar groups and the like. The term "nucleotide linkages" referred to herein includes nucleotide linkages
20 such as phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate, phosphoroamidate, and the like. See *e.g.*, LaPlanche *et al. Nucl. Acids Res.* 14:9081-9093 (1986); Stein *et al. Nucl. Acids Res.* 16:3209-3221 (1988); Zon *et al. Anti-Cancer Drug Design* 6:539-568 (1991); Zon *et al.*, in Eckstein (ed.) Oligonucleotides and Analogues: A Practical
25 Approach, pp. 87-108, Oxford University Press (1991); United States Patent No. 5,151,510; Uhlmann and Peyman *Chemical Reviews* 90:543 (1990), the disclosures of which are hereby incorporated by reference.

Unless specified otherwise, the left hand end of a polynucleotide sequence in sense orientation is the 5' end and the right hand end of the sequence is the 3' end. In
30 addition, the left hand direction of a polynucleotide sequence in sense orientation is referred to as the 5' direction, while the right hand direction of the polynucleotide sequence is referred to as the 3' direction. Further, unless otherwise indicated, each nucleotide sequence is set forth herein as a sequence of deoxyribonucleotides. It is intended, however, that the given sequence be interpreted as would be appropriate to the

polynucleotide composition: for example, if the isolated nucleic acid is composed of RNA, the given sequence intends ribonucleotides, with uridine substituted for thymidine.

The term "allelic variant" refers to one of two or more alternative naturally-occurring forms of a gene, wherein each gene possesses a unique nucleotide sequence.

- 5 In a preferred embodiment, different alleles of a given gene have similar or identical biological properties.

The term "percent sequence identity" in the context of nucleic acid sequences refers to the residues in two sequences which are the same when aligned for maximum correspondence. The length of sequence identity comparison may be over a stretch of at least about nine nucleotides, usually at least about 20 nucleotides, more usually at least about 24 nucleotides, typically at least about 28 nucleotides, more typically at least about 32 nucleotides, and preferably at least about 36 or more nucleotides. There are a number of different algorithms known in the art which can be used to measure nucleotide sequence identity. For instance, polynucleotide sequences can be compared using FASTA, Gap or Bestfit, which are programs in Wisconsin Package Version 10.0, Genetics Computer Group (GCG), Madison, Wisconsin. FASTA, which includes, *e.g.*, the programs FASTA2 and FASTA3, provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences (Pearson, *Methods Enzymol.* 183: 63-98 (1990); Pearson, *Methods Mol. Biol.* 132: 185-219 (2000); Pearson, *Methods Enzymol.* 266: 227-258 (1996); Pearson, *J. Mol. Biol.* 276: 71-84 (1998); herein incorporated by reference). Unless otherwise specified, default parameters for a particular program or algorithm are used. For instance, percent sequence identity between nucleic acid sequences can be determined using FASTA with its default parameters (a word size of 6 and the NOPAM factor for the scoring matrix) or using Gap with its default parameters as provided in GCG Version 6.1, herein incorporated by reference.

A reference to a nucleic acid sequence encompasses its complement unless otherwise specified. Thus, a reference to a nucleic acid molecule having a particular sequence should be understood to encompass its complementary strand, with its complementary sequence. The complementary strand is also useful, *e.g.*, for antisense therapy, hybridization probes and PCR primers.

In the molecular biology art, researchers use the terms "percent sequence identity", "percent sequence similarity" and "percent sequence homology"

-13-

interchangeably. In this application, these terms shall have the same meaning with respect to nucleic acid sequences only.

The term "substantial similarity" or "substantial sequence similarity," when referring to a nucleic acid or fragment thereof, indicates that, when optimally aligned
5 with appropriate nucleotide insertions or deletions with another nucleic acid (or its complementary strand), there is nucleotide sequence identity in at least about 50%, more preferably 60% of the nucleotide bases, usually at least about 70%, more usually at least about 80%, preferably at least about 90%, and more preferably at least about 95-98% of the nucleotide bases, as measured by any well-known algorithm of sequence identity,
10 such as FASTA, BLAST or Gap, as discussed above.

Alternatively, substantial similarity exists when a nucleic acid or fragment thereof hybridizes to another nucleic acid, to a strand of another nucleic acid, or to the complementary strand thereof, under selective hybridization conditions. Typically, selective hybridization will occur when there is at least about 55% sequence identity,
15 preferably at least about 65%, more preferably at least about 75%, and most preferably at least about 90% sequence identity, over a stretch of at least about 14 nucleotides, more preferably at least 17 nucleotides, even more preferably at least 20, 25, 30, 35, 40, 50, 60, 70, 80, 90 or 100 nucleotides.

Nucleic acid hybridization will be affected by such conditions as salt
20 concentration, temperature, solvents, the base composition of the hybridizing species, length of the complementary regions, and the number of nucleotide base mismatches between the hybridizing nucleic acids, as will be readily appreciated by those skilled in the art. "Stringent hybridization conditions" and "stringent wash conditions" in the context of nucleic acid hybridization experiments depend upon a number of different
25 physical parameters. The most important parameters include temperature of hybridization, base composition of the nucleic acids, salt concentration and length of the nucleic acid. One having ordinary skill in the art knows how to vary these parameters to achieve a particular stringency of hybridization. In general, "stringent hybridization" is performed at about 25°C below the thermal melting point (T_m) for the specific DNA
30 hybrid under a particular set of conditions. "Stringent washing" is performed at temperatures about 5°C lower than the T_m for the specific DNA hybrid under a particular set of conditions. The T_m is the temperature at which 50% of the target sequence hybridizes to a perfectly matched probe. See Sambrook (1989), *supra*, p. 9.51, hereby incorporated by reference.

-14-

The T_m for a particular DNA-DNA hybrid can be estimated by the formula:

$$T_m = 81.5^\circ\text{C} + 16.6 (\log_{10}[\text{Na}^+]) + 0.41 (\text{fraction G} + \text{C}) - 0.63 (\% \text{ formamide}) - (600/l)$$

where l is the length of the hybrid in base pairs.

The T_m for a particular RNA-RNA hybrid can be estimated by the formula:

5 $T_m = 79.8^\circ\text{C} + 18.5 (\log_{10}[\text{Na}^+]) + 0.58 (\text{fraction G} + \text{C}) + 11.8 (\text{fraction G} + \text{C})^2 - 0.35$
(% formamide) - (820/l).

The T_m for a particular RNA-DNA hybrid can be estimated by the formula:

$$T_m = 79.8^\circ\text{C} + 18.5(\log_{10}[\text{Na}^+]) + 0.58 (\text{fraction G} + \text{C}) + 11.8 (\text{fraction G} + \text{C})^2 - 0.50$$

(% formamide) - (820/l).

10 In general, the T_m decreases by 1-1.5°C for each 1% of mismatch between two nucleic acid sequences. Thus, one having ordinary skill in the art can alter hybridization and/or washing conditions to obtain sequences that have higher or lower degrees of sequence identity to the target nucleic acid. For instance, to obtain hybridizing nucleic acids that contain up to 10% mismatch from the target nucleic acid sequence, 10-15°C
15 would be subtracted from the calculated T_m of a perfectly matched hybrid, and then the hybridization and washing temperatures adjusted accordingly. Probe sequences may also hybridize specifically to duplex DNA under certain conditions to form triplex or other higher order DNA complexes. The preparation of such probes and suitable hybridization conditions are well-known in the art.

20 An example of stringent hybridization conditions for hybridization of complementary nucleic acid sequences having more than 100 complementary residues on a filter in a Southern or Northern blot or for screening a library is 50% formamide/6X SSC at 42°C for at least ten hours and preferably overnight (approximately 16 hours). Another example of stringent hybridization conditions is 6X SSC at 68°C without
25 formamide for at least ten hours and preferably overnight. An example of moderate stringency hybridization conditions is 6X SSC at 55°C without formamide for at least ten hours and preferably overnight. An example of low stringency hybridization conditions for hybridization of complementary nucleic acid sequences having more than 100 complementary residues on a filter in a Southern or Northern blot or for screening a
30 library is 6X SSC at 42°C for at least ten hours. Hybridization conditions to identify nucleic acid sequences that are similar but not identical can be identified by experimentally changing the hybridization temperature from 68°C to 42°C while keeping the salt concentration constant (6X SSC), or keeping the hybridization temperature and salt concentration constant (e.g. 42°C and 6X SSC) and varying the formamide

concentration from 50% to 0%. Hybridization buffers may also include blocking agents to lower background. These agents are well-known in the art. *See Sambrook et al.* (1989), *supra*, pages 8.46 and 9.46-9.58, herein incorporated by reference. *See also* Ausubel (1992), *supra*, Ausubel (1999), *supra*, and Sambrook (2001), *supra*.

5 Wash conditions also can be altered to change stringency conditions. An example of stringent wash conditions is a 0.2x SSC wash at 65°C for 15 minutes (*see* Sambrook (1989), *supra*, for SSC buffer). Often the high stringency wash is preceded by a low stringency wash to remove excess probe. An exemplary medium stringency wash for duplex DNA of more than 100 base pairs is 1x SSC at 45°C for 15 minutes. An
10 exemplary low stringency wash for such a duplex is 4x SSC at 40°C for 15 minutes. In general, signal-to-noise ratio of 2x or higher than that observed for an unrelated probe in the particular hybridization assay indicates detection of a specific hybridization.

As defined herein, nucleic acid molecules that do not hybridize to each other under stringent conditions are still substantially similar to one another if they encode
15 polypeptides that are substantially identical to each other. This occurs, for example, when a nucleic acid molecule is created synthetically or recombinantly using high codon degeneracy as permitted by the redundancy of the genetic code.

Hybridization conditions for nucleic acid molecules that are shorter than 100 nucleotides in length (*e.g.*, for oligonucleotide probes) may be calculated by the formula:
20 $T_m = 81.5^\circ\text{C} + 16.6(\log_{10}[\text{Na}^+]) + 0.41(\text{fraction G+C}) - (600/N)$,
wherein N is change length and the $[\text{Na}^+]$ is 1 M or less. *See* Sambrook (1989), *supra*, p. 11.46. For hybridization of probes shorter than 100 nucleotides, hybridization is usually performed under stringent conditions (5-10°C below the T_m) using high concentrations (0.1-1.0 pmol/ml) of probe. *Id.* at p. 11.45. Determination of hybridization using
25 mismatched probes, pools of degenerate probes or “guessmers,” as well as hybridization solutions and methods for empirically determining hybridization conditions are well-known in the art. *See, e.g.*, Ausubel (1999), *supra*; Sambrook (1989), *supra*, pp. 11.45-11.57.

The term “digestion” or “digestion of DNA” refers to catalytic cleavage of the
30 DNA with a restriction enzyme that acts only at certain sequences in the DNA. The various restriction enzymes referred to herein are commercially available and their reaction conditions, cofactors and other requirements for use are known and routine to the skilled artisan. For analytical purposes, typically, 1 µg of plasmid or DNA fragment is digested with about 2 units of enzyme in about 20 µl of reaction buffer. For the

-16-

purpose of isolating DNA fragments for plasmid construction, typically 5 to 50 µg of DNA are digested with 20 to 250 units of enzyme in proportionately larger volumes. Appropriate buffers and substrate amounts for particular restriction enzymes are described in standard laboratory manuals, such as those referenced below, and they are specified by commercial suppliers. Incubation times of about 1 hour at 37°C are ordinarily used, but conditions may vary in accordance with standard procedures, the supplier's instructions and the particulars of the reaction. After digestion, reactions may be analyzed, and fragments may be purified by electrophoresis through an agarose or polyacrylamide gel, using well-known methods that are routine for those skilled in the art.

The term "ligation" refers to the process of forming phosphodiester bonds between two or more polynucleotides, which most often are double-stranded DNAs. Techniques for ligation are well-known to the art and protocols for ligation are described in standard laboratory manuals and references, such as, *e.g.*, Sambrook (1989), *supra*.

Genome-derived "single exon probes," are probes that comprise at least part of an exon ("reference exon") and can hybridize detectably under high stringency conditions to transcript-derived nucleic acids that include the reference exon but do not hybridize detectably under high stringency conditions to nucleic acids that lack the reference exon. Single exon probes typically further comprise, contiguous to a first end of the exon portion, a first intronic and/or intergenic sequence that is identically contiguous to the exon in the genome, and may contain a second intronic and/or intergenic sequence that is identically contiguous to the exon in the genome. The minimum length of genome-derived single exon probes is defined by the requirement that the exonic portion be of sufficient length to hybridize under high stringency conditions to transcript-derived nucleic acids, as discussed above. The maximum length of genome-derived single exon probes is defined by the requirement that the probes contain portions of no more than one exon. The single exon probes may contain priming sequences not found in contiguity with the rest of the probe sequence in the genome, which priming sequences are useful for PCR and other amplification-based technologies.

The term "microarray" or "nucleic acid microarray" refers to a substrate-bound collection of plural nucleic acids, hybridization to each of the plurality of bound nucleic acids being separately detectable. The substrate can be solid or porous, planar or non-planar, unitary or distributed. Microarrays or nucleic acid microarrays include all the devices so called in Schena (ed.), DNA Microarrays: A Practical Approach (Practical

Approach Series), Oxford University Press (1999); *Nature Genet.* 21(1)(suppl.):1 - 60 (1999); Schena (ed.), Microarray Biochip: Tools and Technology, Eaton Publishing Company/BioTechniques Books Division (2000). These microarrays include substrate-bound collections of plural nucleic acids in which the plurality of nucleic acids are
5 disposed on a plurality of beads, rather than on a unitary planar substrate, as is described, *inter alia*, in Brenner *et al.*, *Proc. Natl. Acad. Sci. USA* 97(4):1665-1670 (2000).

The term "mutated" when applied to nucleic acid molecules means that nucleotides in the nucleic acid sequence of the nucleic acid molecule may be inserted, deleted or changed compared to a reference nucleic acid sequence. A single alteration
10 may be made at a locus (a point mutation) or multiple nucleotides may be inserted, deleted or changed at a single locus. In addition, one or more alterations may be made at any number of loci within a nucleic acid sequence. In a preferred embodiment, the nucleic acid molecule comprises the wild type nucleic acid sequence encoding a BSP or is a BSNA. The nucleic acid molecule may be mutated by any method known in the art
15 including those mutagenesis techniques described *infra*.

The term "error-prone PCR" refers to a process for performing PCR under conditions where the copying fidelity of the DNA polymerase is low, such that a high rate of point mutations is obtained along the entire length of the PCR product. *See, e.g.*, Leung *et al.*, *Technique* 1: 11-15 (1989) and Caldwell *et al.*, *PCR Methods Applic.* 2: 28-
20 33 (1992).

The term "oligonucleotide-directed mutagenesis" refers to a process which enables the generation of site-specific mutations in any cloned DNA segment of interest. *See, e.g.*, Reidhaar-Olson *et al.*, *Science* 241: 53-57 (1988).

The term "assembly PCR" refers to a process which involves the assembly of a
25 PCR product from a mixture of small DNA fragments. A large number of different PCR reactions occur in parallel in the same vial, with the products of one reaction priming the products of another reaction.

The term "sexual PCR mutagenesis" or "DNA shuffling" refers to a method of error-prone PCR coupled with forced homologous recombination between DNA
30 molecules of different but highly related DNA sequence *in vitro*, caused by random fragmentation of the DNA molecule based on sequence similarity, followed by fixation of the crossover by primer extension in an error-prone PCR reaction. *See, e.g.*, Stemmer, *Proc. Natl. Acad. Sci. U.S.A.* 91: 10747-10751 (1994). DNA shuffling can be carried out between several related genes ("Family shuffling").

The term “*in vivo* mutagenesis” refers to a process of generating random mutations in any cloned DNA of interest which involves the propagation of the DNA in a strain of bacteria such as *E. coli* that carries mutations in one or more of the DNA repair pathways. These “mutator” strains have a higher random mutation rate than that of a wild-type parent. Propagating the DNA in a mutator strain will eventually generate random mutations within the DNA.

The term “cassette mutagenesis” refers to any process for replacing a small region of a double-stranded DNA molecule with a synthetic oligonucleotide “cassette” that differs from the native sequence. The oligonucleotide often contains completely and/or partially randomized native sequence.

The term “recursive ensemble mutagenesis” refers to an algorithm for protein engineering (protein mutagenesis) developed to produce diverse populations of phenotypically related mutants whose members differ in amino acid sequence. This method uses a feedback mechanism to control successive rounds of combinatorial cassette mutagenesis. *See, e.g., Arkin et al., Proc. Natl. Acad. Sci. U.S.A.* 89: 7811-7815 (1992).

The term “exponential ensemble mutagenesis” refers to a process for generating combinatorial libraries with a high percentage of unique and functional mutants, wherein small groups of residues are randomized in parallel to identify, at each altered position, amino acids which lead to functional proteins. *See, e.g., Delegrave et al., Biotechnology Research* 11: 1548-1552 (1993); Arnold, *Current Opinion in Biotechnology* 4: 450-455 (1993). Each of the references mentioned above are hereby incorporated by reference in its entirety.

“Operatively linked” expression control sequences refers to a linkage in which the expression control sequence is contiguous with the gene of interest to control the gene of interest, as well as expression control sequences that act in *trans* or at a distance to control the gene of interest.

The term “expression control sequence” as used herein refers to polynucleotide sequences which are necessary to affect the expression of coding sequences to which they are operatively linked. Expression control sequences are sequences which control the transcription, post-transcriptional events and translation of nucleic acid sequences. Expression control sequences include appropriate transcription initiation, termination, promoter and enhancer sequences; efficient RNA processing signals such as splicing and polyadenylation signals; sequences that stabilize cytoplasmic mRNA; sequences that

-19-

enhance translation efficiency (*e.g.*, ribosome binding sites); sequences that enhance protein stability; and when desired, sequences that enhance protein secretion. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include the promoter, ribosomal binding site, and
5 transcription termination sequence. The term “control sequences” is intended to include, at a minimum, all components whose presence is essential for expression, and can also include additional components whose presence is advantageous, for example, leader sequences and fusion partner sequences.

The term “vector,” as used herein, is intended to refer to a nucleic acid molecule
10 capable of transporting another nucleic acid to which it has been linked. One type of vector is a “plasmid”, which refers to a circular double-stranded DNA loop into which additional DNA segments may be ligated. Other vectors include cosmids, bacterial artificial chromosomes (BAC) and yeast artificial chromosomes (YAC). Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral
15 genome. Viral vectors that infect bacterial cells are referred to as bacteriophages. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication). Other vectors can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable
20 of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as “recombinant expression vectors” (or simply, “expression vectors”). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, “plasmid” and “vector” may be used interchangeably as the plasmid is the most commonly used form of vector.
25 However, the invention is intended to include other forms of expression vectors that serve equivalent functions.

The term “recombinant host cell” (or simply “host cell”), as used herein, is intended to refer to a cell into which an expression vector has been introduced. It should be understood that such terms are intended to refer not only to the particular subject cell
30 but to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term “host cell” as used herein.

-20-

As used herein, the phrase "open reading frame" and the equivalent acronym "ORF" refer to that portion of a transcript-derived nucleic acid that can be translated in its entirety into a sequence of contiguous amino acids. As so defined, an ORF has length, measured in nucleotides, exactly divisible by 3. As so defined, an ORF need not encode
5 the entirety of a natural protein.

As used herein, the phrase "ORF-encoded peptide" refers to the predicted or actual translation of an ORF.

As used herein, the phrase "degenerate variant" of a reference nucleic acid sequence intends all nucleic acid sequences that can be directly translated, using the
10 standard genetic code, to provide an amino acid sequence identical to that translated from the reference nucleic acid sequence.

The term "polypeptide" encompasses both naturally-occurring and non-naturally-occurring proteins and polypeptides, polypeptide fragments and polypeptide mutants, derivatives and analogs. A polypeptide may be monomeric or polymeric. Further, a
15 polypeptide may comprise a number of different modules within a single polypeptide each of which has one or more distinct activities. A preferred polypeptide in accordance with the invention comprises a BSP encoded by a nucleic acid molecule of the instant invention, as well as a fragment, mutant, analog and derivative thereof.

The term "isolated protein" or "isolated polypeptide" is a protein or polypeptide
20 that by virtue of its origin or source of derivation (1) is not associated with naturally associated components that accompany it in its native state, (2) is free of other proteins from the same species (3) is expressed by a cell from a different species, or (4) does not occur in nature. Thus, a polypeptide that is chemically synthesized or synthesized in a cellular system different from the cell from which it naturally originates will be
25 "isolated" from its naturally associated components. A polypeptide or protein may also be rendered substantially free of naturally associated components by isolation, using protein purification techniques well-known in the art.

A protein or polypeptide is "substantially pure," "substantially homogeneous" or "substantially purified" when at least about 60% to 75% of a sample exhibits a single
30 species of polypeptide. The polypeptide or protein may be monomeric or multimeric. A substantially pure polypeptide or protein will typically comprise about 50%, 60%, 70%, 80% or 90% W/W of a protein sample, more usually about 95%, and preferably will be over 99% pure. Protein purity or homogeneity may be indicated by a number of means well-known in the art, such as polyacrylamide gel electrophoresis of a protein sample,

followed by visualizing a single polypeptide band upon staining the gel with a stain well-known in the art. For certain purposes, higher resolution may be provided by using HPLC or other means well-known in the art for purification.

The term "polypeptide fragment" as used herein refers to a polypeptide of the
5 instant invention that has an amino-terminal and/or carboxy-terminal deletion compared to a full-length polypeptide. In a preferred embodiment, the polypeptide fragment is a contiguous sequence in which the amino acid sequence of the fragment is identical to the corresponding positions in the naturally-occurring sequence. Fragments typically are at least 5, 6, 7, 8, 9 or 10 amino acids long, preferably at least 12, 14, 16 or 18 amino acids
10 long, more preferably at least 20 amino acids long, more preferably at least 25, 30, 35, 40 or 45, amino acids, even more preferably at least 50 or 60 amino acids long, and even more preferably at least 70 amino acids long.

A "derivative" refers to polypeptides or fragments thereof that are substantially similar in primary structural sequence but which include, *e.g.*, *in vivo* or *in vitro* chemical
15 and biochemical modifications that are not found in the native polypeptide. Such modifications include, for example, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization,
20 disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as
25 arginylation, and ubiquitination. Other modification include, *e.g.*, labeling with radionuclides, and various enzymatic modifications, as will be readily appreciated by those skilled in the art. A variety of methods for labeling polypeptides and of substituents or labels useful for such purposes are well-known in the art, and include radioactive isotopes such as ^{125}I , ^{32}P , ^{35}S , and ^3H , ligands which bind to labeled
30 antiligands (*e.g.*, antibodies), fluorophores, chemiluminescent agents, enzymes, and antiligands which can serve as specific binding pair members for a labeled ligand. The choice of label depends on the sensitivity required, ease of conjugation with the primer, stability requirements, and available instrumentation. Methods for labeling polypeptides

-22-

are well-known in the art. *See* Ausubel (1992), *supra*; Ausubel (1999), *supra*, herein incorporated by reference.

The term "fusion protein" refers to polypeptides of the instant invention comprising polypeptides or fragments coupled to heterologous amino acid sequences.

5 Fusion proteins are useful because they can be constructed to contain two or more desired functional elements from two or more different proteins. A fusion protein comprises at least 10 contiguous amino acids from a polypeptide of interest, more preferably at least 20 or 30 amino acids, even more preferably at least 40, 50 or 60 amino acids, yet more preferably at least 75, 100 or 125 amino acids. Fusion proteins can be
10 produced recombinantly by constructing a nucleic acid sequence which encodes the polypeptide or a fragment thereof in frame with a nucleic acid sequence encoding a different protein or peptide and then expressing the fusion protein. Alternatively, a fusion protein can be produced chemically by crosslinking the polypeptide or a fragment thereof to another protein.

15 The term "analog" refers to both polypeptide analogs and non-peptide analogs. The term "polypeptide analog" as used herein refers to a polypeptide of the instant invention that is comprised of a segment of at least 25 amino acids that has substantial identity to a portion of an amino acid sequence but which contains non-natural amino acids or non-natural inter-residue bonds. In a preferred embodiment, the analog has the
20 same or similar biological activity as the native polypeptide. Typically, polypeptide analogs comprise a conservative amino acid substitution (or insertion or deletion) with respect to the naturally-occurring sequence. Analogs typically are at least 20 amino acids long, preferably at least 50 amino acids long or longer, and can often be as long as a full-length naturally-occurring polypeptide.

25 The term "non-peptide analog" refers to a compound with properties that are analogous to those of a reference polypeptide of the instant invention. A non-peptide compound may also be termed a "peptide mimetic" or a "peptidomimetic." Such compounds are often developed with the aid of computerized molecular modeling. Peptide mimetics that are structurally similar to useful peptides may be used to produce
30 an equivalent effect. Generally, peptidomimetics are structurally similar to a paradigm polypeptide (*i.e.*, a polypeptide that has a desired biochemical property or pharmacological activity), but have one or more peptide linkages optionally replaced by a linkage selected from the group consisting of: --CH₂NH--, --CH₂S--, --CH₂-CH₂--,
--CH=CH--(cis and trans), --COCH₂--, --CH(OH)CH₂--, and --CH₂SO--, by methods

well-known in the art. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (*e.g.*, D-lysine in place of L-lysine) may also be used to generate more stable peptides. In addition, constrained peptides comprising a consensus sequence or a substantially identical consensus
5 sequence variation may be generated by methods known in the art (Rizo *et al.*, *Ann. Rev. Biochem.* 61:387-418 (1992), incorporated herein by reference). For example, one may add internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

A "polypeptide mutant" or "mutein" refers to a polypeptide of the instant
10 invention whose sequence contains substitutions, insertions or deletions of one or more amino acids compared to the amino acid sequence of a native or wild-type protein. A mutein may have one or more amino acid point substitutions, in which a single amino acid at a position has been changed to another amino acid, one or more insertions and/or deletions, in which one or more amino acids are inserted or deleted, respectively, in the
15 sequence of the naturally-occurring protein, and/or truncations of the amino acid sequence at either or both the amino or carboxy termini. Further, a mutein may have the same or different biological activity as the naturally-occurring protein. For instance, a mutein may have an increased or decreased biological activity. A mutein has at least 50% sequence similarity to the wild type protein, preferred is 60% sequence similarity,
20 more preferred is 70% sequence similarity. Even more preferred are muteins having 80%, 85% or 90% sequence similarity to the wild type protein. In an even more preferred embodiment, a mutein exhibits 95% sequence identity, even more preferably 97%, even more preferably 98% and even more preferably 99%. Sequence similarity may be measured by any common sequence analysis algorithm, such as Gap or Bestfit.

25 Preferred amino acid substitutions are those which: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, (4) alter binding affinity or enzymatic activity, and (5) confer or modify other physicochemical or functional properties of such analogs. For example, single or multiple amino acid substitutions (preferably conservative amino acid
30 substitutions) may be made in the naturally-occurring sequence (preferably in the portion of the polypeptide outside the domain(s) forming intermolecular contacts. In a preferred embodiment, the amino acid substitutions are moderately conservative substitutions or conservative substitutions. In a more preferred embodiment, the amino acid substitutions are conservative substitutions. A conservative amino acid substitution should not

-24-

substantially change the structural characteristics of the parent sequence (e.g., a replacement amino acid should not tend to disrupt a helix that occurs in the parent sequence, or disrupt other types of secondary structure that characterizes the parent sequence). Examples of art-recognized polypeptide secondary and tertiary structures are described in Creighton (ed.), Proteins, Structures and Molecular Principles, W. H. Freeman and Company (1984); Branden *et al.* (ed.), Introduction to Protein Structure, Garland Publishing (1991); Thornton *et al.*, *Nature* 354:105-106 (1991), each of which are incorporated herein by reference.

As used herein, the twenty conventional amino acids and their abbreviations follow conventional usage. See Golub *et al.* (eds.), Immunology - A Synthesis 2nd Ed., Sinauer Associates (1991), which is incorporated herein by reference. Stereoisomers (e.g., D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as α -, α -disubstituted amino acids, N-alkyl amino acids, and other unconventional amino acids may also be suitable components for polypeptides of the present invention. Examples of unconventional amino acids include: 4-hydroxyproline, γ -carboxyglutamate, ϵ -N,N,N-trimethyllysine, ϵ -N-acetyllysine, O-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine, s-N-methylarginine, and other similar amino acids and imino acids (e.g., 4-hydroxyproline). In the polypeptide notation used herein, the lefthand direction is the amino terminal direction and the right hand direction is the carboxy-terminal direction, in accordance with standard usage and convention.

A protein has "homology" or is "homologous" to a protein from another organism if the encoded amino acid sequence of the protein has a similar sequence to the encoded amino acid sequence of a protein of a different organism and has a similar biological activity or function. Alternatively, a protein may have homology or be homologous to another protein if the two proteins have similar amino acid sequences and have similar biological activities or functions. Although two proteins are said to be "homologous," this does not imply that there is necessarily an evolutionary relationship between the proteins. Instead, the term "homologous" is defined to mean that the two proteins have similar amino acid sequences and similar biological activities or functions. In a preferred embodiment, a homologous protein is one that exhibits 50% sequence similarity to the wild type protein, preferred is 60% sequence similarity, more preferred is 70% sequence similarity. Even more preferred are homologous proteins that exhibit 80%, 85% or 90%

sequence similarity to the wild type protein. In a yet more preferred embodiment, a homologous protein exhibits 95%, 97%, 98% or 99% sequence similarity.

When "sequence similarity" is used in reference to proteins or peptides, it is recognized that residue positions that are not identical often differ by conservative amino acid substitutions. In a preferred embodiment, a polypeptide that has "sequence similarity" comprises conservative or moderately conservative amino acid substitutions. A "conservative amino acid substitution" is one in which an amino acid residue is substituted by another amino acid residue having a side chain (R group) with similar chemical properties (*e.g.*, charge or hydrophobicity). In general, a conservative amino acid substitution will not substantially change the functional properties of a protein. In cases where two or more amino acid sequences differ from each other by conservative substitutions, the percent sequence identity or degree of similarity may be adjusted upwards to correct for the conservative nature of the substitution. Means for making this adjustment are well-known to those of skill in the art. *See, e.g.*, Pearson, *Methods Mol. Biol.* 24: 307-31 (1994), herein incorporated by reference.

For instance, the following six groups each contain amino acids that are conservative substitutions for one another:

- 1) Serine (S), Threonine (T);
- 2) Aspartic Acid (D), Glutamic Acid (E);
- 3) Asparagine (N), Glutamine (Q);
- 4) Arginine (R), Lysine (K);
- 5) Isoleucine (I), Leucine (L), Methionine (M), Alanine (A), Valine (V), and
- 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W).

Alternatively, a conservative replacement is any change having a positive value in the PAM250 log-likelihood matrix disclosed in Gonnet *et al.*, *Science* 256: 1443-45 (1992), herein incorporated by reference. A "moderately conservative" replacement is any change having a nonnegative value in the PAM250 log-likelihood matrix.

Sequence similarity for polypeptides, which is also referred to as sequence identity, is typically measured using sequence analysis software. Protein analysis software matches similar sequences using measures of similarity assigned to various substitutions, deletions and other modifications, including conservative amino acid substitutions. For instance, GCG contains programs such as "Gap" and "Bestfit" which can be used with default parameters to determine sequence homology or sequence identity between closely related polypeptides, such as homologous polypeptides from

-26-

different species of organisms or between a wild type protein and a mutein thereof. *See, e.g.*, GCG Version 6.1. Other programs include FASTA, discussed *supra*.

A preferred algorithm when comparing a sequence of the invention to a database containing a large number of sequences from different organisms is the computer
 5 program BLAST, especially blastp or tblastn. *See, e.g.*, Altschul *et al.*, *J. Mol. Biol.* 215: 403-410 (1990); Altschul *et al.*, *Nucleic Acids Res.* 25:3389-402 (1997); herein incorporated by reference. Preferred parameters for blastp are:

	Expectation value:	10 (default)
	Filter:	seg (default)
10	Cost to open a gap:	11 (default)
	Cost to extend a gap:	1 (default)
	Max. alignments:	100 (default)
	Word size:	11 (default)
	No. of descriptions:	100 (default)
15	Penalty Matrix:	BLOSUM62

The length of polypeptide sequences compared for homology will generally be at least about 16 amino acid residues, usually at least about 20 residues, more usually at least about 24 residues, typically at least about 28 residues, and preferably more than about 35 residues. When searching a database containing sequences from a large number
 20 of different organisms, it is preferable to compare amino acid sequences.

Database searching using amino acid sequences can be measured by algorithms other than blastp are known in the art. For instance, polypeptide sequences can be compared using FASTA, a program in GCG Version 6.1. FASTA (*e.g.*, FASTA2 and FASTA3) provides alignments and percent sequence identity of the regions of the best
 25 overlap between the query and search sequences (Pearson (1990), *supra*; Pearson (2000), *supra*). For example, percent sequence identity between amino acid sequences can be determined using FASTA with its default or recommended parameters (a word size of 2 and the PAM250 scoring matrix), as provided in GCG Version 6.1, herein incorporated by reference.

30 An "antibody" refers to an intact immunoglobulin, or to an antigen-binding portion thereof that competes with the intact antibody for specific binding to a molecular species, *e.g.*, a polypeptide of the instant invention. Antigen-binding portions may be produced by recombinant DNA techniques or by enzymatic or chemical cleavage of intact antibodies. Antigen-binding portions include, *inter alia*, Fab, Fab', F(ab')₂, Fv,

-27-

dAb, and complementarity determining region (CDR) fragments, single-chain antibodies (scFv), chimeric antibodies, diabodies and polypeptides that contain at least a portion of an immunoglobulin that is sufficient to confer specific antigen binding to the polypeptide. An Fab fragment is a monovalent fragment consisting of the VL, VH, CL and CH1 domains; an F(ab')₂ fragment is a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; an Fd fragment consists of the VH and CH1 domains; an Fv fragment consists of the VL and VH domains of a single arm of an antibody; and a dAb fragment consists of a VH domain. *See, e.g., Ward et al., Nature* 341: 544-546 (1989).

10 By "bind specifically" and "specific binding" is here intended the ability of the antibody to bind to a first molecular species in preference to binding to other molecular species with which the antibody and first molecular species are admixed. An antibody is said specifically to "recognize" a first molecular species when it can bind specifically to that first molecular species.

15 A single-chain antibody (scFv) is an antibody in which a VL and VH region are paired to form a monovalent molecule via a synthetic linker that enables them to be made as a single protein chain. *See, e.g., Bird et al., Science* 242: 423-426 (1988); *Huston et al., Proc. Natl. Acad. Sci. USA* 85: 5879-5883 (1988). Diabodies are bivalent, bispecific antibodies in which VH and VL domains are expressed on a single polypeptide chain, but using a linker that is too short to allow for pairing between the two domains on the same chain, thereby forcing the domains to pair with complementary domains of another chain and creating two antigen binding sites. *See e.g., Holliger et al., Proc. Natl. Acad. Sci. USA* 90: 6444-6448 (1993); *Poljak et al., Structure* 2: 1121-1123 (1994). One or more CDRs may be incorporated into a molecule either covalently or noncovalently to make it an immunoadhesin. An immunoadhesin may incorporate the CDR(s) as part of a larger polypeptide chain, may covalently link the CDR(s) to another polypeptide chain, or may incorporate the CDR(s) noncovalently. The CDRs permit the immunoadhesin to specifically bind to a particular antigen of interest. A chimeric antibody is an antibody that contains one or more regions from one antibody and one or more regions from one or more other antibodies.

30 An antibody may have one or more binding sites. If there is more than one binding site, the binding sites may be identical to one another or may be different. For instance, a naturally-occurring immunoglobulin has two identical binding sites, a single-

-28-

chain antibody or Fab fragment has one binding site, while a “bispecific” or “bifunctional” antibody has two different binding sites.

An “isolated antibody” is an antibody that (1) is not associated with naturally-associated components, including other naturally-associated antibodies, that accompany it in its native state, (2) is free of other proteins from the same species, (3) is expressed by a cell from a different species, or (4) does not occur in nature. It is known that purified proteins, including purified antibodies, may be stabilized with non-naturally-associated components. The non-naturally-associated component may be a protein, such as albumin (*e.g.*, BSA) or a chemical such as polyethylene glycol (PEG).

10 A “neutralizing antibody” or “an inhibitory antibody” is an antibody that inhibits the activity of a polypeptide or blocks the binding of a polypeptide to a ligand that normally binds to it. An “activating antibody” is an antibody that increases the activity of a polypeptide.

The term “epitope” includes any protein determinant capable of specifically binding to an immunoglobulin or T-cell receptor. Epitopic determinants usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually have specific three-dimensional structural characteristics, as well as specific charge characteristics. An antibody is said to specifically bind an antigen when the dissociation constant is less than 1 μM , preferably less than 100 nM and most preferably less than 10 nM.

The term “patient” as used herein includes human and veterinary subjects.

Throughout this specification and claims, the word “comprise,” or variations such as “comprises” or “comprising,” will be understood to imply the inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers.

The term “breast specific” refers to a nucleic acid molecule or polypeptide that is expressed predominantly in the breast as compared to other tissues in the body. In a preferred embodiment, a “breast specific” nucleic acid molecule or polypeptide is expressed at a level that is 5-fold higher than any other tissue in the body. In a more preferred embodiment, the “breast specific” nucleic acid molecule or polypeptide is expressed at a level that is 10-fold higher than any other tissue in the body, more preferably at least 15-fold, 20-fold, 25-fold, 50-fold or 100-fold higher than any other tissue in the body. Nucleic acid molecule levels may be measured by nucleic acid hybridization, such as Northern blot hybridization, or quantitative PCR. Polypeptide

levels may be measured by any method known to accurately quantitate protein levels, such as Western blot analysis.

Nucleic Acid Molecules, Regulatory Sequences, Vectors, Host Cells and Recombinant Methods of Making Polypeptides

5

Nucleic Acid Molecules

One aspect of the invention provides isolated nucleic acid molecules that are specific to the breast or to breast cells or tissue or that are derived from such nucleic acid molecules. These isolated breast specific nucleic acids (BSNAs) may comprise a cDNA,
10 a genomic DNA, RNA, or a fragment of one of these nucleic acids, or may be a non-naturally-occurring nucleic acid molecule. In a preferred embodiment, the nucleic acid molecule encodes a polypeptide that is specific to breast, a breast-specific polypeptide (BSP). In a more preferred embodiment, the nucleic acid molecule encodes a polypeptide that comprises an amino acid sequence of SEQ ID NO: 172 through 295. In
15 another highly preferred embodiment, the nucleic acid molecule comprises a nucleic acid sequence of SEQ ID NO: 1 through 171.

A BSNA may be derived from a human or from another animal. In a preferred embodiment, the BSNA is derived from a human or other mammal. In a more preferred embodiment, the BSNA is derived from a human or other primate. In an even more
20 preferred embodiment, the BSNA is derived from a human.

By "nucleic acid molecule" for purposes of the present invention, it is also meant to be inclusive of nucleic acid sequences that selectively hybridize to a nucleic acid molecule encoding a BSNA or a complement thereof. The hybridizing nucleic acid molecule may or may not encode a polypeptide or may not encode a BSP. However, in a
25 preferred embodiment, the hybridizing nucleic acid molecule encodes a BSP. In a more preferred embodiment, the invention provides a nucleic acid molecule that selectively hybridizes to a nucleic acid molecule that encodes a polypeptide comprising an amino acid sequence of SEQ ID NO: 172 through 295. In an even more preferred embodiment, the invention provides a nucleic acid molecule that selectively hybridizes to a nucleic
30 acid molecule comprising the nucleic acid sequence of SEQ ID NO: 1 through 171.

In a preferred embodiment, the nucleic acid molecule selectively hybridizes to a nucleic acid molecule encoding a BSP under low stringency conditions. In a more preferred embodiment, the nucleic acid molecule selectively hybridizes to a nucleic acid molecule encoding a BSP under moderate stringency conditions. In a more preferred

-30-

embodiment, the nucleic acid molecule selectively hybridizes to a nucleic acid molecule encoding a BSP under high stringency conditions. In an even more preferred embodiment, the nucleic acid molecule hybridizes under low, moderate or high stringency conditions to a nucleic acid molecule encoding a polypeptide comprising an amino acid sequence of SEQ ID NO: 172 through 295. In a yet more preferred embodiment, the nucleic acid molecule hybridizes under low, moderate or high stringency conditions to a nucleic acid molecule comprising a nucleic acid sequence selected from SEQ ID NO: 1 through 171. In a preferred embodiment of the invention, the hybridizing nucleic acid molecule may be used to express recombinantly a polypeptide of the invention.

By "nucleic acid molecule" as used herein it is also meant to be inclusive of sequences that exhibits substantial sequence similarity to a nucleic acid encoding a BSP or a complement of the encoding nucleic acid molecule. In a preferred embodiment, the nucleic acid molecule exhibits substantial sequence similarity to a nucleic acid molecule encoding human BSP. In a more preferred embodiment, the nucleic acid molecule exhibits substantial sequence similarity to a nucleic acid molecule encoding a polypeptide having an amino acid sequence of SEQ ID NO: 172 through 295. In a preferred embodiment, the similar nucleic acid molecule is one that has at least 60% sequence identity with a nucleic acid molecule encoding a BSP, such as a polypeptide having an amino acid sequence of SEQ ID NO: 172 through 295, more preferably at least 70%, even more preferably at least 80% and even more preferably at least 85%. In a more preferred embodiment, the similar nucleic acid molecule is one that has at least 90% sequence identity with a nucleic acid molecule encoding a BSP, more preferably at least 95%, more preferably at least 97%, even more preferably at least 98%, and still more preferably at least 99%. In another highly preferred embodiment, the nucleic acid molecule is one that has at least 99.5%, 99.6%, 99.7%, 99.8% or 99.9% sequence identity with a nucleic acid molecule encoding a BSP.

In another preferred embodiment, the nucleic acid molecule exhibits substantial sequence similarity to a BSNA or its complement. In a more preferred embodiment, the nucleic acid molecule exhibits substantial sequence similarity to a nucleic acid molecule comprising a nucleic acid sequence of SEQ ID NO: 1 through 171. In a preferred embodiment, the nucleic acid molecule is one that has at least 60% sequence identity with a BSNA, such as one having a nucleic acid sequence of SEQ ID NO: 1 through 171, more preferably at least 70%, even more preferably at least 80% and even more

preferably at least 85%. In a more preferred embodiment, the nucleic acid molecule is one that has at least 90% sequence identity with a BSNA, more preferably at least 95%, more preferably at least 97%, even more preferably at least 98%, and still more preferably at least 99%. In another highly preferred embodiment, the nucleic acid
5 molecule is one that has at least 99.5%, 99.6%, 99.7%, 99.8% or 99.9% sequence identity with a BSNA.

A nucleic acid molecule that exhibits substantial sequence similarity may be one that exhibits sequence identity over its entire length to a BSNA or to a nucleic acid molecule encoding a BSP, or may be one that is similar over only a part of its length. In
10 this case, the part is at least 50 nucleotides of the BSNA or the nucleic acid molecule encoding a BSP, preferably at least 100 nucleotides, more preferably at least 150 or 200 nucleotides, even more preferably at least 250 or 300 nucleotides, still more preferably at least 400 or 500 nucleotides.

The substantially similar nucleic acid molecule may be a naturally-occurring one
15 that is derived from another species, especially one derived from another primate, wherein the similar nucleic acid molecule encodes an amino acid sequence that exhibits significant sequence identity to that of SEQ ID NO: 172 through 295 or demonstrates significant sequence identity to the nucleotide sequence of SEQ ID NO: 1 through 171. The similar nucleic acid molecule may also be a naturally-occurring nucleic acid
20 molecule from a human, when the BSNA is a member of a gene family. The similar nucleic acid molecule may also be a naturally-occurring nucleic acid molecule derived from a non-primate, mammalian species, including without limitation, domesticated species, *e.g.*, dog, cat, mouse, rat, rabbit, hamster, cow, horse and pig; and wild animals, *e.g.*, monkey, fox, lions, tigers, bears, giraffes, zebras, etc. The substantially similar
25 nucleic acid molecule may also be a naturally-occurring nucleic acid molecule derived from a non-mammalian species, such as birds or reptiles. The naturally-occurring substantially similar nucleic acid molecule may be isolated directly from humans or other species. In another embodiment, the substantially similar nucleic acid molecule may be one that is experimentally produced by random mutation of a nucleic acid molecule. In
30 another embodiment, the substantially similar nucleic acid molecule may be one that is experimentally produced by directed mutation of a BSNA. Further, the substantially similar nucleic acid molecule may or may not be a BSNA. However, in a preferred embodiment, the substantially similar nucleic acid molecule is a BSNA.

-32-

By “nucleic acid molecule” it is also meant to be inclusive of allelic variants of a BSNA or a nucleic acid encoding a BSP. For instance, single nucleotide polymorphisms (SNPs) occur frequently in eukaryotic genomes. In fact, more than 1.4 million SNPs have already identified in the human genome, International Human Genome Sequencing Consortium, *Nature* 409: 860-921 (2001). Thus, the sequence determined from one individual of a species may differ from other allelic forms present within the population. Additionally, small deletions and insertions, rather than single nucleotide polymorphisms, are not uncommon in the general population, and often do not alter the function of the protein. Further, amino acid substitutions occur frequently among natural allelic variants, and often do not substantially change protein function.

In a preferred embodiment, the nucleic acid molecule comprising an allelic variant is a variant of a gene, wherein the gene is transcribed into an mRNA that encodes a BSP. In a more preferred embodiment, the gene is transcribed into an mRNA that encodes a BSP comprising an amino acid sequence of SEQ ID NO: 172 through 295. In another preferred embodiment, the allelic variant is a variant of a gene, wherein the gene is transcribed into an mRNA that is a BSNA. In a more preferred embodiment, the gene is transcribed into an mRNA that comprises the nucleic acid sequence of SEQ ID NO: 1 through 171. In a preferred embodiment, the allelic variant is a naturally-occurring allelic variant in the species of interest. In a more preferred embodiment, the species of interest is human.

By “nucleic acid molecule” it is also meant to be inclusive of a part of a nucleic acid sequence of the instant invention. The part may or may not encode a polypeptide, and may or may not encode a polypeptide that is a BSP. However, in a preferred embodiment, the part encodes a BSP. In one aspect, the invention comprises a part of a BSNA. In a second aspect, the invention comprises a part of a nucleic acid molecule that hybridizes or exhibits substantial sequence similarity to a BSNA. In a third aspect, the invention comprises a part of a nucleic acid molecule that is an allelic variant of a BSNA. In a fourth aspect, the invention comprises a part of a nucleic acid molecule that encodes a BSP. A part comprises at least 10 nucleotides, more preferably at least 15, 17, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 300, 350, 400 or 500 nucleotides. The maximum size of a nucleic acid part is one nucleotide shorter than the sequence of the nucleic acid molecule encoding the full-length protein.

-33-

By "nucleic acid molecule" it is also meant to be inclusive of sequence that encoding a fusion protein, a homologous protein, a polypeptide fragment, a mutin or a polypeptide analog, as described below.

Nucleotide sequences of the instantly-described nucleic acids were determined by sequencing a DNA molecule that had resulted, directly or indirectly, from at least one enzymatic polymerization reaction (e.g., reverse transcription and/or polymerase chain reaction) using an automated sequencer (such as the MegaBACE™ 1000, Molecular Dynamics, Sunnyvale, CA, USA). Further, all amino acid sequences of the polypeptides of the present invention were predicted by translation from the nucleic acid sequences so determined, unless otherwise specified.

In a preferred embodiment of the invention, the nucleic acid molecule contains modifications of the native nucleic acid molecule. These modifications include nonnative internucleoside bonds, post-synthetic modifications or altered nucleotide analogues. One having ordinary skill in the art would recognize that the type of modification that can be made will depend upon the intended use of the nucleic acid molecule. For instance, when the nucleic acid molecule is used as a hybridization probe, the range of such modifications will be limited to those that permit sequence-discriminating base pairing of the resulting nucleic acid. When used to direct expression of RNA or protein *in vitro* or *in vivo*, the range of such modifications will be limited to those that permit the nucleic acid to function properly as a polymerization substrate. When the isolated nucleic acid is used as a therapeutic agent, the modifications will be limited to those that do not confer toxicity upon the isolated nucleic acid.

In a preferred embodiment, isolated nucleic acid molecules can include nucleotide analogues that incorporate labels that are directly detectable, such as radiolabels or fluorophores, or nucleotide analogues that incorporate labels that can be visualized in a subsequent reaction, such as biotin or various haptens. In a more preferred embodiment, the labeled nucleic acid molecule may be used as a hybridization probe.

Common radiolabeled analogues include those labeled with ³³P, ³²P, and ³⁵S, such as α-³²P-dATP, α-³²P-dCTP, α-³²P-dGTP, α-³²P-dTTP, α-³²P-3'dATP, α-³²P-ATP, α-³²P-CTP, α-³²P-GTP, α-³²P-UTP, α-³⁵S-dATP, α-³⁵S-GTP, α-³³P-dATP, and the like.

Commercially available fluorescent nucleotide analogues readily incorporated into the nucleic acids of the present invention include Cy3-dCTP, Cy3-dUTP, Cy5-dCTP, Cy3-dUTP (Amersham Pharmacia Biotech, Piscataway, New Jersey, USA), fluorescein-12-dUTP, tetramethylrhodamine-6-dUTP, Texas Red®-5-dUTP, Cascade

Blue®-7-dUTP, BODIPY® FL-14-dUTP, BODIPY® TMR-14-dUTP, BODIPY® TR-14-dUTP, Rhodamine Green™-5-dUTP, Oregon Green® 488-5-dUTP, Texas Red®-12-dUTP, BODIPY® 630/650-14-dUTP, BODIPY® 650/665-14-dUTP, Alexa Fluor® 488-5-dUTP, Alexa Fluor® 532-5-dUTP, Alexa Fluor® 568-5-dUTP, Alexa Fluor® 594-5-dUTP, Alexa Fluor® 546-14-dUTP, fluorescein-12-UTP, tetramethylrhodamine-6-UTP, Texas Red®-5-UTP, Cascade Blue®-7-UTP, BODIPY® FL-14-UTP, BODIPY® TMR-14-UTP, BODIPY® TR-14-UTP, Rhodamine Green™-5-UTP, Alexa Fluor® 488-5-UTP, Alexa Fluor® 546-14-UTP (Molecular Probes, Inc. Eugene, OR, USA). One may also custom synthesize nucleotides having other fluorophores. See Henegariu *et al.*, *Nature Biotechnol.* 18: 345-348 (2000), the disclosure of which is incorporated herein by reference in its entirety.

Haptens that are commonly conjugated to nucleotides for subsequent labeling include biotin (biotin-11-dUTP, Molecular Probes, Inc., Eugene, OR, USA; biotin-21-UTP, biotin-21-dUTP, Clontech Laboratories, Inc., Palo Alto, CA, USA), digoxigenin (DIG-11-dUTP, alkali labile, DIG-11-UTP, Roche Diagnostics Corp., Indianapolis, IN, USA), and dinitrophenyl (dinitrophenyl-11-dUTP, Molecular Probes, Inc., Eugene, OR, USA).

Nucleic acid molecules can be labeled by incorporation of labeled nucleotide analogues into the nucleic acid. Such analogues can be incorporated by enzymatic polymerization, such as by nick translation, random priming, polymerase chain reaction (PCR), terminal transferase tailing, and end-filling of overhangs, for DNA molecules, and *in vitro* transcription driven, *e.g.*, from phage promoters, such as T7, T3, and SP6, for RNA molecules. Commercial kits are readily available for each such labeling approach. Analogues can also be incorporated during automated solid phase chemical synthesis. Labels can also be incorporated after nucleic acid synthesis, with the 5' phosphate and 3' hydroxyl providing convenient sites for post-synthetic covalent attachment of detectable labels.

Other post-synthetic approaches also permit internal labeling of nucleic acids. For example, fluorophores can be attached using a cisplatin reagent that reacts with the N7 of guanine residues (and, to a lesser extent, adenine bases) in DNA, RNA, and PNA to provide a stable coordination complex between the nucleic acid and fluorophore label (Universal Linkage System) (available from Molecular Probes, Inc., Eugene, OR, USA and Amersham Pharmacia Biotech, Piscataway, NJ, USA); see Alers *et al.*, *Genes, Chromosomes & Cancer* 25: 301-305 (1999); Jelsma *et al.*, *J. NIH Res.* 5: 82 (1994);

Van Belkum *et al.*, *BioTechniques* 16: 148-153 (1994), incorporated herein by reference. As another example, nucleic acids can be labeled using a disulfide-containing linker (FastTag™ Reagent, Vector Laboratories, Inc., Burlingame, CA, USA) that is photo- or thermally-coupled to the target nucleic acid using aryl azide chemistry; after reduction, a
5 free thiol is available for coupling to a hapten, fluorophore, sugar, affinity ligand, or other marker.

One or more independent or interacting labels can be incorporated into the nucleic acid molecules of the present invention. For example, both a fluorophore and a moiety that in proximity thereto acts to quench fluorescence can be included to report
10 specific hybridization through release of fluorescence quenching or to report exonucleotidic excision. *See, e.g.*, Tyagi *et al.*, *Nature Biotechnol.* 14: 303-308 (1996); Tyagi *et al.*, *Nature Biotechnol.* 16: 49-53 (1998); Sokol *et al.*, *Proc. Natl. Acad. Sci. USA* 95: 11538-11543 (1998); Kostrikis *et al.*, *Science* 279: 1228-1229 (1998); Marras *et al.*, *Genet. Anal.* 14: 151-156 (1999); U. S. Patent 5,846,726; 5,925,517; 5,925,517;
15 5,723,591 and 5,538,848; Holland *et al.*, *Proc. Natl. Acad. Sci. USA* 88: 7276-7280 (1991); Heid *et al.*, *Genome Res.* 6(10): 986-94 (1996); Kuimelis *et al.*, *Nucleic Acids Symp. Ser.* (37): 255-6 (1997); the disclosures of which are incorporated herein by reference in their entireties.

Nucleic acid molecules of the invention may be modified by altering one or more
20 native phosphodiester internucleoside bonds to more nuclease-resistant, internucleoside bonds. *See* Hartmann *et al.* (eds.), Manual of Antisense Methodology: Perspectives in Antisense Science, Kluwer Law International (1999); Stein *et al.* (eds.), Applied Antisense Oligonucleotide Technology, Wiley-Liss (1998); Chadwick *et al.* (eds.), Oligonucleotides as Therapeutic Agents - Symposium No. 209, John Wiley & Son Ltd
25 (1997); the disclosures of which are incorporated herein by reference in their entireties. Such altered internucleoside bonds are often desired for antisense techniques or for targeted gene correction. *See* Gamper *et al.*, *Nucl. Acids Res.* 28(21): 4332-4339 (2000), the disclosure of which is incorporated herein by reference in its entirety.

Modified oligonucleotide backbones include, without limitation,
30 phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having

-36-

normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Representative United States patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U. S. Patents 3,687,808; 5 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050, the disclosures of which are incorporated herein by reference in their entireties. In a preferred embodiment, the modified internucleoside 10 linkages may be used for antisense techniques.

Other modified oligonucleotide backbones do not include a phosphorus atom, but have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include 15 those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; 20 and others having mixed N, O, S and CH₂ component parts. Representative U.S. patents that teach the preparation of the above backbones include, but are not limited to, U.S. Patent 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 25 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437 and 5,677,439; the disclosures of which are incorporated herein by reference in their entireties.

In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage are replaced with novel groups, such as peptide nucleic acids (PNA). In PNA compounds, the phosphodiester backbone of the nucleic acid is replaced 30 with an amide-containing backbone, in particular by repeating N-(2-aminoethyl) glycine units linked by amide bonds. Nucleobases are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone, typically by methylene carbonyl linkages. PNA can be synthesized using a modified peptide synthesis protocol. PNA oligomers can be synthesized by both Fmoc and tBoc methods. Representative U.S.

patents that teach the preparation of PNA compounds include, but are not limited to, U.S. Patent 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Automated PNA synthesis is readily achievable on commercial synthesizers (see, e.g., "PNA User's Guide," Rev. 2, February 1998, Perseptive Biosystems Part No. 5 60138, Applied Biosystems, Inc., Foster City, CA).

PNA molecules are advantageous for a number of reasons. First, because the PNA backbone is uncharged, PNA/DNA and PNA/RNA duplexes have a higher thermal stability than is found in DNA/DNA and DNA/RNA duplexes. The T_m of a PNA/DNA or PNA/RNA duplex is generally 1°C higher per base pair than the T_m of the 10 corresponding DNA/DNA or DNA/RNA duplex (in 100 mM NaCl). Second, PNA molecules can also form stable PNA/DNA complexes at low ionic strength, under conditions in which DNA/DNA duplex formation does not occur. Third, PNA also demonstrates greater specificity in binding to complementary DNA because a PNA/DNA mismatch is more destabilizing than DNA/DNA mismatch. A single mismatch in mixed 15 a PNA/DNA 15-mer lowers the T_m by 8–20°C (15°C on average). In the corresponding DNA/DNA duplexes, a single mismatch lowers the T_m by 4–16°C (11°C on average). Because PNA probes can be significantly shorter than DNA probes, their specificity is greater. Fourth, PNA oligomers are resistant to degradation by enzymes, and the lifetime of these compounds is extended both *in vivo* and *in vitro* because nucleases and proteases 20 do not recognize the PNA polyamide backbone with nucleobase sidechains. See, e.g., Ray *et al.*, *FASEB J.* 14(9): 1041-60 (2000); Nielsen *et al.*, *Pharmacol Toxicol.* 86(1): 3-7 (2000); Larsen *et al.*, *Biochim Biophys Acta.* 1489(1): 159-66 (1999); Nielsen, *Curr. Opin. Struct. Biol.* 9(3): 353-7 (1999), and Nielsen, *Curr. Opin. Biotechnol.* 10(1): 71-5 (1999), the disclosures of which are incorporated herein by reference in their entireties.

25 Nucleic acid molecules may be modified compared to their native structure throughout the length of the nucleic acid molecule or can be localized to discrete portions thereof. As an example of the latter, chimeric nucleic acids can be synthesized that have discrete DNA and RNA domains and that can be used for targeted gene repair and modified PCR reactions, as further described in U.S. Patents 5,760,012 and 5,731,181, 30 Misra *et al.*, *Biochem.* 37: 1917-1925 (1998); and Finn *et al.*, *Nucl. Acids Res.* 24: 3357-3363 (1996), the disclosures of which are incorporated herein by reference in their entireties.

Unless otherwise specified, nucleic acids of the present invention can include any topological conformation appropriate to the desired use; the term thus explicitly

-38-

comprehends, among others, single-stranded, double-stranded, triplexed, quadruplexed, partially double-stranded, partially-triplexed, partially-quadruplexed, branched, hairpinned, circular, and padlocked conformations. Padlock conformations and their utilities are further described in Banér *et al.*, *Curr. Opin. Biotechnol.* 12: 11-15 (2001);

5 Escude *et al.*, *Proc. Natl. Acad. Sci. USA* 14: 96(19):10603-7 (1999); Nilsson *et al.*, *Science* 265(5181): 2085-8 (1994), the disclosures of which are incorporated herein by reference in their entireties. Triplex and quadruplex conformations, and their utilities, are reviewed in Praseuth *et al.*, *Biochim. Biophys. Acta.* 1489(1): 181-206 (1999); Fox, *Curr. Med. Chem.* 7(1): 17-37 (2000); Kochetkova *et al.*, *Methods Mol. Biol.* 130: 189-201

10 (2000); Chan *et al.*, *J. Mol. Med.* 75(4): 267-82 (1997), the disclosures of which are incorporated herein by reference in their entireties.

Methods for Using Nucleic Acid Molecules as Probes and Primers

The isolated nucleic acid molecules of the present invention can be used as

15 hybridization probes to detect, characterize, and quantify hybridizing nucleic acids in, and isolate hybridizing nucleic acids from, both genomic and transcript-derived nucleic acid samples. When free in solution, such probes are typically, but not invariably, detectably labeled; bound to a substrate, as in a microarray, such probes are typically, but not invariably unlabeled.

20 In one embodiment, the isolated nucleic acids of the present invention can be used as probes to detect and characterize gross alterations in the gene of a BSNA, such as deletions, insertions, translocations, and duplications of the BSNA genomic locus through fluorescence *in situ* hybridization (FISH) to chromosome spreads. *See, e.g.*, Andreeff *et al.* (eds.), Introduction to Fluorescence *In Situ* Hybridization: Principles and

25 Clinical Applications, John Wiley & Sons (1999), the disclosure of which is incorporated herein by reference in its entirety. The isolated nucleic acids of the present invention can be used as probes to assess smaller genomic alterations using, *e.g.*, Southern blot detection of restriction fragment length polymorphisms. The isolated nucleic acid molecules of the present invention can be used as probes to isolate genomic clones that

30 include the nucleic acid molecules of the present invention, which thereafter can be restriction mapped and sequenced to identify deletions, insertions, translocations, and substitutions (single nucleotide polymorphisms, SNPs) at the sequence level.

In another embodiment, the isolated nucleic acid molecules of the present invention can be used as probes to detect, characterize, and quantify BSNA in, and

isolate BSNA from, transcript-derived nucleic acid samples. In one aspect, the isolated nucleic acid molecules of the present invention can be used as hybridization probes to detect, characterize by length, and quantify mRNA by Northern blot of total or poly-A⁺-selected RNA samples. In another aspect, the isolated nucleic acid molecules of the present invention can be used as hybridization probes to detect, characterize by location, and quantify mRNA by *in situ* hybridization to tissue sections. *See, e.g., Schwarczacher et al., In Situ Hybridization*, Springer-Verlag New York (2000), the disclosure of which is incorporated herein by reference in its entirety. In another preferred embodiment, the isolated nucleic acid molecules of the present invention can be used as hybridization probes to measure the representation of clones in a cDNA library or to isolate hybridizing nucleic acid molecules acids from cDNA libraries, permitting sequence level characterization of mRNAs that hybridize to BSNAs, including, without limitations, identification of deletions, insertions, substitutions, truncations, alternatively spliced forms and single nucleotide polymorphisms. In yet another preferred embodiment, the nucleic acid molecules of the instant invention may be used in microarrays.

All of the aforementioned probe techniques are well within the skill in the art, and are described at greater length in standard texts such as Sambrook (2001), *supra*; Ausubel (1999), *supra*; and Walker *et al.* (eds.), *The Nucleic Acids Protocols Handbook*, Humana Press (2000), the disclosures of which are incorporated herein by reference in their entirety.

Thus, in one embodiment, a nucleic acid molecule of the invention may be used as a probe or primer to identify or amplify a second nucleic acid molecule that selectively hybridizes to the nucleic acid molecule of the invention. In a preferred embodiment, the probe or primer is derived from a nucleic acid molecule encoding a BSP. In a more preferred embodiment, the probe or primer is derived from a nucleic acid molecule encoding a polypeptide having an amino acid sequence of SEQ ID NO: 172 through 295. In another preferred embodiment, the probe or primer is derived from a BSNA. In a more preferred embodiment, the probe or primer is derived from a nucleic acid molecule having a nucleotide sequence of SEQ ID NO: 1 through 171.

In general, a probe or primer is at least 10 nucleotides in length, more preferably at least 12, more preferably at least 14 and even more preferably at least 16 or 17 nucleotides in length. In an even more preferred embodiment, the probe or primer is at least 18 nucleotides in length, even more preferably at least 20 nucleotides and even more preferably at least 22 nucleotides in length. Primers and probes may also be longer

-40-

in length. For instance, a probe or primer may be 25 nucleotides in length, or may be 30, 40 or 50 nucleotides in length. Methods of performing nucleic acid hybridization using oligonucleotide probes are well-known in the art. *See, e.g.*, Sambrook *et al.*, 1989, *supra*, Chapter 11 and pp. 11.31-11.32 and 11.40-11.44, which describes radiolabeling of short probes, and pp. 11.45-11.53, which describe hybridization conditions for oligonucleotide probes, including specific conditions for probe hybridization (pp. 11.50-11.51).

Methods of performing primer-directed amplification are also well-known in the art. Methods for performing the polymerase chain reaction (PCR) are compiled, *inter alia*, in McPherson, PCR Basics: From Background to Bench, Springer Verlag (2000); Innis *et al.* (eds.), PCR Applications: Protocols for Functional Genomics, Academic Press (1999); Gelfand *et al.* (eds.), PCR Strategies, Academic Press (1998); Newton *et al.*, PCR, Springer-Verlag New York (1997); Burke (ed.), PCR: Essential Techniques, John Wiley & Son Ltd (1996); White (ed.), PCR Cloning Protocols: From Molecular Cloning to Genetic Engineering, Vol. 67, Humana Press (1996); McPherson *et al.* (eds.), PCR 2: A Practical Approach, Oxford University Press, Inc. (1995); the disclosures of which are incorporated herein by reference in their entireties. Methods for performing RT-PCR are collected, *e.g.*, in Siebert *et al.* (eds.), Gene Cloning and Analysis by RT-PCR, Eaton Publishing Company/Bio Techniques Books Division, 1998; Siebert (ed.), PCR Technique: RT-PCR, Eaton Publishing Company/ BioTechniques Books (1995); the disclosure of which is incorporated herein by reference in its entirety.

PCR and hybridization methods may be used to identify and/or isolate allelic variants, homologous nucleic acid molecules and fragments of the nucleic acid molecules of the invention. PCR and hybridization methods may also be used to identify, amplify and/or isolate nucleic acid molecules that encode homologous proteins, analogs, fusion protein or muteins of the invention. The nucleic acid primers of the present invention can be used to prime amplification of nucleic acid molecules of the invention, using transcript-derived or genomic DNA as template.

The nucleic acid primers of the present invention can also be used, for example, to prime single base extension (SBE) for SNP detection (*See, e.g.*, U.S. Patent 6,004,744, the disclosure of which is incorporated herein by reference in its entirety).

Isothermal amplification approaches, such as rolling circle amplification, are also now well-described. *See, e.g.*, Schweitzer *et al.*, *Curr. Opin. Biotechnol.* 12(1): 21-7 (2001); U.S. Patents 5,854,033 and 5,714,320; and international patent publications WO 97/19193 and WO 00/15779, the disclosures of which are incorporated herein by

reference in their entireties. Rolling circle amplification can be combined with other techniques to facilitate SNP detection. *See, e.g., Lizardi et al., Nature Genet.* 19(3): 225-32 (1998).

Nucleic acid molecules of the present invention may be bound to a substrate
5 either covalently or noncovalently. The substrate can be porous or solid, planar or non-planar, unitary or distributed. The bound nucleic acid molecules may be used as hybridization probes, and may be labeled or unlabeled. In a preferred embodiment, the bound nucleic acid molecules are unlabeled.

In one embodiment, the nucleic acid molecule of the present invention is bound to
10 a porous substrate, *e.g.*, a membrane, typically comprising nitrocellulose, nylon, or positively-charged derivatized nylon. The nucleic acid molecule of the present invention can be used to detect a hybridizing nucleic acid molecule that is present within a labeled nucleic acid sample, *e.g.*, a sample of transcript-derived nucleic acids. In another embodiment, the nucleic acid molecule is bound to a solid substrate, including, without
15 limitation, glass, amorphous silicon, crystalline silicon or plastics. Examples of plastics include, without limitation, polymethylacrylic, polyethylene, polypropylene, polyacrylate, polymethylmethacrylate, polyvinylchloride, polytetrafluoroethylene, polystyrene, polycarbonate, polyacetal, polysulfone, celluloseacetate, cellulosenitrate, nitrocellulose, or mixtures thereof. The solid substrate may be any shape, including
20 rectangular, disk-like and spherical. In a preferred embodiment, the solid substrate is a microscope slide or slide-shaped substrate.

The nucleic acid molecule of the present invention can be attached covalently to a surface of the support substrate or applied to a derivatized surface in a chaotropic agent that facilitates denaturation and adherence by presumed noncovalent interactions, or
25 some combination thereof. The nucleic acid molecule of the present invention can be bound to a substrate to which a plurality of other nucleic acids are concurrently bound, hybridization to each of the plurality of bound nucleic acids being separately detectable. At low density, *e.g.* on a porous membrane, these substrate-bound collections are typically denominated macroarrays; at higher density, typically on a solid support, such
30 as glass, these substrate bound collections of plural nucleic acids are colloquially termed microarrays. As used herein, the term microarray includes arrays of all densities. It is, therefore, another aspect of the invention to provide microarrays that include the nucleic acids of the present invention.

Expression Vectors, Host Cells and Recombinant Methods of Producing Polypeptides

Another aspect of the present invention relates to vectors that comprise one or more of the isolated nucleic acid molecules of the present invention, and host cells in which such vectors have been introduced.

5 The vectors can be used, *inter alia*, for propagating the nucleic acids of the present invention in host cells (cloning vectors), for shuttling the nucleic acids of the present invention between host cells derived from disparate organisms (shuttle vectors), for inserting the nucleic acids of the present invention into host cell chromosomes (insertion vectors), for expressing sense or antisense RNA transcripts of the nucleic acids
10 of the present invention *in vitro* or within a host cell, and for expressing polypeptides encoded by the nucleic acids of the present invention, alone or as fusions to heterologous polypeptides (expression vectors). Vectors of the present invention will often be suitable for several such uses.

 Vectors are by now well-known in the art, and are described, *inter alia*, in Jones
15 *et al.* (eds.), Vectors: Cloning Applications: Essential Techniques (Essential Techniques Series), John Wiley & Son Ltd. (1998); Jones *et al.* (eds.), Vectors: Expression Systems: Essential Techniques (Essential Techniques Series), John Wiley & Son Ltd. (1998); Gacesa *et al.*, Vectors: Essential Data, John Wiley & Sons Ltd. (1995); Cid-Arregui (eds.), Viral Vectors: Basic Science and Gene Therapy, Eaton Publishing Co. (2000);
20 Sambrook (2001), *supra*; Ausubel (1999), *supra*; the disclosures of which are incorporated herein by reference in their entireties. Furthermore, an enormous variety of vectors are available commercially. Use of existing vectors and modifications thereof being well within the skill in the art, only basic features need be described here.

 Nucleic acid sequences may be expressed by operatively linking them to an
25 expression control sequence in an appropriate expression vector and employing that expression vector to transform an appropriate unicellular host. Expression control sequences are sequences which control the transcription, post-transcriptional events and translation of nucleic acid sequences. Such operative linking of a nucleic sequence of this invention to an expression control sequence, of course, includes, if not already part
30 of the nucleic acid sequence, the provision of a translation initiation codon, ATG or GTG, in the correct reading frame upstream of the nucleic acid sequence.

A wide variety of host/expression vector combinations may be employed in expressing the nucleic acid sequences of this invention. Useful expression vectors, for

example, may consist of segments of chromosomal, non-chromosomal and synthetic nucleic acid sequences.

In one embodiment, prokaryotic cells may be used with an appropriate vector. Prokaryotic host cells are often used for cloning and expression. In a preferred
5 embodiment, prokaryotic host cells include *E. coli*, *Pseudomonas*, *Bacillus* and *Streptomyces*. In a preferred embodiment, bacterial host cells are used to express the nucleic acid molecules of the instant invention. Useful expression vectors for bacterial hosts include bacterial plasmids, such as those from *E. coli*, *Bacillus* or *Streptomyces*, including pBluescript, pGEX-2T, pUC vectors, col E1, pCR1, pBR322, pMB9 and their
10 derivatives, wider host range plasmids, such as RP4, phage DNAs, *e.g.*, the numerous derivatives of phage lambda, *e.g.*, NM989, λ GT10 and λ GT11, and other phages, *e.g.*, M13 and filamentous single-stranded phage DNA. Where *E. coli* is used as host, selectable markers are, analogously, chosen for selectivity in gram negative bacteria: *e.g.*, typical markers confer resistance to antibiotics, such as ampicillin, tetracycline,
15 chloramphenicol, kanamycin, streptomycin and zeocin; auxotrophic markers can also be used.

In other embodiments, eukaryotic host cells, such as yeast, insect, mammalian or plant cells, may be used. Yeast cells, typically *S. cerevisiae*, are useful for eukaryotic genetic studies, due to the ease of targeting genetic changes by homologous
20 recombination and the ability to easily complement genetic defects using recombinantly expressed proteins. Yeast cells are useful for identifying interacting protein components, *e.g.* through use of a two-hybrid system. In a preferred embodiment, yeast cells are useful for protein expression. Vectors of the present invention for use in yeast will typically, but not invariably, contain an origin of replication suitable for use in yeast and
25 a selectable marker that is functional in yeast. Yeast vectors include Yeast Integrating plasmids (*e.g.*, YIp5) and Yeast Replicating plasmids (the YRp and YEp series plasmids), Yeast Centromere plasmids (the YCp series plasmids), Yeast Artificial Chromosomes (YACs) which are based on yeast linear plasmids, denoted YLp, pGPD-2, 2μ plasmids and derivatives thereof, and improved shuttle vectors such as those
30 described in Gietz *et al.*, *Gene*, 74: 527-34 (1988) (YIplac, YEplac and YCplac). Selectable markers in yeast vectors include a variety of auxotrophic markers, the most common of which are (in *Saccharomyces cerevisiae*) URA3, HIS3, LEU2, TRP1 and LYS2, which complement specific auxotrophic mutations, such as *ura3-52*, *his3-D1*, *leu2-D1*, *trp1-D1* and *lys2-201*.

-44-

Insect cells are often chosen for high efficiency protein expression. Where the host cells are from *Spodoptera frugiperda*, e.g., Sf9 and Sf21 cell lines, and expresSF™ cells (Protein Sciences Corp., Meriden, CT, USA)), the vector replicative strategy is typically based upon the baculovirus life cycle. Typically, baculovirus transfer vectors
5 are used to replace the wild-type AcMNPV polyhedrin gene with a heterologous gene of interest. Sequences that flank the polyhedrin gene in the wild-type genome are positioned 5' and 3' of the expression cassette on the transfer vectors. Following co-transfection with AcMNPV DNA, a homologous recombination event occurs between these sequences resulting in a recombinant virus carrying the gene of interest and the
10 polyhedrin or p10 promoter. Selection can be based upon visual screening for lacZ fusion activity.

In another embodiment, the host cells may be mammalian cells, which are particularly useful for expression of proteins intended as pharmaceutical agents, and for screening of potential agonists and antagonists of a protein or a physiological pathway.
15 Mammalian vectors intended for autonomous extrachromosomal replication will typically include a viral origin, such as the SV40 origin (for replication in cell lines expressing the large T-antigen, such as COS1 and COS7 cells), the papillomavirus origin, or the EBV origin for long term episomal replication (for use, e.g., in 293-EBNA cells, which constitutively express the EBV EBNA-1 gene product and adenovirus E1A).
20 Vectors intended for integration, and thus replication as part of the mammalian chromosome, can, but need not, include an origin of replication functional in mammalian cells, such as the SV40 origin. Vectors based upon viruses, such as adenovirus, adeno-associated virus, vaccinia virus, and various mammalian retroviruses, will typically replicate according to the viral replicative strategy. Selectable markers for use in
25 mammalian cells include resistance to neomycin (G418), blasticidin, hygromycin and to zeocin, and selection based upon the purine salvage pathway using HAT medium.

Expression in mammalian cells can be achieved using a variety of plasmids, including pSV2, pBC12BI, and p91023, as well as lytic virus vectors (e.g., vaccinia virus, adeno virus, and baculovirus), episomal virus vectors (e.g., bovine papillomavirus),
30 and retroviral vectors (e.g., murine retroviruses). Useful vectors for insect cells include baculoviral vectors and pVL 941.

Plant cells can also be used for expression, with the vector replicon typically derived from a plant virus (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) and selectable markers chosen for suitability in plants.

-45-

It is known that codon usage of different host cells may be different. For example, a plant cell and a human cell may exhibit a difference in codon preference for encoding a particular amino acid. As a result, human mRNA may not be efficiently translated in a plant, bacteria or insect host cell. Therefore, another embodiment of this invention is directed to codon optimization. The codons of the nucleic acid molecules of the invention may be modified to resemble, as much as possible, genes naturally contained within the host cell without altering the amino acid sequence encoded by the nucleic acid molecule.

Any of a wide variety of expression control sequences may be used in these vectors to express the DNA sequences of this invention. Such useful expression control sequences include the expression control sequences associated with structural genes of the foregoing expression vectors. Expression control sequences that control transcription include, *e.g.*, promoters, enhancers and transcription termination sites. Expression control sequences in eukaryotic cells that control post-transcriptional events include splice donor and acceptor sites and sequences that modify the half-life of the transcribed RNA, *e.g.*, sequences that direct poly(A) addition or binding sites for RNA-binding proteins. Expression control sequences that control translation include ribosome binding sites, sequences which direct targeted expression of the polypeptide to or within particular cellular compartments, and sequences in the 5' and 3' untranslated regions that modify the rate or efficiency of translation.

Examples of useful expression control sequences for a prokaryote, *e.g.*, *E. coli*, will include a promoter, often a phage promoter, such as phage lambda pL promoter, the *trc* promoter, a hybrid derived from the *trp* and *lac* promoters, the bacteriophage T7 promoter (in *E. coli* cells engineered to express the T7 polymerase), the TAC or TRC system, the major operator and promoter regions of phage lambda, the control regions of fd coat protein, or the *araBAD* operon. Prokaryotic expression vectors may further include transcription terminators, such as the *aspA* terminator, and elements that facilitate translation, such as a consensus ribosome binding site and translation termination codon, Schomer *et al.*, *Proc. Natl. Acad. Sci. USA* 83: 8506-8510 (1986).

Expression control sequences for yeast cells, typically *S. cerevisiae*, will include a yeast promoter, such as the *CYC1* promoter, the *GAL1* promoter, the *GAL10* promoter, *ADH1* promoter, the promoters of the yeast α -mating system, or the *GPD* promoter, and will typically have elements that facilitate transcription termination, such as the transcription termination signals from the *CYC1* or *ADH1* gene.

-46-

Expression vectors useful for expressing proteins in mammalian cells will include a promoter active in mammalian cells. These promoters include those derived from mammalian viruses, such as the enhancer-promoter sequences from the immediate early gene of the human cytomegalovirus (CMV), the enhancer-promoter sequences from the Rous sarcoma virus long terminal repeat (RSV LTR), the enhancer-promoter from SV40 or the early and late promoters of adenovirus. Other expression control sequences include the promoter for 3-phosphoglycerate kinase or other glycolytic enzymes, the promoters of acid phosphatase. Other expression control sequences include those from the gene comprising the BSNA of interest. Often, expression is enhanced by incorporation of polyadenylation sites, such as the late SV40 polyadenylation site and the polyadenylation signal and transcription termination sequences from the bovine growth hormone (BGH) gene, and ribosome binding sites. Furthermore, vectors can include introns, such as intron II of rabbit β -globin gene and the SV40 splice elements.

Preferred nucleic acid vectors also include a selectable or amplifiable marker gene and means for amplifying the copy number of the gene of interest. Such marker genes are well-known in the art. Nucleic acid vectors may also comprise stabilizing sequences (*e.g.*, ori- or ARS-like sequences and telomere-like sequences), or may alternatively be designed to favor directed or non-directed integration into the host cell genome. In a preferred embodiment, nucleic acid sequences of this invention are inserted in frame into an expression vector that allows high level expression of an RNA which encodes a protein comprising the encoded nucleic acid sequence of interest. Nucleic acid cloning and sequencing methods are well-known to those of skill in the art and are described in an assortment of laboratory manuals, including Sambrook (1989), *supra*, Sambrook (2000), *supra*; and Ausubel (1992), *supra*, Ausubel (1999), *supra*. Product information from manufacturers of biological, chemical and immunological reagents also provide useful information.

Expression vectors may be either constitutive or inducible. Inducible vectors include either naturally inducible promoters, such as the *trc* promoter, which is regulated by the *lac* operon, and the *pL* promoter, which is regulated by tryptophan, the MMTV-LTR promoter, which is inducible by dexamethasone, or can contain synthetic promoters and/or additional elements that confer inducible control on adjacent promoters. Examples of inducible synthetic promoters are the hybrid *Plac/ara-1* promoter and the *PLtetO-1* promoter. The *PLtetO-1* promoter takes advantage of the high expression levels from the *PL* promoter of phage lambda, but replaces the lambda repressor sites with two

-47-

copies of operator 2 of the Tn10 tetracycline resistance operon, causing this promoter to be tightly repressed by the Tet repressor protein and induced in response to tetracycline (Tc) and Tc derivatives such as anhydrotetracycline. Vectors may also be inducible because they contain hormone response elements, such as the glucocorticoid response
5 element (GRE) and the estrogen response element (ERE), which can confer hormone inducibility where vectors are used for expression in cells having the respective hormone receptors. To reduce background levels of expression, elements responsive to ecdysone, an insect hormone, can be used instead, with coexpression of the ecdysone receptor.

In one aspect of the invention, expression vectors can be designed to fuse the
10 expressed polypeptide to small protein tags that facilitate purification and/or visualization. Tags that facilitate purification include a polyhistidine tag that facilitates purification of the fusion protein by immobilized metal affinity chromatography, for example using NiNTA resin (Qiagen Inc., Valencia, CA, USA) or TALON™ resin (cobalt immobilized affinity chromatography medium, Clontech Labs, Palo Alto, CA,
15 USA). The fusion protein can include a chitin-binding tag and self-excising intein, permitting chitin-based purification with self-removal of the fused tag (IMPACT™ system, New England Biolabs, Inc., Beverley, MA, USA). Alternatively, the fusion protein can include a calmodulin-binding peptide tag, permitting purification by calmodulin affinity resin (Stratagene, La Jolla, CA, USA), or a specifically excisable
20 fragment of the biotin carboxylase carrier protein, permitting purification of *in vivo* biotinylated protein using an avidin resin and subsequent tag removal (Promega, Madison, WI, USA). As another useful alternative, the proteins of the present invention can be expressed as a fusion protein with glutathione-S-transferase, the affinity and specificity of binding to glutathione permitting purification using glutathione affinity
25 resins, such as Glutathione-Superflow Resin (Clontech Laboratories, Palo Alto, CA, USA), with subsequent elution with free glutathione. Other tags include, for example, the Xpress epitope, detectable by anti-Xpress antibody (Invitrogen, Carlsbad, CA, USA), a myc tag, detectable by anti-myc tag antibody, the V5 epitope, detectable by anti-V5 antibody (Invitrogen, Carlsbad, CA, USA), FLAG® epitope, detectable by anti-FLAG®
30 antibody (Stratagene, La Jolla, CA, USA), and the HA epitope.

For secretion of expressed proteins, vectors can include appropriate sequences that encode secretion signals, such as leader peptides. For example, the pSec Tag2 vectors (Invitrogen, Carlsbad, CA, USA) are 5.2 kb mammalian expression vectors that

-48-

carry the secretion signal from the V-J2-C region of the mouse Ig kappa-chain for efficient secretion of recombinant proteins from a variety of mammalian cell lines.

Expression vectors can also be designed to fuse proteins encoded by the heterologous nucleic acid insert to polypeptides that are larger than purification and/or
5 identification tags. Useful fusion proteins include those that permit display of the encoded protein on the surface of a phage or cell, fusion to intrinsically fluorescent proteins, such as those that have a green fluorescent protein (GFP)-like chromophore, fusions to the IgG Fc region, and fusion proteins for use in two hybrid systems.

Vectors for phage display fuse the encoded polypeptide to, e.g., the gene III
10 protein (pIII) or gene VIII protein (pVIII) for display on the surface of filamentous phage, such as M13. See Barbas *et al.*, Phage Display: A Laboratory Manual, Cold Spring Harbor Laboratory Press (2001); Kay *et al.* (eds.), Phage Display of Peptides and Proteins: A Laboratory Manual, Academic Press, Inc., (1996); Abelson *et al.* (eds.), Combinatorial Chemistry (Methods in Enzymology, Vol. 267) Academic Press (1996).
15 Vectors for yeast display, e.g. the pYD1 yeast display vector (Invitrogen, Carlsbad, CA, USA), use the α -agglutinin yeast adhesion receptor to display recombinant protein on the surface of *S. cerevisiae*. Vectors for mammalian display, e.g., the pDisplay™ vector (Invitrogen, Carlsbad, CA, USA), target recombinant proteins using an N-terminal cell surface targeting signal and a C-terminal transmembrane anchoring domain of platelet
20 derived growth factor receptor.

A wide variety of vectors now exist that fuse proteins encoded by heterologous nucleic acids to the chromophore of the substrate-independent, intrinsically fluorescent green fluorescent protein from *Aequorea victoria* ("GFP") and its variants. The GFP-like chromophore can be selected from GFP-like chromophores found in naturally occurring
25 proteins, such as *A. victoria* GFP (GenBank accession number AAA27721), *Renilla reniformis* GFP, FP583 (GenBank accession no. AF168419) (DsRed), FP593 (AF272711), FP483 (AF168420), FP484 (AF168424), FP595 (AF246709), FP486 (AF168421), FP538 (AF168423), and FP506 (AF168422), and need include only so much of the native protein as is needed to retain the chromophore's intrinsic
30 fluorescence. Methods for determining the minimal domain required for fluorescence are known in the art. See Li *et al.*, *J. Biol. Chem.* 272: 28545-28549 (1997). Alternatively, the GFP-like chromophore can be selected from GFP-like chromophores modified from those found in nature. The methods for engineering such modified GFP-like chromophores and testing them for fluorescence activity, both alone and as part of

protein fusions, are well-known in the art. See Heim *et al.*, *Curr. Biol.* 6: 178-182 (1996) and Palm *et al.*, *Methods Enzymol.* 302: 378-394 (1999), incorporated herein by reference in its entirety. A variety of such modified chromophores are now commercially available and can readily be used in the fusion proteins of the present invention. These include EGFP ("enhanced GFP"), EBFP ("enhanced blue fluorescent protein"), BFP2, EYFP ("enhanced yellow fluorescent protein"), ECFP ("enhanced cyan fluorescent protein") or Citrine. EGFP (*see, e.g.*, Cormack *et al.*, *Gene* 173: 33-38 (1996); United States Patent Nos. 6,090,919 and 5,804,387) is found on a variety of vectors, both plasmid and viral, which are available commercially (Clontech Labs, Palo Alto, CA, USA); EBFP is optimized for expression in mammalian cells whereas BFP2, which retains the original jellyfish codons, can be expressed in bacteria (*see, e.g.*, Heim *et al.*, *Curr. Biol.* 6: 178-182 (1996) and Cormack *et al.*, *Gene* 173: 33-38 (1996)). Vectors containing these blue-shifted variants are available from Clontech Labs (Palo Alto, CA, USA). Vectors containing EYFP, ECFP (*see, e.g.*, Heim *et al.*, *Curr. Biol.* 6: 178-182 (1996); Miyawaki *et al.*, *Nature* 388: 882-887 (1997)) and Citrine (*see, e.g.*, Heikal *et al.*, *Proc. Natl. Acad. Sci. USA* 97: 11996-12001 (2000)) are also available from Clontech Labs. The GFP-like chromophore can also be drawn from other modified GFPs, including those described in U.S. Patents 6,124,128; 6,096,865; 6,090,919; 6,066,476; 6,054,321; 6,027,881; 5,968,750; 5,874,304; 5,804,387; 5,777,079; 5,741,668; and 5,625,048, the disclosures of which are incorporated herein by reference in their entireties. See also Conn (ed.), Green Fluorescent Protein (Methods in Enzymology, Vol. 302), Academic Press, Inc. (1999). The GFP-like chromophore of each of these GFP variants can usefully be included in the fusion proteins of the present invention.

Fusions to the IgG Fc region increase serum half life of protein pharmaceutical products through interaction with the FcRn receptor (also denominated the FcRp receptor and the Brambell receptor, FcRb), further described in International Patent Application Nos. WO 97/43316, WO 97/34631, WO 96/32478, WO 96/18412.

For long-term, high-yield recombinant production of the proteins, protein fusions, and protein fragments of the present invention, stable expression is preferred. Stable expression is readily achieved by integration into the host cell genome of vectors having selectable markers, followed by selection of these integrants. Vectors such as pUB6/V5-His A, B, and C (Invitrogen, Carlsbad, CA, USA) are designed for high-level stable expression of heterologous proteins in a wide range of mammalian tissue types and

-50-

cell lines. pUB6/V5-His uses the promoter/enhancer sequence from the human ubiquitin C gene to drive expression of recombinant proteins: expression levels in 293, CHO, and NIH3T3 cells are comparable to levels from the CMV and human EF-1a promoters. The *bsd* gene permits rapid selection of stably transfected mammalian cells with the potent
5 antibiotic blasticidin.

Replication incompetent retroviral vectors, typically derived from Moloney murine leukemia virus, also are useful for creating stable transfectants having integrated provirus. The highly efficient transduction machinery of retroviruses, coupled with the availability of a variety of packaging cell lines such as RetroPack™ PT 67, EcoPack2™
10 293, AmphoPack-293, and GP2-293 cell lines (all available from Clontech Laboratories, Palo Alto, CA, USA), allow a wide host range to be infected with high efficiency; varying the multiplicity of infection readily adjusts the copy number of the integrated provirus.

Of course, not all vectors and expression control sequences will function equally
15 well to express the nucleic acid sequences of this invention. Neither will all hosts function equally well with the same expression system. However, one of skill in the art may make a selection among these vectors, expression control sequences and hosts without undue experimentation and without departing from the scope of this invention. For example, in selecting a vector, the host must be considered because the vector must
20 be replicated in it. The vector's copy number, the ability to control that copy number, the ability to control integration, if any, and the expression of any other proteins encoded by the vector, such as antibiotic or other selection markers, should also be considered. The present invention further includes host cells comprising the vectors of the present invention, either present episomally within the cell or integrated, in whole or in part, into
25 the host cell chromosome. Among other considerations, some of which are described above, a host cell strain may be chosen for its ability to process the expressed protein in the desired fashion. Such post-translational modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation, and it is an aspect of the present invention to provide BSPs with such post-
30 translational modifications.

Polypeptides of the invention may be post-translationally modified. Post-translational modifications include phosphorylation of amino acid residues serine, threonine and/or tyrosine, N-linked and/or O-linked glycosylation, methylation, acetylation, prenylation, methylation, acetylation, arginylation, ubiquination and

-51-

racemization. One may determine whether a polypeptide of the invention is likely to be post-translationally modified by analyzing the sequence of the polypeptide to determine if there are peptide motifs indicative of sites for post-translational modification. There are a number of computer programs that permit prediction of post-translational
5 modifications. See, e.g., www.expasy.org (accessed August 31, 2001), which includes PSORT, for prediction of protein sorting signals and localization sites, SignalP, for prediction of signal peptide cleavage sites, MITOPROT and Predotar, for prediction of mitochondrial targeting sequences, NetOGlyc, for prediction of type O-glycosylation sites in mammalian proteins, big-PI Predictor and DGPI, for prediction of prenylation-
10 anchor and cleavage sites, and NetPhos, for prediction of Ser, Thr and Tyr phosphorylation sites in eukaryotic proteins. Other computer programs, such as those included in GCG, also may be used to determine post-translational modification peptide motifs.

General examples of types of post-translational modifications may be found in
15 web sites such as the Delta Mass database <http://www.abrf.org/ABRF/ResearchCommittees/deltamass/deltamass.html> (accessed October 19, 2001); "GlycoSuiteDB: a new curated relational database of glycoprotein glycan structures and their biological sources" Cooper et al. *Nucleic Acids Res.* 29; 332-335 (2001) and <http://www.glycosuite.com/> (accessed October 19, 2001); "O-GLYCBASE version 4.0: a
20 revised database of O-glycosylated proteins" Gupta et al. *Nucleic Acids Research*, 27: 370-372 (1999) and <http://www.cbs.dtu.dk/databases/OGLYCBASE/> (accessed October 19, 2001); "PhosphoBase, a database of phosphorylation sites: release 2.0.", Kreegipuu et al. *Nucleic Acids Res* 27(1):237-239 (1999) and <http://www.cbs.dtu.dk/databases/PhosphoBase/> (accessed October 19, 2001); or <http://pir.georgetown.edu/pirwww/search/textresid.html>
25 <http://pirwww/search/textresid.html> (accessed October 19, 2001).

Tumorigenesis is often accompanied by alterations in the post-translational modifications of proteins. Thus, in another embodiment, the invention provides polypeptides from cancerous cells or tissues that have altered post-translational modifications compared to the post-translational modifications of polypeptides from
30 normal cells or tissues. A number of altered post-translational modifications are known. One common alteration is a change in phosphorylation state, wherein the polypeptide from the cancerous cell or tissue is hyperphosphorylated or hypophosphorylated compared to the polypeptide from a normal tissue, or wherein the polypeptide is phosphorylated on different residues than the polypeptide from a normal cell. Another

-52-

common alteration is a change in glycosylation state, wherein the polypeptide from the cancerous cell or tissue has more or less glycosylation than the polypeptide from a normal tissue, and/or wherein the polypeptide from the cancerous cell or tissue has a different type of glycosylation than the polypeptide from a noncancerous cell or tissue.

- 5 Changes in glycosylation may be critical because carbohydrate-protein and carbohydrate-carbohydrate interactions are important in cancer cell progression, dissemination and invasion. See, e.g., Barchi, *Curr. Pharm. Des.* 6: 485-501 (2000), Verma, *Cancer Biochem. Biophys.* 14: 151-162 (1994) and Dennis et al., *Bioessays* 5: 412-421 (1999).

- Another post-translational modification that may be altered in cancer cells is
10 prenylation. Prenylation is the covalent attachment of a hydrophobic prenyl group (either farnesyl or geranylgeranyl) to a polypeptide. Prenylation is required for localizing a protein to a cell membrane and is often required for polypeptide function. For instance, the Ras superfamily of GTPase signaling proteins must be prenylated for function in a cell. See, e.g., Prendergast et al., *Semin. Cancer Biol.* 10: 443-452 (2000) and Khwaja et al., *Lancet* 355: 741-744 (2000).
15

- Other post-translation modifications that may be altered in cancer cells include, without limitation, polypeptide methylation, acetylation, arginylation or racemization of amino acid residues. In these cases, the polypeptide from the cancerous cell may exhibit either increased or decreased amounts of the post-translational modification compared to
20 the corresponding polypeptides from noncancerous cells.

- Other polypeptide alterations in cancer cells include abnormal polypeptide cleavage of proteins and aberrant protein-protein interactions. Abnormal polypeptide cleavage may be cleavage of a polypeptide in a cancerous cell that does not usually occur in a normal cell, or a lack of cleavage in a cancerous cell, wherein the polypeptide is
25 cleaved in a normal cell. Aberrant protein-protein interactions may be either covalent cross-linking or non-covalent binding between proteins that do not normally bind to each other. Alternatively, in a cancerous cell, a protein may fail to bind to another protein to which it is bound in a noncancerous cell. Alterations in cleavage or in protein-protein interactions may be due to over- or underproduction of a polypeptide in a cancerous cell
30 compared to that in a normal cell, or may be due to alterations in post-translational modifications (see above) of one or more proteins in the cancerous cell. See, e.g., Henschen-Edman, *Ann. N.Y. Acad. Sci.* 936: 580-593 (2001).

Alterations in polypeptide post-translational modifications, as well as changes in polypeptide cleavage and protein-protein interactions, may be determined by any method

-53-

known in the art. For instance, alterations in phosphorylation may be determined by using anti-phosphoserine, anti-phosphothreonine or anti-phosphotyrosine antibodies or by amino acid analysis. Glycosylation alterations may be determined using antibodies specific for different sugar residues, by carbohydrate sequencing, or by alterations in the size of the glycoprotein, which can be determined by, e.g., SDS polyacrylamide gel electrophoresis (PAGE). Other alterations of post-translational modifications, such as prenylation, racemization, methylation, acetylation and arginylation, may be determined by chemical analysis, protein sequencing, amino acid analysis, or by using antibodies specific for the particular post-translational modifications. Changes in protein-protein interactions and in polypeptide cleavage may be analyzed by any method known in the art including, without limitation, non-denaturing PAGE (for non-covalent protein-protein interactions), SDS PAGE (for covalent protein-protein interactions and protein cleavage), chemical cleavage, protein sequencing or immunoassays.

In another embodiment, the invention provides polypeptides that have been post-translationally modified. In one embodiment, polypeptides may be modified enzymatically or chemically, by addition or removal of a post-translational modification. For example, a polypeptide may be glycosylated or deglycosylated enzymatically. Similarly, polypeptides may be phosphorylated using a purified kinase, such as a MAP kinase (e.g., p38, ERK, or JNK) or a tyrosine kinase (e.g., Src or erbB2). A polypeptide may also be modified through synthetic chemistry. Alternatively, one may isolate the polypeptide of interest from a cell or tissue that expresses the polypeptide with the desired post-translational modification. In another embodiment, a nucleic acid molecule encoding the polypeptide of interest is introduced into a host cell that is capable of post-translationally modifying the encoded polypeptide in the desired fashion. If the polypeptide does not contain a motif for a desired post-translational modification, one may alter the post-translational modification by mutating the nucleic acid sequence of a nucleic acid molecule encoding the polypeptide so that it contains a site for the desired post-translational modification. Amino acid sequences that may be post-translationally modified are known in the art. See, e.g., the programs described above on the website www.expasy.org. The nucleic acid molecule is then be introduced into a host cell that is capable of post-translationally modifying the encoded polypeptide. Similarly, one may delete sites that are post-translationally modified by either mutating the nucleic acid sequence so that the encoded polypeptide does not contain the post-translational

modification motif, or by introducing the native nucleic acid molecule into a host cell that is not capable of post-translationally modifying the encoded polypeptide.

In selecting an expression control sequence, a variety of factors should also be considered. These include, for example, the relative strength of the sequence, its
5 controllability, and its compatibility with the nucleic acid sequence of this invention, particularly with regard to potential secondary structures. Unicellular hosts should be selected by consideration of their compatibility with the chosen vector, the toxicity of the product coded for by the nucleic acid sequences of this invention, their secretion characteristics, their ability to fold the polypeptide correctly, their fermentation or culture
10 requirements, and the ease of purification from them of the products coded for by the nucleic acid sequences of this invention.

The recombinant nucleic acid molecules and more particularly, the expression vectors of this invention may be used to express the polypeptides of this invention as recombinant polypeptides in a heterologous host cell. The polypeptides of this invention
15 may be full-length or less than full-length polypeptide fragments recombinantly expressed from the nucleic acid sequences according to this invention. Such polypeptides include analogs, derivatives and muteins that may or may not have biological activity.

Vectors of the present invention will also often include elements that permit *in*
20 *vitro* transcription of RNA from the inserted heterologous nucleic acid. Such vectors typically include a phage promoter, such as that from T7, T3, or SP6, flanking the nucleic acid insert. Often two different such promoters flank the inserted nucleic acid, permitting separate *in vitro* production of both sense and antisense strands.

Transformation and other methods of introducing nucleic acids into a host cell
25 (*e.g.*, conjugation, protoplast transformation or fusion, transfection, electroporation, liposome delivery, membrane fusion techniques, high velocity DNA-coated pellets, viral infection and protoplast fusion) can be accomplished by a variety of methods which are well-known in the art (*See*, for instance, Ausubel, *supra*, and Sambrook *et al.*, *supra*). Bacterial, yeast, plant or mammalian cells are transformed or transfected with an
30 expression vector, such as a plasmid, a cosmid, or the like, wherein the expression vector comprises the nucleic acid of interest. Alternatively, the cells may be infected by a viral expression vector comprising the nucleic acid of interest. Depending upon the host cell, vector, and method of transformation used, transient or stable expression of the polypeptide will be constitutive or inducible. One having ordinary skill in the art will be

-55-

able to decide whether to express a polypeptide transiently or stably, and whether to express the protein constitutively or inducibly.

A wide variety of unicellular host cells are useful in expressing the DNA sequences of this invention. These hosts may include well-known eukaryotic and prokaryotic hosts, such as strains of, fungi, yeast, insect cells such as *Spodoptera frugiperda* (SF9), animal cells such as CHO, as well as plant cells in tissue culture. Representative examples of appropriate host cells include, but are not limited to, bacterial cells, such as *E. coli*, *Caulobacter crescentus*, *Streptomyces* species, and *Salmonella typhimurium*; yeast cells, such as *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Pichia pastoris*, *Pichia methanolica*; insect cell lines, such as those from *Spodoptera frugiperda*, e.g., Sf9 and Sf21 cell lines, and expresSF™ cells (Protein Sciences Corp., Meriden, CT, USA), *Drosophila* S2 cells, and *Trichoplusia ni* High Five® Cells (Invitrogen, Carlsbad, CA, USA); and mammalian cells. Typical mammalian cells include BHK cells, BSC 1 cells, BSC 40 cells, BMT 10 cells, VERO cells, COS1 cells, COS7 cells, Chinese hamster ovary (CHO) cells, 3T3 cells, NIH 3T3 cells, 293 cells, HEPG2 cells, HeLa cells, L cells, MDCK cells, HEK293 cells, WI38 cells, murine ES cell lines (e.g., from strains 129/SV, C57/BL6, DBA-1, 129/SVJ), K562 cells, Jurkat cells, and BW5147 cells. Other mammalian cell lines are well-known and readily available from the American Type Culture Collection (ATCC) (Manassas, VA, USA) and the National Institute of General Medical Sciences (NIGMS) Human Genetic Cell Repository at the Coriell Cell Repositories (Camden, NJ, USA). Cells or cell lines derived from breast are particularly preferred because they may provide a more native post-translational processing. Particularly preferred are human breast cells.

Particular details of the transfection, expression and purification of recombinant proteins are well documented and are understood by those of skill in the art. Further details on the various technical aspects of each of the steps used in recombinant production of foreign genes in bacterial cell expression systems can be found in a number of texts and laboratory manuals in the art. See, e.g., Ausubel (1992), *supra*, Ausubel (1999), *supra*, Sambrook (1989), *supra*, and Sambrook (2001), *supra*, herein incorporated by reference.

Methods for introducing the vectors and nucleic acids of the present invention into the host cells are well-known in the art; the choice of technique will depend primarily upon the specific vector to be introduced and the host cell chosen.

-56-

Nucleic acid molecules and vectors may be introduced into prokaryotes, such as *E. coli*, in a number of ways. For instance, phage lambda vectors will typically be packaged using a packaging extract (e.g., Gigapack® packaging extract, Stratagene, La Jolla, CA, USA), and the packaged virus used to infect *E. coli*.

5 Plasmid vectors will typically be introduced into chemically competent or electrocompetent bacterial cells. *E. coli* cells can be rendered chemically competent by treatment, e.g., with CaCl₂, or a solution of Mg²⁺, Mn²⁺, Ca²⁺, Rb⁺ or K⁺, dimethyl sulfoxide, dithiothreitol, and hexamine cobalt (III), Hanahan, *J. Mol. Biol.* 166(4):557-80 (1983), and vectors introduced by heat shock. A wide variety of chemically competent
10 strains are also available commercially (e.g., Epicurian Coli® XL10-Gold® Ultracompetent Cells (Stratagene, La Jolla, CA, USA); DH5α competent cells (Clontech Laboratories, Palo Alto, CA, USA); and TOP10 Chemically Competent *E. coli* Kit (Invitrogen, Carlsbad, CA, USA)). Bacterial cells can be rendered electrocompetent, that is, competent to take up exogenous DNA by electroporation, by various pre-pulse
15 treatments; vectors are introduced by electroporation followed by subsequent outgrowth in selected media. An extensive series of protocols is provided online in Electroprotocols (BioRad, Richmond, CA, USA) (http://www.biorad.com/LifeScience/pdf/New_Gene_Pulser.pdf).

Vectors can be introduced into yeast cells by spheroplasting, treatment with
20 lithium salts, electroporation, or protoplast fusion. Spheroplasts are prepared by the action of hydrolytic enzymes such as snail-gut extract, usually denoted Glusulase, or Zymolyase, an enzyme from *Arthrobacter luteus*, to remove portions of the cell wall in the presence of osmotic stabilizers, typically 1 M sorbitol. DNA is added to the spheroplasts, and the mixture is co-precipitated with a solution of polyethylene glycol
25 (PEG) and Ca²⁺. Subsequently, the cells are resuspended in a solution of sorbitol, mixed with molten agar and then layered on the surface of a selective plate containing sorbitol.

For lithium-mediated transformation, yeast cells are treated with lithium acetate, which apparently permeabilizes the cell wall, DNA is added and the cells are co-precipitated with PEG. The cells are exposed to a brief heat shock, washed free of
30 PEG and lithium acetate, and subsequently spread on plates containing ordinary selective medium. Increased frequencies of transformation are obtained by using specially-prepared single-stranded carrier DNA and certain organic solvents. Schiestl *et al.*, *Curr. Genet.* 16(5-6): 339-46 (1989).

-57-

For electroporation, freshly-grown yeast cultures are typically washed, suspended in an osmotic protectant, such as sorbitol, mixed with DNA, and the cell suspension pulsed in an electroporation device. Subsequently, the cells are spread on the surface of plates containing selective media. Becker *et al.*, *Methods Enzymol.* 194: 182-187 (1991).

5 The efficiency of transformation by electroporation can be increased over 100-fold by using PEG, single-stranded carrier DNA and cells that are in late log-phase of growth. Larger constructs, such as YACs, can be introduced by protoplast fusion.

Mammalian and insect cells can be directly infected by packaged viral vectors, or transfected by chemical or electrical means. For chemical transfection, DNA can be
10 coprecipitated with CaPO₄ or introduced using liposomal and nonliposomal lipid-based agents. Commercial kits are available for CaPO₄ transfection (CalPhos™ Mammalian Transfection Kit, Clontech Laboratories, Palo Alto, CA, USA), and lipid-mediated transfection can be practiced using commercial reagents, such as LIPOFECTAMINE™
2000, LIPOFECTAMINE™ Reagent, CELLFECTIN® Reagent, and LIPOFECTIN®
15 Reagent (Invitrogen, Carlsbad, CA, USA), DOTAP Liposomal Transfection Reagent, FuGENE 6, X-tremeGENE Q2, DOSPER, (Roche Molecular Biochemicals, Indianapolis, IN USA), Effectene™, PolyFect®, Superfect® (Qiagen, Inc., Valencia, CA, USA). Protocols for electroporating mammalian cells can be found online in Electroprotocols (Bio-Rad, Richmond, CA, USA) ([http://www.bio-rad.com/LifeScience/pdf/](http://www.bio-rad.com/LifeScience/pdf/New_Gene_Pulser.pdf)
20 [New_Gene_Pulser.pdf](http://www.bio-rad.com/LifeScience/pdf/New_Gene_Pulser.pdf)); Norton *et al.* (eds.), Gene Transfer Methods: Introducing DNA into Living Cells and Organisms, BioTechniques Books, Eaton Publishing Co. (2000); incorporated herein by reference in its entirety. Other transfection techniques include transfection by particle bombardment and microinjection. See, e.g., Cheng *et al.*, *Proc. Natl. Acad. Sci. USA* 90(10): 4455-9 (1993); Yang *et al.*, *Proc. Natl. Acad. Sci. USA*
25 87(24): 9568-72 (1990).

Production of the recombinantly produced proteins of the present invention can optionally be followed by purification.

Purification of recombinantly expressed proteins is now well by those skilled in the art. See, e.g., Thorner *et al.* (eds.), Applications of Chimeric Genes and Hybrid
30 Proteins, Part A: Gene Expression and Protein Purification (Methods in Enzymology, Vol. 326), Academic Press (2000); Harbin (ed.), Cloning, Gene Expression and Protein Purification : Experimental Procedures and Process Rationale, Oxford Univ. Press (2001); Marshak *et al.*, Strategies for Protein Purification and Characterization: A Laboratory Course Manual, Cold Spring Harbor Laboratory Press (1996); and Roe (ed.),

Protein Purification Applications, Oxford University Press (2001); the disclosures of which are incorporated herein by reference in their entireties, and thus need not be detailed here.

Briefly, however, if purification tags have been fused through use of an expression vector that appends such tags, purification can be effected, at least in part, by means appropriate to the tag, such as use of immobilized metal affinity chromatography for polyhistidine tags. Other techniques common in the art include ammonium sulfate fractionation, immunoprecipitation, fast protein liquid chromatography (FPLC), high performance liquid chromatography (HPLC), and preparative gel electrophoresis.

10 Polypeptides

Another object of the invention is to provide polypeptides encoded by the nucleic acid molecules of the instant invention. In a preferred embodiment, the polypeptide is a breast specific polypeptide (BSP). In an even more preferred embodiment, the polypeptide is derived from a polypeptide comprising the amino acid sequence of SEQ ID NO: 172 through 295. A polypeptide as defined herein may be produced recombinantly, as discussed *supra*, may be isolated from a cell that naturally expresses the protein, or may be chemically synthesized following the teachings of the specification and using methods well-known to those having ordinary skill in the art.

In another aspect, the polypeptide may comprise a fragment of a polypeptide, wherein the fragment is as defined herein. In a preferred embodiment, the polypeptide fragment is a fragment of a BSP. In a more preferred embodiment, the fragment is derived from a polypeptide comprising the amino acid sequence of SEQ ID NO: 172 through 295. A polypeptide that comprises only a fragment of an entire BSP may or may not be a polypeptide that is also a BSP. For instance, a full-length polypeptide may be breast-specific, while a fragment thereof may be found in other tissues as well as in breast. A polypeptide that is not a BSP, whether it is a fragment, analog, mutein, homologous protein or derivative, is nevertheless useful, especially for immunizing animals to prepare anti-BSP antibodies. However, in a preferred embodiment, the part or fragment is a BSP. Methods of determining whether a polypeptide is a BSP are described *infra*.

Fragments of at least 6 contiguous amino acids are useful in mapping B cell and T cell epitopes of the reference protein. See, e.g., Geysen *et al.*, *Proc. Natl. Acad. Sci. USA* 81: 3998-4002 (1984) and U.S. Patents 4,708,871 and 5,595,915, the disclosures of

-59-

which are incorporated herein by reference in their entireties. Because the fragment need not itself be immunogenic, part of an immunodominant epitope, nor even recognized by native antibody, to be useful in such epitope mapping, all fragments of at least 6 amino acids of the proteins of the present invention have utility in such a study.

5 Fragments of at least 8 contiguous amino acids, often at least 15 contiguous amino acids, are useful as immunogens for raising antibodies that recognize the proteins of the present invention. *See, e.g.,* Lerner, *Nature* 299: 592-596 (1982); Shinnick *et al.*, *Annu. Rev. Microbiol.* 37: 425-46 (1983); Sutcliffe *et al.*, *Science* 219: 660-6 (1983), the disclosures of which are incorporated herein by reference in their entireties. As further
10 described in the above-cited references, virtually all 8-mers, conjugated to a carrier, such as a protein, prove immunogenic, meaning that they are capable of eliciting antibody for the conjugated peptide; accordingly, all fragments of at least 8 amino acids of the proteins of the present invention have utility as immunogens.

 Fragments of at least 8, 9, 10 or 12 contiguous amino acids are also useful as
15 competitive inhibitors of binding of the entire protein, or a portion thereof, to antibodies (as in epitope mapping), and to natural binding partners, such as subunits in a multimeric complex or to receptors or ligands of the subject protein; this competitive inhibition permits identification and separation of molecules that bind specifically to the protein of interest, U.S. Patents 5,539,084 and 5,783,674, incorporated herein by reference in their
20 entireties.

 The protein, or protein fragment, of the present invention is thus at least 6 amino acids in length, typically at least 8, 9, 10 or 12 amino acids in length, and often at least 15 amino acids in length. Often, the protein of the present invention, or fragment thereof, is at least 20 amino acids in length, even 25 amino acids, 30 amino acids, 35 amino acids,
25 or 50 amino acids or more in length. Of course, larger fragments having at least 75 amino acids, 100 amino acids, or even 150 amino acids are also useful, and at times preferred.

 One having ordinary skill in the art can produce fragments of a polypeptide by truncating the nucleic acid molecule, *e.g.*, a BSNA, encoding the polypeptide and then
30 expressing it recombinantly. Alternatively, one can produce a fragment by chemically synthesizing a portion of the full-length polypeptide. One may also produce a fragment by enzymatically cleaving either a recombinant polypeptide or an isolated naturally-occurring polypeptide. Methods of producing polypeptide fragments are well-known in the art. *See, e.g.,* Sambrook (1989), *supra*; Sambrook (2001), *supra*; Ausubel (1992),

-60-

supra; and Ausubel (1999), *supra*. In one embodiment, a polypeptide comprising only a fragment of polypeptide of the invention, preferably a BSP, may be produced by chemical or enzymatic cleavage of a polypeptide. In a preferred embodiment, a polypeptide fragment is produced by expressing a nucleic acid molecule encoding a
5 fragment of the polypeptide, preferably a BSP, in a host cell.

By "polypeptides" as used herein it is also meant to be inclusive of mutants, fusion proteins, homologous proteins and allelic variants of the polypeptides specifically exemplified.

A mutant protein, or mutein, may have the same or different properties compared
10 to a naturally-occurring polypeptide and comprises at least one amino acid insertion, duplication, deletion, rearrangement or substitution compared to the amino acid sequence of a native protein. Small deletions and insertions can often be found that do not alter the function of the protein. In one embodiment, the mutein may or may not be breast-specific. In a preferred embodiment, the mutein is breast-specific. In a preferred
15 embodiment, the mutein is a polypeptide that comprises at least one amino acid insertion, duplication, deletion, rearrangement or substitution compared to the amino acid sequence of SEQ ID NO: 172 through 295. In a more preferred embodiment, the mutein is one that exhibits at least 50% sequence identity, more preferably at least 60% sequence identity, even more preferably at least 70%, yet more preferably at least 80% sequence identity to
20 a BSP comprising an amino acid sequence of SEQ ID NO: 172 through 295. In yet a more preferred embodiment, the mutein exhibits at least 85%, more preferably 90%, even more preferably 95% or 96%, and yet more preferably at least 97%, 98%, 99% or 99.5% sequence identity to a BSP comprising an amino acid sequence of SEQ ID NO: 172 through 295.

25 A mutein may be produced by isolation from a naturally-occurring mutant cell, tissue or organism. A mutein may be produced by isolation from a cell, tissue or organism that has been experimentally mutagenized. Alternatively, a mutein may be produced by chemical manipulation of a polypeptide, such as by altering the amino acid residue to another amino acid residue using synthetic or semi-synthetic chemical
30 techniques. In a preferred embodiment, a mutein may be produced from a host cell comprising an altered nucleic acid molecule compared to the naturally-occurring nucleic acid molecule. For instance, one may produce a mutein of a polypeptide by introducing one or more mutations into a nucleic acid sequence of the invention and then expressing it recombinantly. These mutations may be targeted, in which particular encoded amino

-61-

acids are altered, or may be untargeted, in which random encoded amino acids within the polypeptide are altered. Muteins with random amino acid alterations can be screened for a particular biological activity or property, particularly whether the polypeptide is breast-specific, as described below. Multiple random mutations can be introduced into the

5 gene by methods well-known to the art, *e.g.*, by error-prone PCR, shuffling, oligonucleotide-directed mutagenesis, assembly PCR, sexual PCR mutagenesis, *in vivo* mutagenesis, cassette mutagenesis, recursive ensemble mutagenesis, exponential ensemble mutagenesis and site-specific mutagenesis. Methods of producing muteins with targeted or random amino acid alterations are well-known in the art. *See, e.g.*,

10 Sambrook (1989), *supra*; Sambrook (2001), *supra*; Ausubel (1992), *supra*; and Ausubel (1999), U.S. Patent 5,223,408, and the references discussed *supra*, each herein incorporated by reference.

By "polypeptide" as used herein it is also meant to be inclusive of polypeptides homologous to those polypeptides exemplified herein. In a preferred embodiment, the

15 polypeptide is homologous to a BSP. In an even more preferred embodiment, the polypeptide is homologous to a BSP selected from the group having an amino acid sequence of SEQ ID NO: 172 through 295. In a preferred embodiment, the homologous polypeptide is one that exhibits significant sequence identity to a BSP. In a more preferred embodiment, the polypeptide is one that exhibits significant sequence identity

20 to an comprising an amino acid sequence of SEQ ID NO: 172 through 295. In an even more preferred embodiment, the homologous polypeptide is one that exhibits at least 50% sequence identity, more preferably at least 60% sequence identity, even more preferably at least 70%, yet more preferably at least 80% sequence identity to a BSP comprising an amino acid sequence of SEQ ID NO: 172 through 295. In a yet more

25 preferred embodiment, the homologous polypeptide is one that exhibits at least 85%, more preferably 90%, even more preferably 95% or 96%, and yet more preferably at least 97% or 98% sequence identity to a BSP comprising an amino acid sequence of SEQ ID NO: 172 through 295. In another preferred embodiment, the homologous polypeptide is one that exhibits at least 99%, more preferably 99.5%, even more preferably 99.6%,

30 99.7%, 99.8% or 99.9% sequence identity to a BSP comprising an amino acid sequence of SEQ ID NO: 172 through 295. In a preferred embodiment, the amino acid substitutions are conservative amino acid substitutions as discussed above.

In another embodiment, the homologous polypeptide is one that is encoded by a nucleic acid molecule that selectively hybridizes to a BSNA. In a preferred embodiment,

-62-

the homologous polypeptide is encoded by a nucleic acid molecule that hybridizes to a BSNA under low stringency, moderate stringency or high stringency conditions, as defined herein. In a more preferred embodiment, the BSNA is selected from the group consisting of SEQ ID NO: 1 through 171. In another preferred embodiment, the

5 homologous polypeptide is encoded by a nucleic acid molecule that hybridizes to a nucleic acid molecule that encodes a BSP under low stringency, moderate stringency or high stringency conditions, as defined herein. In a more preferred embodiment, the BSP is selected from the group consisting of SEQ ID NO: 172 through 295.

The homologous polypeptide may be a naturally-occurring one that is derived

10 from another species, especially one derived from another primate, such as chimpanzee, gorilla, rhesus macaque, baboon or gorilla, wherein the homologous polypeptide comprises an amino acid sequence that exhibits significant sequence identity to that of SEQ ID NO: 172 through 295. The homologous polypeptide may also be a naturally-occurring polypeptide from a human, when the BSP is a member of a family of

15 polypeptides. The homologous polypeptide may also be a naturally-occurring polypeptide derived from a non-primate, mammalian species, including without limitation, domesticated species, *e.g.*, dog, cat, mouse, rat, rabbit, guinea pig, hamster, cow, horse, goat or pig. The homologous polypeptide may also be a naturally-occurring polypeptide derived from a non-mammalian species, such as birds or reptiles. The

20 naturally-occurring homologous protein may be isolated directly from humans or other species. Alternatively, the nucleic acid molecule encoding the naturally-occurring homologous polypeptide may be isolated and used to express the homologous polypeptide recombinantly. In another embodiment, the homologous polypeptide may be one that is experimentally produced by random mutation of a nucleic acid molecule and

25 subsequent expression of the nucleic acid molecule. In another embodiment, the homologous polypeptide may be one that is experimentally produced by directed mutation of one or more codons to alter the encoded amino acid of a BSP. Further, the homologous protein may or may not encode polypeptide that is a BSP. However, in a preferred embodiment, the homologous polypeptide encodes a polypeptide that is a BSP.

30 Relatedness of proteins can also be characterized using a second functional test, the ability of a first protein competitively to inhibit the binding of a second protein to an antibody. It is, therefore, another aspect of the present invention to provide isolated proteins not only identical in sequence to those described with particularity herein, but also to provide isolated proteins ("cross-reactive proteins") that competitively inhibit the

-63-

binding of antibodies to all or to a portion of various of the isolated polypeptides of the present invention. Such competitive inhibition can readily be determined using immunoassays well-known in the art.

As discussed above, single nucleotide polymorphisms (SNPs) occur frequently in eukaryotic genomes, and the sequence determined from one individual of a species may differ from other allelic forms present within the population. Thus, by "polypeptide" as used herein it is also meant to be inclusive of polypeptides encoded by an allelic variant of a nucleic acid molecule encoding a BSP. In a preferred embodiment, the polypeptide is encoded by an allelic variant of a gene that encodes a polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO: 172 through 295. In a yet more preferred embodiment, the polypeptide is encoded by an allelic variant of a gene that has the nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through 171.

In another embodiment, the invention provides polypeptides which comprise derivatives of a polypeptide encoded by a nucleic acid molecule according to the instant invention. In a preferred embodiment, the polypeptide is a BSP. In a preferred embodiment, the polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NO: 172 through 295, or is a mutein, allelic variant, homologous protein or fragment thereof. In a preferred embodiment, the derivative has been acetylated, carboxylated, phosphorylated, glycosylated or ubiquitinated. In another preferred embodiment, the derivative has been labeled with, *e.g.*, radioactive isotopes such as ^{125}I , ^{32}P , ^{35}S , and ^3H . In another preferred embodiment, the derivative has been labeled with fluorophores, chemiluminescent agents, enzymes, and antiligands that can serve as specific binding pair members for a labeled ligand.

Polypeptide modifications are well-known to those of skill and have been described in great detail in the scientific literature. Several particularly common modifications, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, for instance, are described in most basic texts, such as, for instance Creighton, Protein Structure and Molecular Properties, 2nd ed., W. H. Freeman and Company (1993). Many detailed reviews are available on this subject, such as, for example, those provided by Wold, in Johnson (ed.), Posttranslational Covalent Modification of Proteins, pgs. 1-12, Academic Press (1983); Seifter *et al.*, *Meth. Enzymol.* 182: 626-646 (1990) and Rattan *et al.*, *Ann. N.Y. Acad. Sci.* 663: 48-62 (1992).

-64-

It will be appreciated, as is well-known and as noted above, that polypeptides are not always entirely linear. For instance, polypeptides may be branched as a result of ubiquitination, and they may be circular, with or without branching, generally as a result of posttranslation events, including natural processing event and events brought about by human manipulation which do not occur naturally. Circular, branched and branched circular polypeptides may be synthesized by non-translation natural process and by entirely synthetic methods, as well. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. In fact, blockage of the amino or carboxyl group in a polypeptide, or both, by a covalent modification, is common in naturally occurring and synthetic polypeptides and such modifications may be present in polypeptides of the present invention, as well. For instance, the amino terminal residue of polypeptides made in *E. coli*, prior to proteolytic processing, almost invariably will be N-formylmethionine.

Useful post-synthetic (and post-translational) modifications include conjugation to detectable labels, such as fluorophores. A wide variety of amine-reactive and thiol-reactive fluorophore derivatives have been synthesized that react under nondenaturing conditions with N-terminal amino groups and epsilon amino groups of lysine residues, on the one hand, and with free thiol groups of cysteine residues, on the other.

Kits are available commercially that permit conjugation of proteins to a variety of amine-reactive or thiol-reactive fluorophores: Molecular Probes, Inc. (Eugene, OR, USA), *e.g.*, offers kits for conjugating proteins to Alexa Fluor 350, Alexa Fluor 430, Fluorescein-EX, Alexa Fluor 488, Oregon Green 488, Alexa Fluor 532, Alexa Fluor 546, Alexa Fluor 546, Alexa Fluor 568, Alexa Fluor 594, and Texas Red-X.

A wide variety of other amine-reactive and thiol-reactive fluorophores are available commercially (Molecular Probes, Inc., Eugene, OR, USA), including Alexa Fluor® 350, Alexa Fluor® 488, Alexa Fluor® 532, Alexa Fluor® 546, Alexa Fluor® 568, Alexa Fluor® 594, Alexa Fluor® 647 (monoclonal antibody labeling kits available from Molecular Probes, Inc., Eugene, OR, USA), BODIPY dyes, such as BODIPY 493/503, BODIPY FL, BODIPY R6G, BODIPY 530/550, BODIPY TMR, BODIPY 558/568, BODIPY 564/570, BODIPY 576/589, BODIPY 581/591, BODIPY TR, BODIPY 630/650, BODIPY 650/665, Cascade Blue, Cascade Yellow, Dansyl, lissamine rhodamine B, Marina Blue, Oregon Green 488, Oregon Green 514, Pacific Blue, rhodamine 6G, rhodamine green, rhodamine red, tetramethylrhodamine, Texas Red (available from Molecular Probes, Inc., Eugene, OR, USA).

-65-

The polypeptides of the present invention can also be conjugated to fluorophores, other proteins, and other macromolecules, using bifunctional linking reagents. Common homobifunctional reagents include, *e.g.*, APG, AEDP, BASED, BMB, BMDB, BMH, BMOE, BM[PEO]3, BM[PEO]4, BS3, BSOCOES, DFDNB, DMA, DMP, DMS, DPDPB, DSG, DSP (Lomant's Reagent), DSS, DST, DTBP, DTME, DTSSP, EGS, HBVS, Sulfo-BSOCOES, Sulfo-DST, Sulfo-EGS (all available from Pierce, Rockford, IL, USA); common heterobifunctional cross-linkers include ABH, AMAS, ANB-NOS, APDP, ASBA, BMFA, BMPH, BMPS, EDC, EMCA, EMCH, EMCS, KMUA, KMUH, GMBS, LC-SMCC, LC-SPDP, MBS, M2C2H, MPBH, MSA, NHS-ASA, PDPH, PMPI, SADP, SAED, SAND, SANPAH, SASD, SATP, SBAP, SFAD, SIA, SIAB, SMCC, SMPB, SMPH, SMPT, SPDP, Sulfo-EMCS, Sulfo-GMBS, Sulfo-HSAB, Sulfo-KMUS, Sulfo-LC-SPDP, Sulfo-MBS, Sulfo-NHS-LC-ASA, Sulfo-SADP, Sulfo-SANPAH, Sulfo-SIAB, Sulfo-SMCC, Sulfo-SMPB, Sulfo-LC-SMPT, SVSB, TFCS (all available Pierce, Rockford, IL, USA).

The polypeptides, fragments, and fusion proteins of the present invention can be conjugated, using such cross-linking reagents, to fluorophores that are not amine- or thiol-reactive. Other labels that usefully can be conjugated to the polypeptides, fragments, and fusion proteins of the present invention include radioactive labels, echosonographic contrast reagents, and MRI contrast agents.

The polypeptides, fragments, and fusion proteins of the present invention can also usefully be conjugated using cross-linking agents to carrier proteins, such as KLH, bovine thyroglobulin, and even bovine serum albumin (BSA), to increase immunogenicity for raising anti-BSP antibodies.

The polypeptides, fragments, and fusion proteins of the present invention can also usefully be conjugated to polyethylene glycol (PEG); PEGylation increases the serum half-life of proteins administered intravenously for replacement therapy. Delgado *et al.*, *Crit. Rev. Ther. Drug Carrier Syst.* 9(3-4): 249-304 (1992); Scott *et al.*, *Curr. Pharm. Des.* 4(6): 423-38 (1998); DeSantis *et al.*, *Curr. Opin. Biotechnol.* 10(4): 324-30 (1999), incorporated herein by reference in their entireties. PEG monomers can be attached to the protein directly or through a linker, with PEGylation using PEG monomers activated with tresyl chloride (2,2,2-trifluoroethanesulphonyl chloride) permitting direct attachment under mild conditions.

In yet another embodiment, the invention provides analogs of a polypeptide encoded by a nucleic acid molecule according to the instant invention. In a preferred

-66-

embodiment, the polypeptide is a BSP. In a more preferred embodiment, the analog is derived from a polypeptide having part or all of the amino acid sequence of SEQ ID NO: 172 through 295. In a preferred embodiment, the analog is one that comprises one or more substitutions of non-natural amino acids or non-native inter-residue bonds

5 compared to the naturally-occurring polypeptide. In general, the non-peptide analog is structurally similar to a BSP, but one or more peptide linkages is replaced by a linkage selected from the group consisting of --CH₂NH--, --CH₂S--, --CH₂-CH₂--,

--CH=CH--(cis and trans), --COCH₂--, --CH(OH)CH₂-- and --CH₂SO--. In another embodiment, the non-peptide analog comprises substitution of one or more amino acids

10 of a BSP with a D-amino acid of the same type or other non-natural amino acid in order to generate more stable peptides. D-amino acids can readily be incorporated during chemical peptide synthesis: peptides assembled from D-amino acids are more resistant to proteolytic attack; incorporation of D-amino acids can also be used to confer specific three-dimensional conformations on the peptide. Other amino acid analogues commonly

15 added during chemical synthesis include ornithine, norleucine, phosphorylated amino acids (typically phosphoserine, phosphothreonine, phosphotyrosine), L-malonyltyrosine, a non-hydrolyzable analog of phosphotyrosine (*see, e.g., Kole et al., Biochem. Biophys. Res. Com.* 209: 817-821 (1995)), and various halogenated phenylalanine derivatives.

Non-natural amino acids can be incorporated during solid phase chemical

20 synthesis or by recombinant techniques, although the former is typically more common. Solid phase chemical synthesis of peptides is well established in the art. Procedures are described, inter alia, in Chan *et al.* (eds.), Fmoc Solid Phase Peptide Synthesis: A Practical Approach (Practical Approach Series), Oxford Univ. Press (March 2000); Jones, Amino Acid and Peptide Synthesis (Oxford Chemistry Primers, No 7), Oxford

25 Univ. Press (1992); and Bodanszky, Principles of Peptide Synthesis (Springer Laboratory), Springer Verlag (1993); the disclosures of which are incorporated herein by reference in their entireties.

Amino acid analogues having detectable labels are also usefully incorporated during synthesis to provide derivatives and analogs. Biotin, for example can be added

30 using biotinoyl-(9-fluorenylmethoxycarbonyl)-L-lysine (Fmoc biocytin) (Molecular Probes, Eugene, OR, USA). Biotin can also be added enzymatically by incorporation into a fusion protein of a *E. coli* BirA substrate peptide. The Fmoc and tBOC derivatives of dabcyL-L-lysine (Molecular Probes, Inc., Eugene, OR, USA) can be used to incorporate the dabcyL chromophore at selected sites in the peptide sequence during

-67-

synthesis. The aminonaphthalene derivative EDANS, the most common fluorophore for pairing with the dabcyI quencher in fluorescence resonance energy transfer (FRET) systems, can be introduced during automated synthesis of peptides by using EDANS-FMOC-L-glutamic acid or the corresponding *t*BOC derivative (both from
5 Molecular Probes, Inc., Eugene, OR, USA). Tetramethylrhodamine fluorophores can be incorporated during automated FMOC synthesis of peptides using (FMOC)-TMR-L-lysine (Molecular Probes, Inc. Eugene, OR, USA).

Other useful amino acid analogues that can be incorporated during chemical synthesis include aspartic acid, glutamic acid, lysine, and tyrosine analogues having allyl
10 side-chain protection (Applied Biosystems, Inc., Foster City, CA, USA); the allyl side chain permits synthesis of cyclic, branched-chain, sulfonated, glycosylated, and phosphorylated peptides.

A large number of other FMOC-protected non-natural amino acid analogues capable of incorporation during chemical synthesis are available commercially,
15 including, *e.g.*, Fmoc-2-aminobicyclo[2.2.1]heptane-2-carboxylic acid, Fmoc-3-endo-aminobicyclo[2.2.1]heptane-2-endo-carboxylic acid, Fmoc-3-exo-aminobicyclo[2.2.1]heptane-2-exo-carboxylic acid, Fmoc-3-endo-amino-bicyclo[2.2.1]hept-5-ene-2-endo-carboxylic acid, Fmoc-3-exo-amino-bicyclo[2.2.1]hept-5-ene-2-exo-carboxylic acid, Fmoc-cis-2-amino-1-cyclohexanecarboxylic acid, Fmoc-
20 trans-2-amino-1-cyclohexanecarboxylic acid, Fmoc-1-amino-1-cyclopentanecarboxylic acid, Fmoc-cis-2-amino-1-cyclopentanecarboxylic acid, Fmoc-1-amino-1-cyclopropanecarboxylic acid, Fmoc-D-2-amino-4-(ethylthio)butyric acid, Fmoc-L-2-amino-4-(ethylthio)butyric acid, Fmoc-L-buthionine, Fmoc-S-methyl-L-Cysteine, Fmoc-2-aminobenzoic acid (anthranillic acid), Fmoc-3-aminobenzoic acid, Fmoc-4-
25 aminobenzoic acid, Fmoc-2-aminobenzophenone-2'-carboxylic acid, Fmoc-N-(4-aminobenzoyl)- β -alanine, Fmoc-2-amino-4,5-dimethoxybenzoic acid, Fmoc-4-aminohippuric acid, Fmoc-2-amino-3-hydroxybenzoic acid, Fmoc-2-amino-5-hydroxybenzoic acid, Fmoc-3-amino-4-hydroxybenzoic acid, Fmoc-4-amino-3-hydroxybenzoic acid, Fmoc-4-amino-2-hydroxybenzoic acid, Fmoc-5-amino-2-
30 hydroxybenzoic acid, Fmoc-2-amino-3-methoxybenzoic acid, Fmoc-4-amino-3-methoxybenzoic acid, Fmoc-2-amino-3-methylbenzoic acid, Fmoc-2-amino-5-methylbenzoic acid, Fmoc-2-amino-6-methylbenzoic acid, Fmoc-3-amino-2-methylbenzoic acid, Fmoc-3-amino-4-methylbenzoic acid, Fmoc-4-amino-3-methylbenzoic acid, Fmoc-3-amino-2-naphtioic acid, Fmoc-D,L-3-amino-3-

-68-

phenylpropionic acid, Fmoc-L-Methyldopa, Fmoc-2-amino-4,6-dimethyl-3-pyridinecarboxylic acid, Fmoc-D,L-amino-2-thiophenacetic acid, Fmoc-4-(carboxymethyl)piperazine, Fmoc-4-carboxypiperazine, Fmoc-4-(carboxymethyl)homopiperazine, Fmoc-4-phenyl-4-piperidinecarboxylic acid, Fmoc-L-1,2,3,4-tetrahydronorharman-3-carboxylic acid, Fmoc-L-thiazolidine-4-carboxylic acid, all available from The Peptide Laboratory (Richmond, CA, USA).

Non-natural residues can also be added biosynthetically by engineering a suppressor tRNA, typically one that recognizes the UAG stop codon, by chemical aminoacylation with the desired unnatural amino acid. Conventional site-directed mutagenesis is used to introduce the chosen stop codon UAG at the site of interest in the protein gene. When the acylated suppressor tRNA and the mutant gene are combined in an *in vitro* transcription/translation system, the unnatural amino acid is incorporated in response to the UAG codon to give a protein containing that amino acid at the specified position. Liu *et al.*, *Proc. Natl Acad. Sci. USA* 96(9): 4780-5 (1999); Wang *et al.*, *Science* 292(5516): 498-500 (2001).

Fusion Proteins

The present invention further provides fusions of each of the polypeptides and fragments of the present invention to heterologous polypeptides. In a preferred embodiment, the polypeptide is a BSP. In a more preferred embodiment, the polypeptide that is fused to the heterologous polypeptide comprises part or all of the amino acid sequence of SEQ ID NO: 172 through 295, or is a mutein, homologous polypeptide, analog or derivative thereof. In an even more preferred embodiment, the nucleic acid molecule encoding the fusion protein comprises all or part of the nucleic acid sequence of SEQ ID NO: 1 through 171, or comprises all or part of a nucleic acid sequence that selectively hybridizes or is homologous to a nucleic acid molecule comprising a nucleic acid sequence of SEQ ID NO: 1 through 171.

The fusion proteins of the present invention will include at least one fragment of the protein of the present invention, which fragment is at least 6, typically at least 8, often at least 15, and usefully at least 16, 17, 18, 19, or 20 amino acids long. The fragment of the protein of the present to be included in the fusion can usefully be at least 25 amino acids long, at least 50 amino acids long, and can be at least 75, 100, or even 150 amino acids long. Fusions that include the entirety of the proteins of the present invention have particular utility.

The heterologous polypeptide included within the fusion protein of the present invention is at least 6 amino acids in length, often at least 8 amino acids in length, and usefully at least 15, 20, and 25 amino acids in length. Fusions that include larger polypeptides, such as the IgG Fc region, and even entire proteins (such as GFP
5 chromophore-containing proteins) are particular useful.

As described above in the description of vectors and expression vectors of the present invention, which discussion is incorporated here by reference in its entirety, heterologous polypeptides to be included in the fusion proteins of the present invention can usefully include those designed to facilitate purification and/or visualization of
10 recombinantly-expressed proteins. *See, e.g.*, Ausubel, Chapter 16, (1992), *supra*. Although purification tags can also be incorporated into fusions that are chemically synthesized, chemical synthesis typically provides sufficient purity that further purification by HPLC suffices; however, visualization tags as above described retain their utility even when the protein is produced by chemical synthesis, and when so
15 included render the fusion proteins of the present invention useful as directly detectable markers of the presence of a polypeptide of the invention.

As also discussed above, heterologous polypeptides to be included in the fusion proteins of the present invention can usefully include those that facilitate secretion of recombinantly expressed proteins — into the periplasmic space or extracellular milieu for
20 prokaryotic hosts, into the culture medium for eukaryotic cells — through incorporation of secretion signals and/or leader sequences. For example, a His⁶ tagged protein can be purified on a Ni affinity column and a GST fusion protein can be purified on a glutathione affinity column. Similarly, a fusion protein comprising the Fc domain of IgG can be purified on a Protein A or Protein G column and a fusion protein comprising an
25 epitope tag such as myc can be purified using an immunoaffinity column containing an anti-c-myc antibody. It is preferable that the epitope tag be separated from the protein encoded by the essential gene by an enzymatic cleavage site that can be cleaved after purification. See also the discussion of nucleic acid molecules encoding fusion proteins that may be expressed on the surface of a cell.

30 Other useful protein fusions of the present invention include those that permit use of the protein of the present invention as bait in a yeast two-hybrid system. *See Bartel et al. (eds.), The Yeast Two-Hybrid System*, Oxford University Press (1997); *Zhu et al., Yeast Hybrid Technologies*, Eaton Publishing (2000); *Fields et al., Trends Genet.* 10(8): 286-92 (1994); *Mendelsohn et al., Curr. Opin. Biotechnol.* 5(5): 482-6 (1994); *Luban et*

al., *Curr. Opin. Biotechnol.* 6(1): 59-64 (1995); Allen *et al.*, *Trends Biochem. Sci.* 20(12): 511-6 (1995); Drees, *Curr. Opin. Chem. Biol.* 3(1): 64-70 (1999); Topcu *et al.*, *Pharm. Res.* 17(9): 1049-55 (2000); Fashena *et al.*, *Gene* 250(1-2): 1-14 (2000); ; Colas *et al.*, (1996) Genetic selection of peptide aptamers that recognize and inhibit cyclin-dependent kinase 2. *Nature* 380, 548-550; Norman, T. *et al.*, (1999) Genetic selection of peptide inhibitors of biological pathways. *Science* 285, 591-595, Fabbri *et al.*, (1999) Inhibition of mammalian cell proliferation by genetically selected peptide aptamers that functionally antagonize E2F activity. *Oncogene* 18, 4357-4363; Xu *et al.*, (1997) Cells that register logical relationships among proteins. *Proc Natl Acad Sci U S A.* 94, 12473-12478; Yang, *et al.*, (1995) Protein-peptide interactions analyzed with the yeast two-hybrid system. *Nuc. Acids Res.* 23, 1152-1156; Kolonin *et al.*, (1998) Targeting cyclin-dependent kinases in *Drosophila* with peptide aptamers. *Proc Natl Acad Sci U S A* 95, 14266-14271; Cohen *et al.*, (1998) An artificial cell-cycle inhibitor isolated from a combinatorial library. *Proc Natl Acad Sci U S A* 95, 14272-14277; Uetz, P.; Giot, L.; al, e.; Fields, S.; Rothberg, J. M. (2000) A comprehensive analysis of protein-protein interactions in *Saccharomyces cerevisiae*. *Nature* 403, 623-627; Ito, *et al.*, (2001) A comprehensive two-hybrid analysis to explore the yeast protein interactome. *Proc Natl Acad Sci U S A* 98, 4569-4574, the disclosures of which are incorporated herein by reference in their entirety. Typically, such fusion is to either *E. coli* LexA or yeast GAL4 DNA binding domains. Related bait plasmids are available that express the bait fused to a nuclear localization signal.

Other useful fusion proteins include those that permit display of the encoded protein on the surface of a phage or cell, fusions to intrinsically fluorescent proteins, such as green fluorescent protein (GFP), and fusions to the IgG Fc region, as described above, which discussion is incorporated here by reference in its entirety.

The polypeptides and fragments of the present invention can also usefully be fused to protein toxins, such as *Pseudomonas* exotoxin A, *diphtheria* toxin, *shiga* toxin A, *anthrax* toxin lethal factor, ricin, in order to effect ablation of cells that bind or take up the proteins of the present invention.

Fusion partners include, *inter alia*, *myc*, hemagglutinin (HA), GST, immunoglobulins, β -galactosidase, biotin *trpE*, protein A, β -lactamase, α -amylase, maltose binding protein, alcohol dehydrogenase, polyhistidine (for example, six histidine at the amino and/or carboxyl terminus of the polypeptide), *lacZ*, green fluorescent protein (GFP), yeast α mating factor, GAL4 transcription activation or DNA binding domain,

luciferase, and serum proteins such as ovalbumin, albumin and the constant domain of IgG. *See, e.g.*, Ausubel (1992), *supra* and Ausubel (1999), *supra*. Fusion proteins may also contain sites for specific enzymatic cleavage, such as a site that is recognized by enzymes such as Factor XIII, trypsin, pepsin, or any other enzyme known in the art.

- 5 Fusion proteins will typically be made by either recombinant nucleic acid methods, as described above, chemically synthesized using techniques well-known in the art (*e.g.*, a Merrifield synthesis), or produced by chemical cross-linking.

Another advantage of fusion proteins is that the epitope tag can be used to bind the fusion protein to a plate or column through an affinity linkage for screening binding
10 proteins or other molecules that bind to the BSP.

As further described below, the isolated polypeptides, muteins, fusion proteins, homologous proteins or allelic variants of the present invention can readily be used as specific immunogens to raise antibodies that specifically recognize BSPs, their allelic variants and homologues. The antibodies, in turn, can be used, *inter alia*, specifically to
15 assay for the polypeptides of the present invention, particularly BSPs, *e.g.* by ELISA for detection of protein fluid samples, such as serum, by immunohistochemistry or laser scanning cytometry, for detection of protein in tissue samples, or by flow cytometry, for detection of intracellular protein in cell suspensions, for specific antibody-mediated isolation and/or purification of BSPs, as for example by immunoprecipitation, and for use
20 as specific agonists or antagonists of BSPs.

One may determine whether polypeptides including muteins, fusion proteins, homologous proteins or allelic variants are functional by methods known in the art. For instance, residues that are tolerant of change while retaining function can be identified by altering the protein at known residues using methods known in the art, such as alanine
25 scanning mutagenesis, Cunningham *et al.*, *Science* 244(4908): 1081-5 (1989); transposon linker scanning mutagenesis, Chen *et al.*, *Gene* 263(1-2): 39-48 (2001); combinations of homolog- and alanine-scanning mutagenesis, Jin *et al.*, *J. Mol. Biol.* 226(3): 851-65 (1992); combinatorial alanine scanning, Weiss *et al.*, *Proc. Natl. Acad. Sci USA* 97(16): 8950-4 (2000), followed by functional assay. Transposon linker scanning kits are
30 available commercially (New England Biolabs, Beverly, MA, USA, catalog. no. E7-102S; EZ::TN™ In-Frame Linker Insertion Kit, catalogue no. EZI04KN, Epicentre Technologies Corporation, Madison, WI, USA).

Purification of the polypeptides including fragments, homologous polypeptides, muteins, analogs, derivatives and fusion proteins is well-known and within the skill of

-72-

one having ordinary skill in the art. *See, e.g.*, Scopes, Protein Purification, 2d ed. (1987). Purification of recombinantly expressed polypeptides is described above. Purification of chemically-synthesized peptides can readily be effected, *e.g.*, by HPLC.

Accordingly, it is an aspect of the present invention to provide the isolated
5 proteins of the present invention in pure or substantially pure form in the presence of absence of a stabilizing agent. Stabilizing agents include both proteinaceous or non-proteinaceous material and are well-known in the art. Stabilizing agents, such as albumin and polyethylene glycol (PEG) are known and are commercially available.

Although high levels of purity are preferred when the isolated proteins of the
10 present invention are used as therapeutic agents, such as in vaccines and as replacement therapy, the isolated proteins of the present invention are also useful at lower purity. For example, partially purified proteins of the present invention can be used as immunogens to raise antibodies in laboratory animals.

In preferred embodiments, the purified and substantially purified proteins of the
15 present invention are in compositions that lack detectable ampholytes, acrylamide monomers, bis-acrylamide monomers, and polyacrylamide.

The polypeptides, fragments, analogs, derivatives and fusions of the present invention can usefully be attached to a substrate. The substrate can be porous or solid, planar or non-planar; the bond can be covalent or noncovalent.

20 For example, the polypeptides, fragments, analogs, derivatives and fusions of the present invention can usefully be bound to a porous substrate, commonly a membrane, typically comprising nitrocellulose, polyvinylidene fluoride (PVDF), or cationically derivatized, hydrophilic PVDF; so bound, the proteins, fragments, and fusions of the present invention can be used to detect and quantify antibodies, *e.g.* in serum, that bind
25 specifically to the immobilized protein of the present invention.

As another example, the polypeptides, fragments, analogs, derivatives and fusions of the present invention can usefully be bound to a substantially nonporous substrate, such as plastic, to detect and quantify antibodies, *e.g.* in serum, that bind specifically to the immobilized protein of the present invention. Such plastics include
30 polymethylacrylic, polyethylene, polypropylene, polyacrylate, polymethylmethacrylate, polyvinylchloride, polytetrafluoroethylene, polystyrene, polycarbonate, polyacetal, polysulfone, celluloseacetate, cellulosenitrate, nitrocellulose, or mixtures thereof; when the assay is performed in a standard microtiter dish, the plastic is typically polystyrene.

-73-

The polypeptides, fragments, analogs, derivatives and fusions of the present invention can also be attached to a substrate suitable for use as a surface enhanced laser desorption ionization source; so attached, the protein, fragment, or fusion of the present invention is useful for binding and then detecting secondary proteins that bind with
5 sufficient affinity or avidity to the surface-bound protein to indicate biologic interaction there between. The proteins, fragments, and fusions of the present invention can also be attached to a substrate suitable for use in surface plasmon resonance detection; so attached, the protein, fragment, or fusion of the present invention is useful for binding and then detecting secondary proteins that bind with sufficient affinity or avidity to the
10 surface-bound protein to indicate biological interaction there between.

Antibodies

In another aspect, the invention provides antibodies, including fragments and derivatives thereof, that bind specifically to polypeptides encoded by the nucleic acid
15 molecules of the invention, as well as antibodies that bind to fragments, muteins, derivatives and analogs of the polypeptides. In a preferred embodiment, the antibodies are specific for a polypeptide that is a BSP, or a fragment, mutein, derivative, analog or fusion protein thereof. In a more preferred embodiment, the antibodies are specific for a polypeptide that comprises SEQ ID NO: 172 through 295, or a fragment, mutein,
20 derivative, analog or fusion protein thereof.

The antibodies of the present invention can be specific for linear epitopes, discontinuous epitopes, or conformational epitopes of such proteins or protein fragments, either as present on the protein in its native conformation or, in some cases, as present on the proteins as denatured, as, *e.g.*, by solubilization in SDS. New epitopes may be also
25 due to a difference in post translational modifications (PTMs) in disease versus normal tissue. For example, a particular site on a BSP may be glycosylated in cancerous cells, but not glycosylated in normal cells or visa versa. In addition, alternative splice forms of a BSP may be indicative of cancer. Differential degradation of the C or N-terminus of a BSP may also be a marker or target for anticancer therapy. For example, a BSP may
30 be N-terminal degraded in cancer cells exposing new epitopes to which antibodies may selectively bind for diagnostic or therapeutic uses.

As is well-known in the art, the degree to which an antibody can discriminate as among molecular species in a mixture will depend, in part, upon the conformational relatedness of the species in the mixture; typically, the antibodies of the present invention

will discriminate over adventitious binding to non-BSP polypeptides by at least 2-fold, more typically by at least 5-fold, typically by more than 10-fold, 25-fold, 50-fold, 75-fold, and often by more than 100-fold, and on occasion by more than 500-fold or 1000-fold. When used to detect the proteins or protein fragments of the present invention, the
5 antibody of the present invention is sufficiently specific when it can be used to determine the presence of the protein of the present invention in samples derived from human breast.

Typically, the affinity or avidity of an antibody (or antibody multimer, as in the case of an IgM pentamer) of the present invention for a protein or protein fragment of the
10 present invention will be at least about 1×10^{-6} molar (M), typically at least about 5×10^{-7} M, 1×10^{-7} M, with affinities and avidities of at least 1×10^{-8} M, 5×10^{-9} M, 1×10^{-10} M and up to 1×10^{-13} M proving especially useful.

The antibodies of the present invention can be naturally-occurring forms, such as IgG, IgM, IgD, IgE, IgY, and IgA, from any avian, reptilian, or mammalian species.

15 Human antibodies can, but will infrequently, be drawn directly from human donors or human cells. In this case, antibodies to the proteins of the present invention will typically have resulted from fortuitous immunization, such as autoimmune immunization, with the protein or protein fragments of the present invention. Such antibodies will typically, but will not invariably, be polyclonal. In addition, individual
20 polyclonal antibodies may be isolated and cloned to generate monoclonals.

Human antibodies are more frequently obtained using transgenic animals that express human immunoglobulin genes, which transgenic animals can be affirmatively immunized with the protein immunogen of the present invention. Human Ig-transgenic mice capable of producing human antibodies and methods of producing human
25 antibodies therefrom upon specific immunization are described, *inter alia*, in U.S. Patents 6,162,963; 6,150,584; 6,114,598; 6,075,181; 5,939,598; 5,877,397; 5,874,299; 5,814,318; 5,789,650; 5,770,429; 5,661,016; 5,633,425; 5,625,126; 5,569,825; 5,545,807; 5,545,806, and 5,591,669, the disclosures of which are incorporated herein by reference in their entireties. Such antibodies are typically monoclonal, and are typically
30 produced using techniques developed for production of murine antibodies.

Human antibodies are particularly useful, and often preferred, when the antibodies of the present invention are to be administered to human beings as *in vivo* diagnostic or therapeutic agents, since recipient immune response to the administered

-75-

antibody will often be substantially less than that occasioned by administration of an antibody derived from another species, such as mouse.

IgG, IgM, IgD, IgE, IgY, and IgA antibodies of the present invention can also be obtained from other species, including mammals such as rodents (typically mouse, but
5 also rat, guinea pig, and hamster) lagomorphs, typically rabbits, and also larger mammals, such as sheep, goats, cows, and horses, and other egg laying birds or reptiles such as chickens or alligators. For example, avian antibodies may be generated using techniques described in WO 00/29444, published 25 May 2000, the contents of which are hereby incorporated in their entirety. In such cases, as with the transgenic human-
10 antibody-producing non-human mammals, fortuitous immunization is not required, and the non-human mammal is typically affirmatively immunized, according to standard immunization protocols, with the protein or protein fragment of the present invention.

As discussed above, virtually all fragments of 8 or more contiguous amino acids of the proteins of the present invention can be used effectively as immunogens when
15 conjugated to a carrier, typically a protein such as bovine thyroglobulin, keyhole limpet hemocyanin, or bovine serum albumin, conveniently using a bifunctional linker such as those described elsewhere above, which discussion is incorporated by reference here.

Immunogenicity can also be conferred by fusion of the polypeptide and fragments of the present invention to other moieties. For example, peptides of the present invention
20 can be produced by solid phase synthesis on a branched polylysine core matrix; these multiple antigenic peptides (MAPs) provide high purity, increased avidity, accurate chemical definition and improved safety in vaccine development. Tam *et al.*, *Proc. Natl. Acad. Sci. USA* 85: 5409-5413 (1988); Posnett *et al.*, *J. Biol. Chem.* 263: 1719-1725 (1988).

25 Protocols for immunizing non-human mammals or avian species are well-established in the art. See Harlow *et al.* (eds.), Using Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory (1998); Coligan *et al.* (eds.), Current Protocols in Immunology, John Wiley & Sons, Inc. (2001); Zola, Monoclonal Antibodies: Preparation and Use of Monoclonal Antibodies and Engineered Antibody Derivatives (Basics: From
30 Background to Bench), Springer Verlag (2000); Gross M, Speck *J.Dtsch. Tierarztl. Wochenschr.* 103: 417-422 (1996), the disclosures of which are incorporated herein by reference. Immunization protocols often include multiple immunizations, either with or without adjuvants such as Freund's complete adjuvant and Freund's incomplete adjuvant, and may include naked DNA immunization (Moss, *Semin. Immunol.* 2: 317-327 (1990).

-76-

Antibodies from non-human mammals and avian species can be polyclonal or monoclonal, with polyclonal antibodies having certain advantages in immunohistochemical detection of the proteins of the present invention and monoclonal antibodies having advantages in identifying and distinguishing particular epitopes of the proteins of the present invention. Antibodies from avian species may have particular
5 advantage in detection of the proteins of the present invention, in human serum or tissues (Vikingo et al., *Biosens. Bioelectron.* 13: 1257-1262 (1998).

Following immunization, the antibodies of the present invention can be produced using any art-accepted technique. Such techniques are well-known in the art, Coligan,
10 *supra*; Zola, *supra*; Howard *et al.* (eds.), Basic Methods in Antibody Production and Characterization, CRC Press (2000); Harlow, *supra*; Davis (ed.), Monoclonal Antibody Protocols, Vol. 45, Humana Press (1995); Delves (ed.), Antibody Production: Essential Techniques, John Wiley & Son Ltd (1997); Kenney, Antibody Solution: An Antibody Methods Manual, Chapman & Hall (1997), incorporated herein by reference in their
15 entireties, and thus need not be detailed here.

Briefly, however, such techniques include, *inter alia*, production of monoclonal antibodies by hybridomas and expression of antibodies or fragments or derivatives thereof from host cells engineered to express immunoglobulin genes or fragments thereof. These two methods of production are not mutually exclusive: genes encoding
20 antibodies specific for the proteins or protein fragments of the present invention can be cloned from hybridomas and thereafter expressed in other host cells. Nor need the two necessarily be performed together: *e.g.*, genes encoding antibodies specific for the proteins and protein fragments of the present invention can be cloned directly from B
25 cells known to be specific for the desired protein, as further described in U.S. Patent 5,627,052, the disclosure of which is incorporated herein by reference in its entirety, or from antibody-displaying phage.

Recombinant expression in host cells is particularly useful when fragments or derivatives of the antibodies of the present invention are desired.

Host cells for recombinant production of either whole antibodies, antibody
30 fragments, or antibody derivatives can be prokaryotic or eukaryotic.

Prokaryotic hosts are particularly useful for producing phage displayed antibodies of the present invention.

The technology of phage-displayed antibodies, in which antibody variable region fragments are fused, for example, to the gene III protein (pIII) or gene VIII protein

(pVIII) for display on the surface of filamentous phage, such as M13, is by now well-established. *See, e.g.,* Sidhu, *Curr. Opin. Biotechnol.* 11(6): 610-6 (2000); Griffiths *et al., Curr. Opin. Biotechnol.* 9(1): 102-8 (1998); Hoogenboom *et al., Immunotechnology*, 4(1): 1-20 (1998); Rader *et al., Current Opinion in Biotechnology* 8: 503-508 (1997);
5 Aujame *et al., Human Antibodies* 8: 155-168 (1997); Hoogenboom, *Trends in Biotechnol.* 15: 62-70 (1997); de Kruif *et al.,* 17: 453-455 (1996); Barbas *et al., Trends in Biotechnol.* 14: 230-234 (1996); Winter *et al., Ann. Rev. Immunol.* 433-455 (1994).
Techniques and protocols required to generate, propagate, screen (pan), and use the antibody fragments from such libraries have recently been compiled. *See, e.g.,* Barbas
10 (2001), *supra*; Kay, *supra*; Abelson, *supra*, the disclosures of which are incorporated herein by reference in their entireties.

Typically, phage-displayed antibody fragments are scFv fragments or Fab fragments; when desired, full length antibodies can be produced by cloning the variable regions from the displaying phage into a complete antibody and expressing the full length
15 antibody in a further prokaryotic or a eukaryotic host cell.

Eukaryotic cells are also useful for expression of the antibodies, antibody fragments, and antibody derivatives of the present invention.

For example, antibody fragments of the present invention can be produced in *Pichia pastoris* and in *Saccharomyces cerevisiae*. *See, e.g.,* Takahashi *et al., Biosci.*
20 *Biotechnol. Biochem.* 64(10): 2138-44 (2000); Freyre *et al., J. Biotechnol.* 76(2-3):1 57-63 (2000); Fischer *et al., Biotechnol. Appl. Biochem.* 30 (Pt 2): 117-20 (1999); Pennell *et al., Res. Immunol.* 149(6): 599-603 (1998); Eldin *et al., J. Immunol. Methods.* 201(1): 67-75 (1997);, Frenken *et al., Res. Immunol.* 149(6): 589-99 (1998); Shusta *et al., Nature Biotechnol.* 16(8): 773-7 (1998), the disclosures of which are incorporated herein
25 by reference in their entireties.

Antibodies, including antibody fragments and derivatives, of the present invention can also be produced in insect cells. *See, e.g.,* Li *et al., Protein Expr. Purif.* 21(1): 121-8 (2001); Ailor *et al., Biotechnol. Bioeng.* 58(2-3): 196-203 (1998); Hsu *et al., Biotechnol. Prog.* 13(1): 96-104 (1997); Edelman *et al., Immunology* 91(1): 13-9 (1997);
30 and Nesbit *et al., J. Immunol. Methods* 151(1-2): 201-8 (1992), the disclosures of which are incorporated herein by reference in their entireties.

Antibodies and fragments and derivatives thereof of the present invention can also be produced in plant cells, particularly maize or tobacco, Giddings *et al., Nature Biotechnol.* 18(11): 1151-5 (2000); Gavilondo *et al., Biotechniques* 29(1): 128-38 (2000);

Fischer *et al.*, *J. Biol. Regul. Homeost. Agents* 14(2): 83-92 (2000); Fischer *et al.*, *Biotechnol. Appl. Biochem.* 30 (Pt 2): 113-6 (1999); Fischer *et al.*, *Biol. Chem.* 380(7-8): 825-39 (1999); Russell, *Curr. Top. Microbiol. Immunol.* 240: 119-38 (1999); and Ma *et al.*, *Plant Physiol.* 109(2): 341-6 (1995), the disclosures of which are incorporated herein
5 by reference in their entireties.

Antibodies, including antibody fragments and derivatives, of the present invention can also be produced in transgenic, non-human, mammalian milk. *See, e.g.* Pollock *et al.*, *J. Immunol Methods.* 231: 147-57 (1999); Young *et al.*, *Res. Immunol.* 149: 609-10 (1998); Limonta *et al.*, *Immunotechnology* 1: 107-13 (1995), the disclosures
10 of which are incorporated herein by reference in their entireties.

Mammalian cells useful for recombinant expression of antibodies, antibody fragments, and antibody derivatives of the present invention include CHO cells, COS cells, 293 cells, and myeloma cells.

Verma *et al.*, *J. Immunol. Methods* 216(1-2):165-81 (1998), herein incorporated
15 by reference, review and compare bacterial, yeast, insect and mammalian expression systems for expression of antibodies.

Antibodies of the present invention can also be prepared by cell free translation, as further described in Merk *et al.*, *J. Biochem.* (Tokyo) 125(2): 328-33 (1999) and Ryabova *et al.*, *Nature Biotechnol.* 15(1): 79-84 (1997), and in the milk of transgenic
20 animals, as further described in Pollock *et al.*, *J. Immunol. Methods* 231(1-2): 147-57 (1999), the disclosures of which are incorporated herein by reference in their entireties.

The invention further provides antibody fragments that bind specifically to one or more of the proteins and protein fragments of the present invention, to one or more of the proteins and protein fragments encoded by the isolated nucleic acids of the present
25 invention, or the binding of which can be competitively inhibited by one or more of the proteins and protein fragments of the present invention or one or more of the proteins and protein fragments encoded by the isolated nucleic acids of the present invention.

Among such useful fragments are Fab, Fab', Fv, F(ab)'₂, and single chain Fv (scFv) fragments. Other useful fragments are described in Hudson, *Curr. Opin.*
30 *Biotechnol.* 9(4): 395-402 (1998).

It is also an aspect of the present invention to provide antibody derivatives that bind specifically to one or more of the proteins and protein fragments of the present invention, to one or more of the proteins and protein fragments encoded by the isolated nucleic acids of the present invention, or the binding of which can be competitively

-79-

inhibited by one or more of the proteins and protein fragments of the present invention or one or more of the proteins and protein fragments encoded by the isolated nucleic acids of the present invention.

Among such useful derivatives are chimeric, primatized, and humanized
5 antibodies; such derivatives are less immunogenic in human beings, and thus more suitable for *in vivo* administration, than are unmodified antibodies from non-human mammalian species. Another useful derivative is PEGylation to increase the serum half life of the antibodies.

Chimeric antibodies typically include heavy and/or light chain variable regions
10 (including both CDR and framework residues) of immunoglobulins of one species, typically mouse, fused to constant regions of another species, typically human. *See, e.g.,* United States Patent No. 5,807,715; Morrison *et al.*, *Proc. Natl. Acad. Sci USA* 81(21): 6851-5 (1984); Sharon *et al.*, *Nature* 309(5966): 364-7 (1984); Takeda *et al.*, *Nature* 314(6010): 452-4 (1985), the disclosures of which are incorporated herein by reference in
15 their entireties. Primatized and humanized antibodies typically include heavy and/or light chain CDRs from a murine antibody grafted into a non-human primate or human antibody V region framework, usually further comprising a human constant region, Riechmann *et al.*, *Nature* 332(6162): 323-7 (1988); Co *et al.*, *Nature* 351(6326): 501-2 (1991); United States Patent Nos. 6,054,297; 5,821,337; 5,770,196; 5,766,886;
20 5,821,123; 5,869,619; 6,180,377; 6,013,256; 5,693,761; and 6,180,370, the disclosures of which are incorporated herein by reference in their entireties.

Other useful antibody derivatives of the invention include heteromeric antibody complexes and antibody fusions, such as diabodies (bispecific antibodies), single-chain diabodies, and intrabodies.

25 It is contemplated that the nucleic acids encoding the antibodies of the present invention can be operably joined to other nucleic acids forming a recombinant vector for cloning or for expression of the antibodies of the invention. The present invention includes any recombinant vector containing the coding sequences, or part thereof, whether for eukaryotic transduction, transfection or gene therapy. Such vectors may be
30 prepared using conventional molecular biology techniques, known to those with skill in the art, and would comprise DNA encoding sequences for the immunoglobulin V-regions including framework and CDRs or parts thereof, and a suitable promoter either with or without a signal sequence for intracellular transport. Such vectors may be transduced or transfected into eukaryotic cells or used for gene therapy (Marasco *et al.*, *Proc. Natl.*

Acad. Sci. (USA) 90: 7889-7893 (1993); Duan et al., *Proc. Natl. Acad. Sci. (USA)* 91: 5075-5079 (1994), by conventional techniques, known to those with skill in the art.

The antibodies of the present invention, including fragments and derivatives thereof, can usefully be labeled. It is, therefore, another aspect of the present invention to
5 provide labeled antibodies that bind specifically to one or more of the proteins and protein fragments of the present invention, to one or more of the proteins and protein fragments encoded by the isolated nucleic acids of the present invention, or the binding of which can be competitively inhibited by one or more of the proteins and protein fragments of the present invention or one or more of the proteins and protein fragments
10 encoded by the isolated nucleic acids of the present invention.

The choice of label depends, in part, upon the desired use.

For example, when the antibodies of the present invention are used for immunohistochemical staining of tissue samples, the label is preferably an enzyme that catalyzes production and local deposition of a detectable product.

15 Enzymes typically conjugated to antibodies to permit their immunohistochemical visualization are well-known, and include alkaline phosphatase, β -galactosidase, glucose oxidase, horseradish peroxidase (HRP), and urease. Typical substrates for production and deposition of visually detectable products include o-nitrophenyl-beta-D-galactopyranoside (ONPG); o-phenylenediamine dihydrochloride (OPD); p-nitrophenyl
20 phosphate (PNPP); p-nitrophenyl-beta-D-galactopyranoside (PNPG); 3',3'-diaminobenzidine (DAB); 3-amino-9-ethylcarbazole (AEC); 4-chloro-1-naphthol (CN); 5-bromo-4-chloro-3-indolyl-phosphate (BCIP); ABTS®; BluoGal; iodinitrotetrazolium (INT); nitroblue tetrazolium chloride (NBT); phenazine methosulfate (PMS); phenolphthalein monophosphate (PMP); tetramethyl benzidine (TMB); tetranitroblue
25 tetrazolium (TNBT); X-Gal; X-Gluc; and X-Glucoside.

Other substrates can be used to produce products for local deposition that are luminescent. For example, in the presence of hydrogen peroxide (H_2O_2), horseradish peroxidase (HRP) can catalyze the oxidation of cyclic diacylhydrazides, such as luminol. Immediately following the oxidation, the luminol is in an excited state (intermediate
30 reaction product), which decays to the ground state by emitting light. Strong enhancement of the light emission is produced by enhancers, such as phenolic compounds. Advantages include high sensitivity, high resolution, and rapid detection without radioactivity and requiring only small amounts of antibody. See, e.g., Thorpe et al., *Methods Enzymol.* 133: 331-53 (1986); Kricka et al., *J. Immunoassay* 17(1): 67-83

(1996); and Lundqvist *et al.*, *J. Biolumin. Chemilumin.* 10(6): 353-9 (1995), the disclosures of which are incorporated herein by reference in their entireties. Kits for such enhanced chemiluminescent detection (ECL) are available commercially.

The antibodies can also be labeled using colloidal gold.

5 As another example, when the antibodies of the present invention are used, *e.g.*, for flow cytometric detection, for scanning laser cytometric detection, or for fluorescent immunoassay, they can usefully be labeled with fluorophores.

There are a wide variety of fluorophore labels that can usefully be attached to the antibodies of the present invention.

10 For flow cytometric applications, both for extracellular detection and for intracellular detection, common useful fluorophores can be fluorescein isothiocyanate (FITC), allophycocyanin (APC), R-phycoerythrin (PE), peridinin chlorophyll protein (PerCP), Texas Red, Cy3, Cy5, fluorescence resonance energy tandem fluorophores such as PerCP-Cy5.5, PE-Cy5, PE-Cy5.5, PE-Cy7, PE-Texas Red, and APC-Cy7.

15 Other fluorophores include, *inter alia*, Alexa Fluor® 350, Alexa Fluor® 488, Alexa Fluor® 532, Alexa Fluor® 546, Alexa Fluor® 568, Alexa Fluor® 594, Alexa Fluor® 647 (monoclonal antibody labeling kits available from Molecular Probes, Inc., Eugene, OR, USA), BODIPY dyes, such as BODIPY 493/503, BODIPY FL, BODIPY R6G, BODIPY 530/550, BODIPY TMR, BODIPY 558/568, BODIPY 558/568,
20 BODIPY 564/570, BODIPY 576/589, BODIPY 581/591, BODIPY TR, BODIPY 630/650, BODIPY 650/665, Cascade Blue, Cascade Yellow, Dansyl, lissamine rhodamine B, Marina Blue, Oregon Green 488, Oregon Green 514, Pacific Blue, rhodamine 6G, rhodamine green, rhodamine red, tetramethylrhodamine, Texas Red (available from Molecular Probes, Inc., Eugene, OR, USA), and Cy2, Cy3, Cy3.5, Cy5,
25 Cy5.5, Cy7, all of which are also useful for fluorescently labeling the antibodies of the present invention.

For secondary detection using labeled avidin, streptavidin, captavidin or neutravidin, the antibodies of the present invention can usefully be labeled with biotin.

When the antibodies of the present invention are used, *e.g.*, for Western blotting
30 applications, they can usefully be labeled with radioisotopes, such as ³³P, ³²P, ³⁵S, ³H, and ¹²⁵I.

As another example, when the antibodies of the present invention are used for radioimmunotherapy, the label can usefully be ²²⁸Th, ²²⁷Ac, ²²⁵Ac, ²²³Ra, ²¹³Bi, ²¹²Pb,

-82-

^{212}Bi , ^{211}At , ^{203}Pb , ^{194}Os , ^{188}Re , ^{186}Re , ^{153}Sm , ^{149}Tb , ^{131}I , ^{125}I , ^{111}In , ^{105}Rh , $^{99\text{m}}\text{Tc}$, ^{97}Ru , ^{90}Y , ^{90}Sr , ^{88}Y , ^{72}Se , ^{67}Cu , or ^{47}Sc .

As another example, when the antibodies of the present invention are to be used for *in vivo* diagnostic use, they can be rendered detectable by conjugation to MRI contrast agents, such as gadolinium diethylenetriaminepentaacetic acid (DTPA), Lauffer *et al.*, *Radiology* 207(2): 529-38 (1998), or by radioisotopic labeling.

As would be understood, use of the labels described above is not restricted to the application for which they are mentioned.

The antibodies of the present invention, including fragments and derivatives thereof, can also be conjugated to toxins, in order to target the toxin's ablative action to cells that display and/or express the proteins of the present invention. Commonly, the antibody in such immunotoxins is conjugated to *Pseudomonas* exotoxin A, diphtheria toxin, shiga toxin A, anthrax toxin lethal factor, or ricin. See Hall (ed.), Immunotoxin Methods and Protocols (Methods in Molecular Biology, vol. 166), Humana Press (2000); and Frankel *et al.* (eds.), Clinical Applications of Immunotoxins, Springer-Verlag (1998), the disclosures of which are incorporated herein by reference in their entireties.

The antibodies of the present invention can usefully be attached to a substrate, and it is, therefore, another aspect of the invention to provide antibodies that bind specifically to one or more of the proteins and protein fragments of the present invention, to one or more of the proteins and protein fragments encoded by the isolated nucleic acids of the present invention, or the binding of which can be competitively inhibited by one or more of the proteins and protein fragments of the present invention or one or more of the proteins and protein fragments encoded by the isolated nucleic acids of the present invention, attached to a substrate.

Substrates can be porous or nonporous, planar or nonplanar.

For example, the antibodies of the present invention can usefully be conjugated to filtration media, such as NHS-activated Sepharose or CNBr-activated Sepharose for purposes of immunoaffinity chromatography.

For example, the antibodies of the present invention can usefully be attached to paramagnetic microspheres, typically by biotin-streptavidin interaction, which microspheres can then be used for isolation of cells that express or display the proteins of the present invention. As another example, the antibodies of the present invention can usefully be attached to the surface of a microtiter plate for ELISA.

-83-

As noted above, the antibodies of the present invention can be produced in prokaryotic and eukaryotic cells. It is, therefore, another aspect of the present invention to provide cells that express the antibodies of the present invention, including hybridoma cells, B cells, plasma cells, and host cells recombinantly modified to express the
5 antibodies of the present invention.

In yet a further aspect, the present invention provides aptamers evolved to bind specifically to one or more of the proteins and protein fragments of the present invention, to one or more of the proteins and protein fragments encoded by the isolated nucleic acids of the present invention, or the binding of which can be competitively inhibited by
10 one or more of the proteins and protein fragments of the present invention or one or more of the proteins and protein fragments encoded by the isolated nucleic acids of the present invention.

In sum, one of skill in the art, provided with the teachings of this invention, has available a variety of methods which may be used to alter the biological properties of the
15 antibodies of this invention including methods which would increase or decrease the stability or half-life, immunogenicity, toxicity, affinity or yield of a given antibody molecule, or to alter it in any other way that may render it more suitable for a particular application.

Transgenic Animals and Cells

20

In another aspect, the invention provides transgenic cells and non-human organisms comprising nucleic acid molecules of the invention. In a preferred embodiment, the transgenic cells and non-human organisms comprise a nucleic acid molecule encoding a BSP. In a preferred embodiment, the BSP comprises an amino acid
25 sequence selected from SEQ ID NO: 172 through 295, or a fragment, mutein, homologous protein or allelic variant thereof. In another preferred embodiment, the transgenic cells and non-human organism comprise a BSNA of the invention, preferably a BSNA comprising a nucleotide sequence selected from the group consisting of SEQ ID NO: 1 through 171, or a part, substantially similar nucleic acid molecule, allelic variant
30 or hybridizing nucleic acid molecule thereof.

In another embodiment, the transgenic cells and non-human organisms have a targeted disruption or replacement of the endogenous orthologue of the human BSG. The transgenic cells can be embryonic stem cells or somatic cells. The transgenic non-human organisms can be chimeric, nonchimeric heterozygotes, and nonchimeric

homozygotes. Methods of producing transgenic animals are well-known in the art. *See, e.g.,* Hogan *et al.*, Manipulating the Mouse Embryo: A Laboratory Manual, 2d ed., Cold Spring Harbor Press (1999); Jackson *et al.*, Mouse Genetics and Transgenics: A Practical Approach, Oxford University Press (2000); and Pinkert, Transgenic Animal Technology: A Laboratory Handbook, Academic Press (1999).

Any technique known in the art may be used to introduce a nucleic acid molecule of the invention into an animal to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection. (*see, e.g.,* Paterson *et al.*, *Appl. Microbiol. Biotechnol.* 40: 691-698 (1994); Carver *et al.*, *Biotechnology* 11: 1263-1270 (1993); Wright *et al.*, *Biotechnology* 9: 830-834 (1991); and U.S. Patent 4,873,191 (1989) retrovirus-mediated gene transfer into germ lines, blastocysts or embryos (*see, e.g.,* Van der Putten *et al.*, *Proc. Natl. Acad. Sci., USA* 82: 6148-6152 (1985)); gene targeting in embryonic stem cells (*see, e.g.,* Thompson *et al.*, *Cell* 56: 313-321 (1989)); electroporation of cells or embryos (*see, e.g.,* Lo, 1983, *Mol. Cell. Biol.* 3: 1803-1814 (1983)); introduction using a gene gun (*see, e.g.,* Ulmer *et al.*, *Science* 259: 1745-49 (1993); introducing nucleic acid constructs into embryonic pluripotent stem cells and transferring the stem cells back into the blastocyst; and sperm-mediated gene transfer (*see, e.g.,* Lavitrano *et al.*, *Cell* 57: 717-723 (1989)).

Other techniques include, for example, nuclear transfer into enucleated oocytes of nuclei from cultured embryonic, fetal, or adult cells induced to quiescence (*see, e.g.,* Campell *et al.*, *Nature* 380: 64-66 (1996); Wilmut *et al.*, *Nature* 385: 810-813 (1997)). The present invention provides for transgenic animals that carry the transgene (*i.e.*, a nucleic acid molecule of the invention) in all their cells, as well as animals which carry the transgene in some, but not all their cells, *i.e.*, mosaic animals or chimeric animals.

The transgene may be integrated as a single transgene or as multiple copies, such as in concatamers, *e.g.*, head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, *e.g.*, the teaching of Lasko *et al. et al.*, *Proc. Natl. Acad. Sci. USA* 89: 6232-6236 (1992). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to

verify that integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques which include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and reverse transcriptase-PCR
5 (RT-PCR). Samples of transgenic gene-expressing tissue may also be evaluated immunocytochemically or immunohistochemically using antibodies specific for the transgene product.

Once the founder animals are produced, they may be bred, inbred, outbred, or crossbred to produce colonies of the particular animal. Examples of such breeding
10 strategies include, but are not limited to: outbreeding of founder animals with more than one integration site in order to establish separate lines; inbreeding of separate lines in order to produce compound transgenics that express the transgene at higher levels because of the effects of additive expression of each transgene; crossing of heterozygous transgenic animals to produce animals homozygous for a given integration site in order to
15 both augment expression and eliminate the need for screening of animals by DNA analysis; crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; and breeding to place the transgene on a distinct background that is appropriate for an experimental model of interest.

Transgenic animals of the invention have uses which include, but are not limited to,
20 to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

Methods for creating a transgenic animal with a disruption of a targeted gene are
25 also well-known in the art. In general, a vector is designed to comprise some nucleotide sequences homologous to the endogenous targeted gene. The vector is introduced into a cell so that it may integrate, via homologous recombination with chromosomal sequences, into the endogenous gene, thereby disrupting the function of the endogenous gene. The transgene may also be selectively introduced into a particular cell type, thus
30 inactivating the endogenous gene in only that cell type. *See, e.g., Gu et al., Science 265: 103-106 (1994).* The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. *See, e.g., Smithies et al., Nature 317: 230-234 (1985); Thomas et al., Cell 51: 503-512 (1987); Thompson et al., Cell 5: 313-321 (1989).*

In one embodiment, a mutant, non-functional nucleic acid molecule of the invention (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous nucleic acid sequence (either the coding regions or regulatory regions of the gene) can be used, with or without a selectable marker and/or a negative selectable
5 marker, to transfect cells that express polypeptides of the invention *in vivo*. In another embodiment, techniques known in the art are used to generate knockouts in cells that contain, but do not express the gene of interest. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the targeted gene. Such approaches are particularly suited in research and agricultural fields where modifications
10 to embryonic stem cells can be used to generate animal offspring with an inactive targeted gene. *See, e.g.,* Thomas, *supra* and Thompson, *supra*. However this approach can be routinely adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors that will be apparent to those of skill in the art.

15 In further embodiments of the invention, cells that are genetically engineered to express the polypeptides of the invention, or alternatively, that are genetically engineered not to express the polypeptides of the invention (*e.g.*, knockouts) are administered to a patient *in vivo*. Such cells may be obtained from an animal or patient or an MHC compatible donor and can include, but are not limited to fibroblasts, bone marrow cells,
20 blood cells (*e.g.*, lymphocytes), adipocytes, muscle cells, endothelial cells etc. The cells are genetically engineered *in vitro* using recombinant DNA techniques to introduce the coding sequence of polypeptides of the invention into the cells, or alternatively, to disrupt the coding sequence and/or endogenous regulatory sequence associated with the polypeptides of the invention, *e.g.*, by transduction (using viral vectors, and preferably
25 vectors that integrate the transgene into the cell genome) or transfection procedures, including, but not limited to, the use of plasmids, cosmids, YACs, naked DNA, electroporation, liposomes, etc.

The coding sequence of the polypeptides of the invention can be placed under the control of a strong constitutive or inducible promoter or promoter/enhancer to achieve
30 expression, and preferably secretion, of the polypeptides of the invention. The engineered cells which express and preferably secrete the polypeptides of the invention can be introduced into the patient systemically, *e.g.*, in the circulation, or intraperitoneally.

Alternatively, the cells can be incorporated into a matrix and implanted in the body, *e.g.*, genetically engineered fibroblasts can be implanted as part of a skin graft;

genetically engineered endothelial cells can be implanted as part of a lymphatic or vascular graft. *See, e.g.*, U.S. Patents 5,399,349 and 5,460,959, each of which is incorporated by reference herein in its entirety.

When the cells to be administered are non-autologous or non-MHC compatible
5 cells, they can be administered using well-known techniques which prevent the development of a host immune response against the introduced cells. For example, the cells may be introduced in an encapsulated form which, while allowing for an exchange of components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

10 Transgenic and "knock-out" animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

15 Computer Readable Means

A further aspect of the invention relates to a computer readable means for storing the nucleic acid and amino acid sequences of the instant invention. In a preferred embodiment, the invention provides a computer readable means for storing SEQ ID NO:
20 1 through 171 and SEQ ID NO: 172 through 295 as described herein, as the complete set of sequences or in any combination. The records of the computer readable means can be accessed for reading and display and for interface with a computer system for the application of programs allowing for the location of data upon a query for data meeting certain criteria, the comparison of sequences, the alignment or ordering of sequences meeting a set of criteria, and the like.

25 The nucleic acid and amino acid sequences of the invention are particularly useful as components in databases useful for search analyses as well as in sequence analysis algorithms. As used herein, the terms "nucleic acid sequences of the invention" and "amino acid sequences of the invention" mean any detectable chemical or physical characteristic of a polynucleotide or polypeptide of the invention that is or may be
30 reduced to or stored in a computer readable form. These include, without limitation, chromatographic scan data or peak data, photographic data or scan data therefrom, and mass spectrographic data.

This invention provides computer readable media having stored thereon sequences of the invention. A computer readable medium may comprise one or more of the following: a nucleic acid sequence comprising a sequence of a nucleic acid sequence of the invention; an amino acid sequence comprising an amino acid sequence of the invention; a set of nucleic acid sequences wherein at least one of said sequences comprises the sequence of a nucleic acid sequence of the invention; a set of amino acid sequences wherein at least one of said sequences comprises the sequence of an amino acid sequence of the invention; a data set representing a nucleic acid sequence comprising the sequence of one or more nucleic acid sequences of the invention; a data set representing a nucleic acid sequence encoding an amino acid sequence comprising the sequence of an amino acid sequence of the invention; a set of nucleic acid sequences wherein at least one of said sequences comprises the sequence of a nucleic acid sequence of the invention; a set of amino acid sequences wherein at least one of said sequences comprises the sequence of an amino acid sequence of the invention; a data set representing a nucleic acid sequence comprising the sequence of a nucleic acid sequence of the invention; a data set representing a nucleic acid sequence encoding an amino acid sequence comprising the sequence of an amino acid sequence of the invention. The computer readable medium can be any composition of matter used to store information or data, including, for example, commercially available floppy disks, tapes, hard drives, compact disks, and video disks.

Also provided by the invention are methods for the analysis of character sequences, particularly genetic sequences. Preferred methods of sequence analysis include, for example, methods of sequence homology analysis, such as identity and similarity analysis, RNA structure analysis, sequence assembly, cladistic analysis, sequence motif analysis, open reading frame determination, nucleic acid base calling, and sequencing chromatogram peak analysis.

A computer-based method is provided for performing nucleic acid sequence identity or similarity identification. This method comprises the steps of providing a nucleic acid sequence comprising the sequence of a nucleic acid of the invention in a computer readable medium; and comparing said nucleic acid sequence to at least one nucleic acid or amino acid sequence to identify sequence identity or similarity.

A computer-based method is also provided for performing amino acid homology identification, said method comprising the steps of: providing an amino acid sequence comprising the sequence of an amino acid of the invention in a computer readable

medium; and comparing said an amino acid sequence to at least one nucleic acid or an amino acid sequence to identify homology.

A computer-based method is still further provided for assembly of overlapping nucleic acid sequences into a single nucleic acid sequence, said method comprising the steps of: providing a first nucleic acid sequence comprising the sequence of a nucleic acid of the invention in a computer readable medium; and screening for at least one overlapping region between said first nucleic acid sequence and a second nucleic acid sequence.

Diagnostic Methods for Breast Cancer

10

The present invention also relates to quantitative and qualitative diagnostic assays and methods for detecting, diagnosing, monitoring, staging and predicting cancers by comparing expression of a BSNA or a BSP in a human patient that has or may have breast cancer, or who is at risk of developing breast cancer, with the expression of a BSNA or a BSP in a normal human control. For purposes of the present invention, "expression of a BSNA" or "BSNA expression" means the quantity of BSG mRNA that can be measured by any method known in the art or the level of transcription that can be measured by any method known in the art in a cell, tissue, organ or whole patient. Similarly, the term "expression of a BSP" or "BSP expression" means the amount of BSP that can be measured by any method known in the art or the level of translation of a BSG BSNA that can be measured by any method known in the art.

15
20

The present invention provides methods for diagnosing breast cancer in a patient, in particular squamous cell carcinoma, by analyzing for changes in levels of BSNA or BSP in cells, tissues, organs or bodily fluids compared with levels of BSNA or BSP in cells, tissues, organs or bodily fluids of preferably the same type from a normal human control, wherein an increase, or decrease in certain cases, in levels of a BSNA or BSP in the patient versus the normal human control is associated with the presence of breast cancer or with a predilection to the disease. In another preferred embodiment, the present invention provides methods for diagnosing breast cancer in a patient by analyzing changes in the structure of the mRNA of a BSG compared to the mRNA from a normal control. These changes include, without limitation, aberrant splicing, alterations in polyadenylation and/or alterations in 5' nucleotide capping. In yet another preferred embodiment, the present invention provides methods for diagnosing breast cancer in a patient by analyzing changes in a BSP compared to a BSP from a normal control. These

25

30

changes include, *e.g.*, alterations in glycosylation and/or phosphorylation of the BSP or subcellular BSP localization.

In a preferred embodiment, the expression of a BSNA is measured by determining the amount of an mRNA that encodes an amino acid sequence selected from SEQ ID NO: 172 through 295, a homolog, an allelic variant, or a fragment thereof. In a more preferred embodiment, the BSNA expression that is measured is the level of expression of a BSNA mRNA selected from SEQ ID NO: 1 through 171, or a hybridizing nucleic acid, homologous nucleic acid or allelic variant thereof, or a part of any of these nucleic acids. BSNA expression may be measured by any method known in the art, such as those described *supra*, including measuring mRNA expression by Northern blot, quantitative or qualitative reverse transcriptase PCR (RT-PCR), microarray, dot or slot blots or *in situ* hybridization. *See, e.g.*, Ausubel (1992), *supra*; Ausubel (1999), *supra*; Sambrook (1989), *supra*; and Sambrook (2001), *supra*. BSNA transcription may be measured by any method known in the art including using a reporter gene hooked up to the promoter of a BSG of interest or doing nuclear run-off assays. Alterations in mRNA structure, *e.g.*, aberrant splicing variants, may be determined by any method known in the art, including, RT-PCR followed by sequencing or restriction analysis. As necessary, BSNA expression may be compared to a known control, such as normal breast nucleic acid, to detect a change in expression.

In another preferred embodiment, the expression of a BSP is measured by determining the level of a BSP having an amino acid sequence selected from the group consisting of SEQ ID NO: 172 through 295, a homolog, an allelic variant, or a fragment thereof. Such levels are preferably determined in at least one of cells, tissues, organs and/or bodily fluids, including determination of normal and abnormal levels. Thus, for instance, a diagnostic assay in accordance with the invention for diagnosing over- or underexpression of BSNA or BSP compared to normal control bodily fluids, cells, or tissue samples may be used to diagnose the presence of breast cancer. The expression level of a BSP may be determined by any method known in the art, such as those described *supra*. In a preferred embodiment, the BSP expression level may be determined by radioimmunoassays, competitive-binding assays, ELISA, Western blot, FACS, immunohistochemistry, immunoprecipitation, proteomic approaches: two-dimensional gel electrophoresis (2D electrophoresis) and non-gel-based approaches such as mass spectrometry or protein interaction profiling. *See, e.g.*, Harlow (1999), *supra*; Ausubel (1992), *supra*; and Ausubel (1999), *supra*. Alterations in the BSP

-91-

structure may be determined by any method known in the art, including, *e.g.*, using antibodies that specifically recognize phosphoserine, phosphothreonine or phosphotyrosine residues, two-dimensional polyacrylamide gel electrophoresis (2D PAGE) and/or chemical analysis of amino acid residues of the protein. *Id.*

5 In a preferred embodiment, a radioimmunoassay (RIA) or an ELISA is used. An antibody specific to a BSP is prepared if one is not already available. In a preferred embodiment, the antibody is a monoclonal antibody. The anti-BSP antibody is bound to a solid support and any free protein binding sites on the solid support are blocked with a protein such as bovine serum albumin. A sample of interest is incubated with the
10 antibody on the solid support under conditions in which the BSP will bind to the anti-BSP antibody. The sample is removed, the solid support is washed to remove unbound material, and an anti-BSP antibody that is linked to a detectable reagent (a radioactive substance for RIA and an enzyme for ELISA) is added to the solid support and incubated under conditions in which binding of the BSP to the labeled antibody will occur. After
15 binding, the unbound labeled antibody is removed by washing. For an ELISA, one or more substrates are added to produce a colored reaction product that is based upon the amount of a BSP in the sample. For an RIA, the solid support is counted for radioactive decay signals by any method known in the art. Quantitative results for both RIA and ELISA typically are obtained by reference to a standard curve.

20 Other methods to measure BSP levels are known in the art. For instance, a competition assay may be employed wherein an anti-BSP antibody is attached to a solid support and an allocated amount of a labeled BSP and a sample of interest are incubated with the solid support. The amount of labeled BSP detected which is attached to the solid support can be correlated to the quantity of a BSP in the sample.

25 Of the proteomic approaches, 2D PAGE is a well-known technique. Isolation of individual proteins from a sample such as serum is accomplished using sequential separation of proteins by isoelectric point and molecular weight. Typically, polypeptides are first separated by isoelectric point (the first dimension) and then separated by size using an electric current (the second dimension). In general, the second dimension is
30 perpendicular to the first dimension. Because no two proteins with different sequences are identical on the basis of both size and charge, the result of 2D PAGE is a roughly square gel in which each protein occupies a unique spot. Analysis of the spots with chemical or antibody probes, or subsequent protein microsequencing can reveal the relative abundance of a given protein and the identity of the proteins in the sample.

-92-

Expression levels of a BSNA can be determined by any method known in the art, including PCR and other nucleic acid methods, such as ligase chain reaction (LCR) and nucleic acid sequence based amplification (NASBA), can be used to detect malignant cells for diagnosis and monitoring of various malignancies. For example,

5 reverse-transcriptase PCR (RT-PCR) is a powerful technique which can be used to detect the presence of a specific mRNA population in a complex mixture of thousands of other mRNA species. In RT-PCR, an mRNA species is first reverse transcribed to complementary DNA (cDNA) with use of the enzyme reverse transcriptase; the cDNA is then amplified as in a standard PCR reaction.

10 Hybridization to specific DNA molecules (*e.g.*, oligonucleotides) arrayed on a solid support can be used to both detect the expression of and quantitate the level of expression of one or more BSNAs of interest. In this approach, all or a portion of one or more BSNAs is fixed to a substrate. A sample of interest, which may comprise RNA, *e.g.*, total RNA or polyA-selected mRNA, or a complementary DNA (cDNA) copy of the
15 RNA is incubated with the solid support under conditions in which hybridization will occur between the DNA on the solid support and the nucleic acid molecules in the sample of interest. Hybridization between the substrate-bound DNA and the nucleic acid molecules in the sample can be detected and quantitated by several means, including, without limitation, radioactive labeling or fluorescent labeling of the nucleic acid
20 molecule or a secondary molecule designed to detect the hybrid.

The above tests can be carried out on samples derived from a variety of cells, bodily fluids and/or tissue extracts such as homogenates or solubilized tissue obtained from a patient. Tissue extracts are obtained routinely from tissue biopsy and autopsy material. Bodily fluids useful in the present invention include blood, urine, saliva or any
25 other bodily secretion or derivative thereof. By blood it is meant to include whole blood, plasma, serum or any derivative of blood. In a preferred embodiment, the specimen tested for expression of BSNA or BSP includes, without limitation, breast tissue, fluid obtained by bronchial alveolar lavage (BAL), sputum, breast cells grown in cell culture, blood, serum, lymph node tissue and lymphatic fluid. In another preferred embodiment,
30 especially when metastasis of a primary breast cancer is known or suspected, specimens include, without limitation, tissues from brain, bone, bone marrow, liver, adrenal glands and colon. In general, the tissues may be sampled by biopsy, including, without limitation, needle biopsy, *e.g.*, transthoracic needle aspiration, cervical mediastinoscopy, endoscopic lymph node biopsy, video-assisted thoracoscopy, exploratory thoracotomy,

bone marrow biopsy and bone marrow aspiration. See Scott, *supra* and Franklin, pp. 529-570, in Kane, *supra*. For early and inexpensive detection, assaying for changes in BSNA or BSPs in cells in sputum samples may be particularly useful. Methods of obtaining and analyzing sputum samples is disclosed in Franklin, *supra*.

- 5 All the methods of the present invention may optionally include determining the expression levels of one or more other cancer markers in addition to determining the expression level of a BSNA or BSP. In many cases, the use of another cancer marker will decrease the likelihood of false positives or false negatives. In one embodiment, the one or more other cancer markers include other BSNA or BSPs as disclosed herein.
- 10 Other cancer markers useful in the present invention will depend on the cancer being tested and are known to those of skill in the art. In a preferred embodiment, at least one other cancer marker in addition to a particular BSNA or BSP is measured. In a more preferred embodiment, at least two other additional cancer markers are used. In an even more preferred embodiment, at least three, more preferably at least five, even more
- 15 preferably at least ten additional cancer markers are used.

Diagnosing

- In one aspect, the invention provides a method for determining the expression levels and/or structural alterations of one or more BSNA and/or BSPs in a sample from a patient suspected of having breast cancer. In general, the method comprises the steps
- 20 of obtaining the sample from the patient, determining the expression level or structural alterations of a BSNA and/or BSP and then ascertaining whether the patient has breast cancer from the expression level of the BSNA or BSP. In general, if high expression relative to a control of a BSNA or BSP is indicative of breast cancer, a diagnostic assay is considered positive if the level of expression of the BSNA or BSP is at least two times
- 25 higher, and more preferably are at least five times higher, even more preferably at least ten times higher, than in preferably the same cells, tissues or bodily fluid of a normal human control. In contrast, if low expression relative to a control of a BSNA or BSP is indicative of breast cancer, a diagnostic assay is considered positive if the level of expression of the BSNA or BSP is at least two times lower, more preferably are at least
- 30 five times lower, even more preferably at least ten times lower than in preferably the same cells, tissues or bodily fluid of a normal human control. The normal human control may be from a different patient or from uninvolved tissue of the same patient.

-94-

The present invention also provides a method of determining whether breast cancer has metastasized in a patient. One may identify whether the breast cancer has metastasized by measuring the expression levels and/or structural alterations of one or more BSNA and/or BSPs in a variety of tissues. The presence of a BSNA or BSP in a certain tissue at levels higher than that of corresponding noncancerous tissue (*e.g.*, the same tissue from another individual) is indicative of metastasis if high level expression of a BSNA or BSP is associated with breast cancer. Similarly, the presence of a BSNA or BSP in a tissue at levels lower than that of corresponding noncancerous tissue is indicative of metastasis if low level expression of a BSNA or BSP is associated with breast cancer. Further, the presence of a structurally altered BSNA or BSP that is associated with breast cancer is also indicative of metastasis.

In general, if high expression relative to a control of a BSNA or BSP is indicative of metastasis, an assay for metastasis is considered positive if the level of expression of the BSNA or BSP is at least two times higher, and more preferably are at least five times higher, even more preferably at least ten times higher, than in preferably the same cells, tissues or bodily fluid of a normal human control. In contrast, if low expression relative to a control of a BSNA or BSP is indicative of metastasis, an assay for metastasis is considered positive if the level of expression of the BSNA or BSP is at least two times lower, more preferably are at least five times lower, even more preferably at least ten times lower than in preferably the same cells, tissues or bodily fluid of a normal human control.

The BSNA or BSP of this invention may be used as element in an array or a multi-analyte test to recognize expression patterns associated with breast cancers or other breast related disorders. In addition, the sequences of either the nucleic acids or proteins may be used as elements in a computer program for pattern recognition of breast disorders.

Staging

The invention also provides a method of staging breast cancer in a human patient. The method comprises identifying a human patient having breast cancer and analyzing cells, tissues or bodily fluids from such human patient for expression levels and/or structural alterations of one or more BSNA and/or BSPs. First, one or more tumors from a variety of patients are staged according to procedures well-known in the art, and the expression level of one or more BSNA and/or BSPs is determined for each stage to obtain a

-95-

standard expression level for each BSNA and BSP. Then, the BSNA or BSP expression levels are determined in a biological sample from a patient whose stage of cancer is not known. The BSNA or BSP expression levels from the patient are then compared to the standard expression level. By comparing the expression level of the BSNAs and BSPs
5 from the patient to the standard expression levels, one may determine the stage of the tumor. The same procedure may be followed using structural alterations of a BSNA or BSP to determine the stage of a breast cancer.

Monitoring

Further provided is a method of monitoring breast cancer in a human patient.
10 One may monitor a human patient to determine whether there has been metastasis and, if there has been, when metastasis began to occur. One may also monitor a human patient to determine whether a preneoplastic lesion has become cancerous. One may also monitor a human patient to determine whether a therapy, *e.g.*, chemotherapy, radiotherapy or surgery, has decreased or eliminated the breast cancer. The method
15 comprises identifying a human patient that one wants to monitor for breast cancer, periodically analyzing cells, tissues or bodily fluids from such human patient for expression levels of one or more BSNAs or BSPs, and comparing the BSNA or BSP levels over time to those BSNA or BSP expression levels obtained previously. Patients may also be monitored by measuring one or more structural alterations in a BSNA or
20 BSP that are associated with breast cancer.

If increased expression of a BSNA or BSP is associated with metastasis, treatment failure, or conversion of a preneoplastic lesion to a cancerous lesion, then detecting an increase in the expression level of a BSNA or BSP indicates that the tumor is metastasizing, that treatment has failed or that the lesion is cancerous, respectively.
25 One having ordinary skill in the art would recognize that if this were the case, then a decreased expression level would be indicative of no metastasis, effective therapy or failure to progress to a neoplastic lesion. If decreased expression of a BSNA or BSP is associated with metastasis, treatment failure, or conversion of a preneoplastic lesion to a cancerous lesion, then detecting an decrease in the expression level of a BSNA or BSP
30 indicates that the tumor is metastasizing, that treatment has failed or that the lesion is cancerous, respectively. In a preferred embodiment, the levels of BSNAs or BSPs are determined from the same cell type, tissue or bodily fluid as prior patient samples.

-96-

Monitoring a patient for onset of breast cancer metastasis is periodic and preferably is done on a quarterly basis, but may be done more or less frequently.

The methods described herein can further be utilized as prognostic assays to identify subjects having or at risk of developing a disease or disorder associated with increased or decreased expression levels of a BSNA and/or BSP. The present invention provides a method in which a test sample is obtained from a human patient and one or more BSNAs and/or BSPs are detected. The presence of higher (or lower) BSNA or BSP levels as compared to normal human controls is diagnostic for the human patient being at risk for developing cancer, particularly breast cancer. The effectiveness of therapeutic agents to decrease (or increase) expression or activity of one or more BSNAs and/or BSPs of the invention can also be monitored by analyzing levels of expression of the BSNAs and/or BSPs in a human patient in clinical trials or in *in vitro* screening assays such as in human cells. In this way, the gene expression pattern can serve as a marker, indicative of the physiological response of the human patient or cells, as the case may be, to the agent being tested.

Detection of Genetic Lesions or Mutations

The methods of the present invention can also be used to detect genetic lesions or mutations in a BSG, thereby determining if a human with the genetic lesion is susceptible to developing breast cancer or to determine what genetic lesions are responsible, or are partly responsible, for a person's existing breast cancer. Genetic lesions can be detected, for example, by ascertaining the existence of a deletion, insertion and/or substitution of one or more nucleotides from the BSGs of this invention, a chromosomal rearrangement of BSG, an aberrant modification of BSG (such as of the methylation pattern of the genomic DNA), or allelic loss of a BSG. Methods to detect such lesions in the BSG of this invention are known to those having ordinary skill in the art following the teachings of the specification.

Methods of Detecting Noncancerous Breast Diseases

The invention also provides a method for determining the expression levels and/or structural alterations of one or more BSNAs and/or BSPs in a sample from a patient suspected of having or known to have a noncancerous breast disease. In general, the method comprises the steps of obtaining a sample from the patient, determining the expression level or structural alterations of a BSNA and/or BSP, comparing the

-97-

expression level or structural alteration of the BSNA or BSP to a normal breast control, and then ascertaining whether the patient has a noncancerous breast disease. In general, if high expression relative to a control of a BSNA or BSP is indicative of a particular noncancerous breast disease, a diagnostic assay is considered positive if the level of
5 expression of the BSNA or BSP is at least two times higher, and more preferably are at least five times higher, even more preferably at least ten times higher, than in preferably the same cells, tissues or bodily fluid of a normal human control. In contrast, if low expression relative to a control of a BSNA or BSP is indicative of a noncancerous breast
10 or BSP is at least two times lower, more preferably are at least five times lower, even more preferably at least ten times lower than in preferably the same cells, tissues or bodily fluid of a normal human control. The normal human control may be from a different patient or from uninvolved tissue of the same patient.

One having ordinary skill in the art may determine whether a BSNA and/or BSP
15 is associated with a particular noncancerous breast disease by obtaining breast tissue from a patient having a noncancerous breast disease of interest and determining which BSNAs and/or BSPs are expressed in the tissue at either a higher or a lower level than in normal breast tissue. In another embodiment, one may determine whether a BSNA or BSP exhibits structural alterations in a particular noncancerous breast disease state by
20 obtaining breast tissue from a patient having a noncancerous breast disease of interest and determining the structural alterations in one or more BSNAs and/or BSPs relative to normal breast tissue.

25 Methods for Identifying Breast Tissue

In another aspect, the invention provides methods for identifying breast tissue. These methods are particularly useful in, *e.g.*, forensic science, breast cell differentiation and development, and in tissue engineering.

In one embodiment, the invention provides a method for determining whether a
30 sample is breast tissue or has breast tissue-like characteristics. The method comprises the steps of providing a sample suspected of comprising breast tissue or having breast tissue-like characteristics, determining whether the sample expresses one or more BSNAs and/or BSPs, and, if the sample expresses one or more BSNAs and/or BSPs, concluding that the sample comprises breast tissue. In a preferred embodiment, the BSNA encodes a
35 polypeptide having an amino acid sequence selected from SEQ ID NO: 172 through 295,

-98-

- or a homolog, allelic variant or fragment thereof. In a more preferred embodiment, the BSNA has a nucleotide sequence selected from SEQ ID NO: 1 through 171, or a hybridizing nucleic acid, an allelic variant or a part thereof. Determining whether a sample expresses a BSNA can be accomplished by any method known in the art.
- 5 Preferred methods include hybridization to microarrays, Northern blot hybridization, and quantitative or qualitative RT-PCR. In another preferred embodiment, the method can be practiced by determining whether a BSP is expressed. Determining whether a sample expresses a BSP can be accomplished by any method known in the art. Preferred methods include Western blot, ELISA, RIA and 2D PAGE. In one embodiment, the BSP
- 10 has an amino acid sequence selected from SEQ ID NO: 172 through 295, or a homolog, allelic variant or fragment thereof. In another preferred embodiment, the expression of at least two BSNA and/or BSPs is determined. In a more preferred embodiment, the expression of at least three, more preferably four and even more preferably five BSNA and/or BSPs are determined.
- 15 In one embodiment, the method can be used to determine whether an unknown tissue is breast tissue. This is particularly useful in forensic science, in which small, damaged pieces of tissues that are not identifiable by microscopic or other means are recovered from a crime or accident scene. In another embodiment, the method can be used to determine whether a tissue is differentiating or developing into breast tissue.
- 20 This is important in monitoring the effects of the addition of various agents to cell or tissue culture, *e.g.*, in producing new breast tissue by tissue engineering. These agents include, *e.g.*, growth and differentiation factors, extracellular matrix proteins and culture medium. Other factors that may be measured for effects on tissue development and differentiation include gene transfer into the cells or tissues, alterations in pH,
- 25 aqueous:air interface and various other culture conditions.

Methods for Producing and Modifying Breast Tissue

- In another aspect, the invention provides methods for producing engineered breast tissue or cells. In one embodiment, the method comprises the steps of providing cells,
- 30 introducing a BSNA or a BSG into the cells, and growing the cells under conditions in which they exhibit one or more properties of breast tissue cells. In a preferred embodiment, the cells are pluripotent. As is well-known in the art, normal breast tissue comprises a large number of different cell types. Thus, in one embodiment, the engineered breast tissue or cells comprises one of these cell types. In another

embodiment, the engineered breast tissue or cells comprises more than one breast cell type. Further, the culture conditions of the cells or tissue may require manipulation in order to achieve full differentiation and development of the breast cell tissue. Methods for manipulating culture conditions are well-known in the art.

5 Nucleic acid molecules encoding one or more BSPs are introduced into cells, preferably pluripotent cells. In a preferred embodiment, the nucleic acid molecules encode BSPs having amino acid sequences selected from SEQ ID NO: 172 through 295, or homologous proteins, analogs, allelic variants or fragments thereof. In a more preferred embodiment, the nucleic acid molecules have a nucleotide sequence selected
10 from SEQ ID NO: 1 through 171, or hybridizing nucleic acids, allelic variants or parts thereof. In another highly preferred embodiment, a BSG is introduced into the cells. Expression vectors and methods of introducing nucleic acid molecules into cells are well-known in the art and are described in detail, *supra*.

15 Artificial breast tissue may be used to treat patients who have lost some or all of their breast function.

Pharmaceutical Compositions

In another aspect, the invention provides pharmaceutical compositions comprising the nucleic acid molecules, polypeptides, antibodies, antibody derivatives,
20 antibody fragments, agonists, antagonists, and inhibitors of the present invention. In a preferred embodiment, the pharmaceutical composition comprises a BSNA or part thereof. In a more preferred embodiment, the BSNA has a nucleotide sequence selected from the group consisting of SEQ ID NO: 1 through 171, a nucleic acid that hybridizes thereto, an allelic variant thereof, or a nucleic acid that has substantial sequence identity
25 thereto. In another preferred embodiment, the pharmaceutical composition comprises a BSP or fragment thereof. In a more preferred embodiment, the BSP having an amino acid sequence that is selected from the group consisting of SEQ ID NO: 172 through 295, a polypeptide that is homologous thereto, a fusion protein comprising all or a portion of the polypeptide, or an analog or derivative thereof. In another preferred embodiment, the
30 pharmaceutical composition comprises an anti-BSP antibody, preferably an antibody that specifically binds to a BSP having an amino acid that is selected from the group consisting of SEQ ID NO: 172 through 295, or an antibody that binds to a polypeptide that is homologous thereto, a fusion protein comprising all or a portion of the polypeptide, or an analog or derivative thereof.

-100-

Such a composition typically contains from about 0.1 to 90% by weight of a therapeutic agent of the invention formulated in and/or with a pharmaceutically acceptable carrier or excipient.

Pharmaceutical formulation is a well-established art, and is further described in
5 Gennaro (ed.), Remington: The Science and Practice of Pharmacy, 20th ed., Lippincott, Williams & Wilkins (2000); Ansel *et al.*, Pharmaceutical Dosage Forms and Drug Delivery Systems, 7th ed., Lippincott Williams & Wilkins (1999); and Kibbe (ed.), Handbook of Pharmaceutical Excipients American Pharmaceutical Association, 3rd ed. (2000), the disclosures of which are incorporated herein by reference in their entireties,
10 and thus need not be described in detail herein.

Briefly, formulation of the pharmaceutical compositions of the present invention will depend upon the route chosen for administration. The pharmaceutical compositions utilized in this invention can be administered by various routes including both enteral and
15 parenteral routes, including oral, intravenous, intramuscular, subcutaneous, inhalation, topical, sublingual, rectal, intra-arterial, intramedullary, intrathecal, intraventricular, transmucosal, transdermal, intranasal, intraperitoneal, intrapulmonary, and intrauterine.

Oral dosage forms can be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Solid formulations of the compositions for oral administration can contain
20 suitable carriers or excipients, such as carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, or microcrystalline cellulose; gums including arabic and tragacanth; proteins such as gelatin and collagen; inorganics, such as kaolin, calcium
25 carbonate, dicalcium phosphate, sodium chloride; and other agents such as acacia and alginic acid.

Agents that facilitate disintegration and/or solubilization can be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate, microcrystalline cellulose, corn starch, sodium starch glycolate, and
30 alginic acid.

Tablet binders that can be used include acacia, methylcellulose, sodium carboxymethylcellulose, polyvinylpyrrolidone (Povidone™), hydroxypropyl methylcellulose, sucrose, starch and ethylcellulose.

-101-

Lubricants that can be used include magnesium stearates, stearic acid, silicone fluid, talc, waxes, oils, and colloidal silica.

Fillers, agents that facilitate disintegration and/or solubilization, tablet binders and lubricants, including the aforementioned, can be used singly or in combination.

5 Solid oral dosage forms need not be uniform throughout. For example, dragee cores can be used in conjunction with suitable coatings, such as concentrated sugar solutions, which can also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures.

10 Oral dosage forms of the present invention include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds can be dissolved or
15 suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Additionally, dyestuffs or pigments can be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, *i.e.*, dosage.

Liquid formulations of the pharmaceutical compositions for oral (enteral)
20 administration are prepared in water or other aqueous vehicles and can contain various suspending agents such as methylcellulose, alginates, tragacanth, pectin, kelgin, carrageenan, acacia, polyvinylpyrrolidone, and polyvinyl alcohol. The liquid formulations can also include solutions, emulsions, syrups and elixirs containing, together with the active compound(s), wetting agents, sweeteners, and coloring and
25 flavoring agents.

The pharmaceutical compositions of the present invention can also be formulated for parenteral administration. Formulations for parenteral administration can be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions.

For intravenous injection, water soluble versions of the compounds of the present
30 invention are formulated in, or if provided as a lyophilate, mixed with, a physiologically acceptable fluid vehicle, such as 5% dextrose ("D5"), physiologically buffered saline, 0.9% saline, Hanks' solution, or Ringer's solution. Intravenous formulations may include carriers, excipients or stabilizers including, without limitation, calcium, human serum albumin, citrate, acetate, calcium chloride, carbonate, and other salts.

-102-

Intramuscular preparations, *e.g.* a sterile formulation of a suitable soluble salt form of the compounds of the present invention, can be dissolved and administered in a pharmaceutical excipient such as Water-for-Injection, 0.9% saline, or 5% glucose solution. Alternatively, a suitable insoluble form of the compound can be prepared and
5 administered as a suspension in an aqueous base or a pharmaceutically acceptable oil base, such as an ester of a long chain fatty acid (*e.g.*, ethyl oleate), fatty oils such as sesame oil, triglycerides, or liposomes.

Parenteral formulations of the compositions can contain various carriers such as vegetable oils, dimethylacetamide, dimethylformamide, ethyl lactate, ethyl carbonate,
10 isopropyl myristate, ethanol, polyols (glycerol, propylene glycol, liquid polyethylene glycol, and the like).

Aqueous injection suspensions can also contain substances that increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Non-lipid polycationic amino polymers can also be used for delivery. Optionally, the
15 suspension can also contain suitable stabilizers or agents that increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

Pharmaceutical compositions of the present invention can also be formulated to permit injectable, long-term, deposition. Injectable depot forms may be made by forming microencapsulated matrices of the compound in biodegradable polymers such as
20 polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in microemulsions that are compatible with body tissues.

25 The pharmaceutical compositions of the present invention can be administered topically.

For topical use the compounds of the present invention can also be prepared in suitable forms to be applied to the skin, or mucus membranes of the nose and throat, and can take the form of lotions, creams, ointments, liquid sprays or inhalants, drops,
30 tinctures, lozenges, or throat paints. Such topical formulations further can include chemical compounds such as dimethylsulfoxide (DMSO) to facilitate surface penetration of the active ingredient. In other transdermal formulations, typically in patch-delivered formulations, the pharmaceutically active compound is formulated with one or more skin penetrants, such as 2-N-methyl-pyrrolidone (NMP) or Azone. A topical semi-solid

-103-

ointment formulation typically contains a concentration of the active ingredient from about 1 to 20%, *e.g.*, 5 to 10%, in a carrier such as a pharmaceutical cream base.

For application to the eyes or ears, the compounds of the present invention can be presented in liquid or semi-liquid form formulated in hydrophobic or hydrophilic bases
5 as ointments, creams, lotions, paints or powders.

For rectal administration the compounds of the present invention can be administered in the form of suppositories admixed with conventional carriers such as cocoa butter, wax or other glyceride.

Inhalation formulations can also readily be formulated. For inhalation, various
10 powder and liquid formulations can be prepared. For aerosol preparations, a sterile formulation of the compound or salt form of the compound may be used in inhalers, such as metered dose inhalers, and nebulizers. Aerosolized forms may be especially useful for treating respiratory disorders.

Alternatively, the compounds of the present invention can be in powder form for
15 reconstitution in the appropriate pharmaceutically acceptable carrier at the time of delivery.

The pharmaceutically active compound in the pharmaceutical compositions of the present invention can be provided as the salt of a variety of acids, including but not limited to hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acid. Salts
20 tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms.

After pharmaceutical compositions have been prepared, they are packaged in an appropriate container and labeled for treatment of an indicated condition.

The active compound will be present in an amount effective to achieve the
25 intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

A "therapeutically effective dose" refers to that amount of active ingredient, for example BSP polypeptide, fusion protein, or fragments thereof, antibodies specific for BSP, agonists, antagonists or inhibitors of BSP, which ameliorates the signs or symptoms
30 of the disease or prevents progression thereof; as would be understood in the medical arts, cure, although desired, is not required.

The therapeutically effective dose of the pharmaceutical agents of the present invention can be estimated initially by *in vitro* tests, such as cell culture assays, followed by assay in model animals, usually mice, rats, rabbits, dogs, or pigs. The animal model

can also be used to determine an initial preferred concentration range and route of administration.

For example, the ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population) can be determined in one or more cell culture of animal model systems. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as LD50/ED50. Pharmaceutical compositions that exhibit large therapeutic indices are preferred.

The data obtained from cell culture assays and animal studies are used in formulating an initial dosage range for human use, and preferably provide a range of circulating concentrations that includes the ED50 with little or no toxicity. After administration, or between successive administrations, the circulating concentration of active agent varies within this range depending upon pharmacokinetic factors well-known in the art, such as the dosage form employed, sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors specific to the subject requiring treatment. Factors that can be taken into account by the practitioner include the severity of the disease state, general health of the subject, age, weight, gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions can be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from 0.1 to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Where the therapeutic agent is a protein or antibody of the present invention, the therapeutic protein or antibody agent typically is administered at a daily dosage of 0.01 mg to 30 mg/kg of body weight of the patient (*e.g.*, 1 mg/kg to 5 mg/kg). The pharmaceutical formulation can be administered in multiple doses per day, if desired, to achieve the total desired daily dose.

Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

Conventional methods, known to those of ordinary skill in the art of medicine, can be used to administer the pharmaceutical formulation(s) of the present invention to

the patient. The pharmaceutical compositions of the present invention can be administered alone, or in combination with other therapeutic agents or interventions.

Therapeutic Methods

5 The present invention further provides methods of treating subjects having defects in a gene of the invention, *e.g.*, in expression, activity, distribution, localization, and/or solubility, which can manifest as a disorder of breast function. As used herein, “treating” includes all medically-acceptable types of therapeutic intervention, including palliation and prophylaxis (prevention) of disease. The term “treating” encompasses any
10 improvement of a disease, including minor improvements. These methods are discussed below.

Gene Therapy and Vaccines

 The isolated nucleic acids of the present invention can also be used to drive *in vivo* expression of the polypeptides of the present invention. *In vivo* expression can be
15 driven from a vector, typically a viral vector, often a vector based upon a replication incompetent retrovirus, an adenovirus, or an adeno-associated virus (AAV), for purpose of gene therapy. *In vivo* expression can also be driven from signals endogenous to the nucleic acid or from a vector, often a plasmid vector, such as pVAX1 (Invitrogen, Carlsbad, CA, USA), for purpose of “naked” nucleic acid vaccination, as further
20 described in U.S. Patents 5,589,466; 5,679,647; 5,804,566; 5,830,877; 5,843,913; 5,880,104; 5,958,891; 5,985,847; 6,017,897; 6,110,898; and 6,204,250, the disclosures of which are incorporated herein by reference in their entireties. For cancer therapy, it is preferred that the vector also be tumor-selective. *See, e.g., Doronin et al., J. Virol. 75: 3314-24 (2001).*

25 In another embodiment of the therapeutic methods of the present invention, a therapeutically effective amount of a pharmaceutical composition comprising a nucleic acid of the present invention is administered. The nucleic acid can be delivered in a vector that drives expression of a BSP, fusion protein, or fragment thereof, or without such vector. Nucleic acid compositions that can drive expression of a BSP are
30 administered, for example, to complement a deficiency in the native BSP, or as DNA vaccines. Expression vectors derived from virus, replication deficient retroviruses, adenovirus, adeno-associated (AAV) virus, herpes virus, or vaccinia virus can be used as can plasmids. *See, e.g., Cid-Arregui, supra.* In a preferred embodiment, the nucleic acid

-106-

molecule encodes a BSP having the amino acid sequence of SEQ ID NO: 172 through 295, or a fragment, fusion protein, allelic variant or homolog thereof.

In still other therapeutic methods of the present invention, pharmaceutical compositions comprising host cells that express a BSP, fusions, or fragments thereof can be administered. In such cases, the cells are typically autologous, so as to circumvent xenogeneic or allotypic rejection, and are administered to complement defects in BSP production or activity. In a preferred embodiment, the nucleic acid molecules in the cells encode a BSP having the amino acid sequence of SEQ ID NO: 172 through 295, or a fragment, fusion protein, allelic variant or homolog thereof.

10 *Antisense Administration*

Antisense nucleic acid compositions, or vectors that drive expression of a BSG antisense nucleic acid, are administered to downregulate transcription and/or translation of a BSG in circumstances in which excessive production, or production of aberrant protein, is the pathophysiologic basis of disease.

15 Antisense compositions useful in therapy can have a sequence that is complementary to coding or to noncoding regions of a BSG. For example, oligonucleotides derived from the transcription initiation site, *e.g.*, between positions -10 and +10 from the start site, are preferred.

Catalytic antisense compositions, such as ribozymes, that are capable of sequence-specific hybridization to BSG transcripts, are also useful in therapy. *See, e.g.*, Phylactou, *Adv. Drug Deliv. Rev.* 44(2-3): 97-108 (2000); Phylactou *et al.*, *Hum. Mol. Genet.* 7(10): 1649-53 (1998); Rossi, *Ciba Found. Symp.* 209: 195-204 (1997); and Sigurdsson *et al.*, *Trends Biotechnol.* 13(8): 286-9 (1995), the disclosures of which are incorporated herein by reference in their entireties.

25 Other nucleic acids useful in the therapeutic methods of the present invention are those that are capable of triplex helix formation in or near the BSG genomic locus. Such triplexing oligonucleotides are able to inhibit transcription. *See, e.g.*, Intody *et al.*, *Nucleic Acids Res.* 28(21): 4283-90 (2000); McGuffie *et al.*, *Cancer Res.* 60(14): 3790-9 (2000), the disclosures of which are incorporated herein by reference. Pharmaceutical compositions comprising such triplex forming oligos (TFOs) are administered in circumstances in which excessive production, or production of aberrant protein, is a pathophysiologic basis of disease.

-107-

In a preferred embodiment, the antisense molecule is derived from a nucleic acid molecule encoding a BSP, preferably a BSP comprising an amino acid sequence of SEQ ID NO: 172 through 295, or a fragment, allelic variant or homolog thereof. In a more preferred embodiment, the antisense molecule is derived from a nucleic acid molecule
5 having a nucleotide sequence of SEQ ID NO: 1 through 171, or a part, allelic variant, substantially similar or hybridizing nucleic acid thereof.

Polypeptide Administration

In one embodiment of the therapeutic methods of the present invention, a therapeutically effective amount of a pharmaceutical composition comprising a BSP, a
10 fusion protein, fragment, analog or derivative thereof is administered to a subject with a clinically-significant BSP defect.

Protein compositions are administered, for example, to complement a deficiency in native BSP. In other embodiments, protein compositions are administered as a vaccine to elicit a humoral and/or cellular immune response to BSP. The immune response can
15 be used to modulate activity of BSP or, depending on the immunogen, to immunize against aberrant or aberrantly expressed forms, such as mutant or inappropriately expressed isoforms. In yet other embodiments, protein fusions having a toxic moiety are administered to ablate cells that aberrantly accumulate BSP.

In a preferred embodiment, the polypeptide is a BSP comprising an amino acid
20 sequence of SEQ ID NO: 172 through 295, or a fusion protein, allelic variant, homolog, analog or derivative thereof. In a more preferred embodiment, the polypeptide is encoded by a nucleic acid molecule having a nucleotide sequence of SEQ ID NO: 1 through 171, or a part, allelic variant, substantially similar or hybridizing nucleic acid thereof.

Antibody, Agonist and Antagonist Administration

In another embodiment of the therapeutic methods of the present invention, a therapeutically effective amount of a pharmaceutical composition comprising an antibody (including fragment or derivative thereof) of the present invention is administered. As is well-known, antibody compositions are administered, for example,
30 to antagonize activity of BSP, or to target therapeutic agents to sites of BSP presence and/or accumulation. In a preferred embodiment, the antibody specifically binds to a BSP comprising an amino acid sequence of SEQ ID NO: 172 through 295, or a fusion protein, allelic variant, homolog, analog or derivative thereof. In a more preferred

-108-

embodiment, the antibody specifically binds to a BSP encoded by a nucleic acid molecule having a nucleotide sequence of SEQ ID NO: 1 through 171, or a part, allelic variant, substantially similar or hybridizing nucleic acid thereof.

The present invention also provides methods for identifying modulators which
5 bind to a BSP or have a modulatory effect on the expression or activity of a BSP. Modulators which decrease the expression or activity of BSP (antagonists) are believed to be useful in treating breast cancer. Such screening assays are known to those of skill in the art and include, without limitation, cell-based assays and cell-free assays. Small molecules predicted via computer imaging to specifically bind to regions of a BSP can
10 also be designed, synthesized and tested for use in the imaging and treatment of breast cancer. Further, libraries of molecules can be screened for potential anticancer agents by assessing the ability of the molecule to bind to the BSPs identified herein. Molecules identified in the library as being capable of binding to a BSP are key candidates for further evaluation for use in the treatment of breast cancer. In a preferred embodiment,
15 these molecules will downregulate expression and/or activity of a BSP in cells.

In another embodiment of the therapeutic methods of the present invention, a pharmaceutical composition comprising a non-antibody antagonist of BSP is administered. Antagonists of BSP can be produced using methods generally known in the art. In particular, purified BSP can be used to screen libraries of pharmaceutical
20 agents, often combinatorial libraries of small molecules, to identify those that specifically bind and antagonize at least one activity of a BSP.

In other embodiments a pharmaceutical composition comprising an agonist of a BSP is administered. Agonists can be identified using methods analogous to those used to identify antagonists.

25 In a preferred embodiment, the antagonist or agonist specifically binds to and antagonizes or agonizes, respectively, a BSP comprising an amino acid sequence of SEQ ID NO: 172 through 295, or a fusion protein, allelic variant, homolog, analog or derivative thereof. In a more preferred embodiment, the antagonist or agonist specifically binds to and antagonizes or agonizes, respectively, a BSP encoded by a
30 nucleic acid molecule having a nucleotide sequence of SEQ ID NO: 1 through 171, or a part, allelic variant, substantially similar or hybridizing nucleic acid thereof.

Targeting Breast Tissue

The invention also provides a method in which a polypeptide of the invention, or an antibody thereto, is linked to a therapeutic agent such that it can be delivered to the

-109-

breast or to specific cells in the breast. In a preferred embodiment, an anti-BSP antibody is linked to a therapeutic agent and is administered to a patient in need of such therapeutic agent. The therapeutic agent may be a toxin, if breast tissue needs to be selectively destroyed. This would be useful for targeting and killing breast cancer cells.

- 5 In another embodiment, the therapeutic agent may be a growth or differentiation factor, which would be useful for promoting breast cell function.

In another embodiment, an anti-BSP antibody may be linked to an imaging agent that can be detected using, *e.g.*, magnetic resonance imaging, CT or PET. This would be useful for determining and monitoring breast function, identifying breast cancer tumors,
10 and identifying noncancerous breast diseases.

EXAMPLES

Example 1: Gene Expression analysis

BSGs were identified by mRNA subtraction analysis using standard methods. The sequences were extended using GeneBank sequences, Incyte's proprietary database.

- 15 From the nucleotide sequences, predicted amino acid sequences were prepared.
DEX0306_1, DEX0306_2 correspond to SEQ ID NO.1, 2 etc. DEX0157 was the parent sequence found in the mRNA subtractions.

	DEX0306_1	DEX0157_1	DEX0306_172
	DEX0306_2	flex DEX0157_1	
20	DEX0306_3	DEX0157_2	DEX0306_173
	DEX0306_4	flex DEX0157_2	
	DEX0306_5	DEX0157_3	DEX0306_174
	DEX0306_6	flex DEX0157_3	
	DEX0306_7	DEX0157_4	DEX0306_175
25	DEX0306_8	flex DEX0157_4	
	DEX0306_9	DEX0157_5	DEX0306_176
	DEX0306_10	flex DEX0157_5	
	DEX0306_11	DEX0157_6	DEX0306_177
	DEX0306_12	flex DEX0157_6	
30	DEX0306_13	DEX0157_7	DEX0306_178
	DEX0306_14	DEX0157_8	DEX0306_179
	DEX0306_15	DEX0157_9	DEX0306_180
	DEX0306_16	flex DEX0157_9	
	DEX0306_17	DEX0157_10	DEX0306_181
35	DEX0306_18	flex DEX0157_10	DEX0306_182
	DEX0306_19	DEX0157_11	DEX0306_183
	DEX0306_20	flex DEX0157_11	
	DEX0306_21	DEX0157_12	DEX0306_184
	DEX0306_22	flex DEX0157_12	
40	DEX0306_23	DEX0157_13	DEX0306_185
	DEX0306_24	flex DEX0157_13	
	DEX0306_25	DEX0157_14	DEX0306_186
	DEX0306_26	flex DEX0157_14	
	DEX0306_27	DEX0157_15	DEX0306_187
45	DEX0306_28	flex DEX0157_15	
	DEX0306_29	DEX0157_16	DEX0306_188

-110-

DEX0306_30 DEX0157_17 DEX0306_189
DEX0306_31 flex DEX0157_17 DEX0306_190
DEX0306_32 DEX0157_18 DEX0306_191
DEX0306_33 flex DEX0157_18
5 DEX0306_34 DEX0157_19 DEX0306_192
DEX0306_35 DEX0157_20 DEX0306_193
DEX0306_36 flex DEX0157_20 DEX0306_194
DEX0306_37 DEX0157_21
DEX0306_38 DEX0157_22 DEX0306_195
10 DEX0306_39 flex DEX0157_22
DEX0306_40 DEX0157_23 DEX0306_196
DEX0306_41 flex DEX0157_23
DEX0306_42 DEX0157_24 DEX0306_197
DEX0306_43 DEX0157_25 DEX0306_198
15 DEX0306_44 flex DEX0157_25 DEX0306_199
DEX0306_45 DEX0157_26 DEX0306_200
DEX0306_46 DEX0157_27 DEX0306_201
DEX0306_47 flex DEX0157_27
DEX0306_48 DEX0157_28 DEX0306_202
20 DEX0306_49 flex DEX0157_28
DEX0306_50 DEX0157_29 DEX0306_203
DEX0306_51 flex DEX0157_29
DEX0306_52 DEX0157_30 DEX0306_204
DEX0306_53 flex DEX0157_30 DEX0306_205
25 DEX0306_54 DEX0157_31 DEX0306_206
DEX0306_55 flex DEX0157_31
DEX0306_56 DEX0157_32 DEX0306_207
DEX0306_57 flex DEX0157_32
DEX0306_58 DEX0157_33 DEX0306_208
30 DEX0306_59 flex DEX0157_33
DEX0306_60 DEX0157_34
DEX0306_61 flex DEX0157_34
DEX0306_62 DEX0157_35 DEX0306_209
DEX0306_63 DEX0157_36 DEX0306_210
35 DEX0306_64 flex DEX0157_36
DEX0306_65 DEX0157_37 DEX0306_211
DEX0306_66 flex DEX0157_37 DEX0306_212
DEX0306_67 DEX0157_38 DEX0306_213
DEX0306_68 DEX0157_39 DEX0306_214
40 DEX0306_69 flex DEX0157_39 DEX0306_215
DEX0306_70 DEX0157_40 DEX0306_216
DEX0306_71 flex DEX0157_40 DEX0306_217
DEX0306_72 DEX0157_41 DEX0306_218
DEX0306_73 flex DEX0157_41 DEX0306_219
45 DEX0306_74 DEX0157_42 DEX0306_220
DEX0306_75 flex DEX0157_42
DEX0306_76 DEX0157_43 DEX0306_221
DEX0306_77 flex DEX0157_43
DEX0306_78 DEX0157_44 DEX0306_222
50 DEX0306_79 flex DEX0157_44
DEX0306_80 DEX0157_45 DEX0306_223
DEX0306_81 flex DEX0157_45 DEX0306_224
DEX0306_82 DEX0157_46 DEX0306_225
DEX0306_83 DEX0157_47 DEX0306_226
55 DEX0306_84 DEX0157_48 DEX0306_227
DEX0306_85 DEX0157_49 DEX0306_228
DEX0306_86 flex DEX0157_49 DEX0306_229
DEX0306_87 DEX0157_50 DEX0306_230
DEX0306_88 flex DEX0157_50 DEX0306_231
60 DEX0306_89 DEX0157_51 DEX0306_232
DEX0306_90 flex DEX0157_51
DEX0306_91 DEX0157_52 DEX0306_233

-111-

DEX0306_92 DEX0157_53 DEX0306_234
DEX0306_93 flex DEX0157_53 DEX0306_235
DEX0306_94 DEX0157_54 DEX0306_236
DEX0306_95 flex DEX0157_54
5 DEX0306_96 DEX0157_55 DEX0306_237
DEX0306_97 DEX0157_56 DEX0306_238
DEX0306_98 flex DEX0157_56 DEX0306_239
DEX0306_99 DEX0157_57 DEX0306_240
DEX0306_100 DEX0157_58 DEX0306_241
10 DEX0306_101 flex DEX0157_58
DEX0306_102 DEX0157_60 DEX0306_242
DEX0306_103 flex DEX0157_60 DEX0306_243
DEX0306_104 DEX0157_61 DEX0306_244
DEX0306_105 flex DEX0157_61 DEX0306_245
15 DEX0306_106 DEX0157_62 DEX0306_246
DEX0306_107 flex DEX0157_62 DEX0306_247
DEX0306_108 DEX0157_63 DEX0306_248
DEX0306_109 flex DEX0157_63
DEX0306_110 DEX0157_64 DEX0306_249
20 DEX0306_111 flex DEX0157_64 DEX0306_250
DEX0306_112 DEX0157_65 DEX0306_251
DEX0306_113 DEX0157_66 DEX0306_252
DEX0306_114 DEX0157_67 DEX0306_253
DEX0306_115 DEX0157_68 DEX0306_254
25 DEX0306_116 flex DEX0157_68 DEX0306_255
DEX0306_117 DEX0157_69 DEX0306_256
DEX0306_118 flex DEX0157_69 DEX0306_257
DEX0306_119 DEX0157_70 DEX0306_258
DEX0306_120 flex DEX0157_70
30 DEX0306_121 DEX0157_71 DEX0306_259
DEX0306_122 flex DEX0157_71
DEX0306_123 DEX0157_72 DEX0306_260
DEX0306_124 flex DEX0157_72 DEX0306_261
DEX0306_125 DEX0157_73 DEX0306_262
35 DEX0306_126 flex DEX0157_73 DEX0306_263
DEX0306_127 DEX0157_74 DEX0306_264
DEX0306_128 flex DEX0157_74
DEX0306_129 DEX0157_75 DEX0306_265
DEX0306_130 DEX0157_76 DEX0306_266
40 DEX0306_131 flex DEX0157_76 DEX0306_267
DEX0306_132 DEX0157_77 DEX0306_268
DEX0306_133 flex DEX0157_77
DEX0306_134 DEX0157_78 DEX0306_269
DEX0306_135 flex DEX0157_78 DEX0306_270
45 DEX0306_136 DEX0157_79 DEX0306_271
DEX0306_137 flex DEX0157_79 DEX0306_272
DEX0306_138 DEX0157_80 DEX0306_273
DEX0306_139 DEX0157_81 DEX0306_274
DEX0306_140 flex DEX0157_81 DEX0306_275
50 DEX0306_141 DEX0157_82 DEX0306_276
DEX0306_142 flex DEX0157_82
DEX0306_143 DEX0157_83 DEX0306_277
DEX0306_144 flex DEX0157_83
DEX0306_145 DEX0157_85 DEX0306_278
55 DEX0306_146 flex DEX0157_85
DEX0306_147 DEX0157_86 DEX0306_279
DEX0306_148 flex DEX0157_86 DEX0306_280
DEX0306_149 DEX0157_87 DEX0306_281
DEX0306_150 flex DEX0157_87
60 DEX0306_151 DEX0157_88 DEX0306_282
DEX0306_152 flex DEX0157_88
DEX0306_153 DEX0157_89 DEX0306_283

-112-

DEX0306_154 flex DEX0157_89
 DEX0306_155 DEX0157_90 DEX0306_284
 DEX0306_156 flex DEX0157_90 DEX0306_285
 DEX0306_157 DEX0157_93 DEX0306_286
 5 DEX0306_158 DEX0157_94 DEX0306_287
 DEX0306_159 flex DEX0157_94
 DEX0306_160 DEX0157_95 DEX0306_288
 DEX0306_161 flex DEX0157_95
 DEX0306_162 DEX0157_96 DEX0306_289
 10 DEX0306_163 DEX0157_97 DEX0306_290
 DEX0306_164 flex DEX0157_97
 DEX0306_165 DEX0157_98 DEX0306_291
 DEX0306_166 DEX0157_99 DEX0306_292
 DEX0306_167 DEX0157_100 DEX0306_293
 15 DEX0306_168 flex DEX0157_100
 DEX0306_169 DEX0157_101 DEX0306_294
 DEX0306_170 DEX0157_102 DEX0306_295
 DEX0306_171 flex DEX0157_102

20 Example 2: Relative Quantitation of Gene Expression

Real-Time quantitative PCR with fluorescent Taqman probes is a quantitation
 detection system utilizing the 5'-3' nuclease activity of Taq DNA polymerase. The
 method uses an internal fluorescent oligonucleotide probe (Taqman) labeled with a 5'
 reporter dye and a downstream, 3' quencher dye. During PCR, the 5'-3' nuclease activity
 25 of Taq DNA polymerase releases the reporter, whose fluorescence can then be detected
 by the laser detector of the Model 7700 Sequence Detection System (PE Applied
 Biosystems, Foster City, CA, USA). Amplification of an endogenous control is used to
 standardize the amount of sample RNA added to the reaction and normalize for Reverse
 Transcriptase (RT) efficiency. Either cyclophilin, glyceraldehyde-3-phosphate
 30 dehydrogenase (GAPDH), ATPase, or 18S ribosomal RNA (rRNA) is used as this
 endogenous control. To calculate relative quantitation between all the samples studied,
 the target RNA levels for one sample were used as the basis for comparative results
 (calibrator). Quantitation relative to the "calibrator" can be obtained using the standard
 curve method or the comparative method (User Bulletin #2: ABI PRISM 7700 Sequence
 35 Detection System).

The tissue distribution and the level of the target gene are evaluated for every
 sample in normal and cancer tissues. Total RNA is extracted from normal tissues, cancer
 tissues, and from cancers and the corresponding matched adjacent tissues. Subsequently,
 first strand cDNA is prepared with reverse transcriptase and the polymerase chain
 40 reaction is done using primers and Taqman probes specific to each target gene. The
 results are analyzed using the ABI PRISM 7700 Sequence Detector. The absolute
 numbers are relative levels of expression of the target gene in a particular tissue
 compared to the calibrator tissue.

One of ordinary skill can design appropriate primers. The relative levels of expression of the BSNA versus normal tissues and other cancer tissues can then be determined. All the values are compared to a normal tissue (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

The relative levels of expression of the BSNA in pairs of matching samples and 1 cancer and 1 normal/normal adjacent of tissue may also be determined. All the values are compared to a normal tissue (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual.

In the analysis of matching samples, BSNA show a high degree of tissue specificity for the tissue of interest. These results confirm the tissue specificity results obtained with normal pooled samples.

Further, the level of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual are compared. This comparison provides an indication of specificity for the cancer stage (*e.g.* higher levels of mRNA expression in the cancer sample compared to the normal adjacent).

Altogether, the high level of tissue specificity, plus the mRNA overexpression in matching samples tested are indicative of SEQ ID NO: 1 through 171 being diagnostic markers for cancer.

Example 2B: Custom Microarray Experiment

Custom oligonucleotide microarrays were provided by Agilent Technologies, Inc. (Palo Alto, CA). The microarrays were fabricated by Agilent using their technology for the *in-situ* synthesis of 60mer oligonucleotides (Hughes, et al. 2001, Nature Biotechnology 19:342-347). The 60mer microarray probes were designed by Agilent, from gene sequences provided by diaDexus, using Agilent proprietary algorithms. Whenever possible two different 60mers were designed for each gene of interest.

All microarray experiments were two-color experiments and were performed using Agilent-recommended protocols and reagents. Briefly, each microarray was hybridized with cRNAs synthesized from polyA+ RNA, isolated from cancer and normal tissues, labeled with fluorescent dyes Cyanine3 and Cyanine5 (NEN Life Science Products, Inc., Boston, MA) using a linear amplification method (Agilent). In each experiment, the experimental sample was polyA+ RNA isolated from cancer tissue from

-114-

a single individual and the reference sample was a pool of polyA+ RNA isolated from normal tissues of the same organ as the cancerous tissue (*i.e.* normal breast tissue in experiments with breast cancer samples). Hybridizations were carried out at 60°C, overnight using Agilent *in-situ* hybridization buffer. Following washing, arrays were
5 scanned with a GenePix 4000B Microarray Scanner (Axon Instruments, Inc., Union City, CA). The resulting images were analyzed with GenePix Pro 3.0 Microarray Acquisition and Analysis Software (Axon). A total of 36 experiments comparing the expression patterns of breast cancer derived polyA+ RNA (9 stage 1 cancers, 23 stage 2 cancers, 4 stage 3 cancers) to polyA+ RNA isolated from a pool of 10 normal breast tissues were
10 analyzed.

Data normalization and expression profiling were done with Expressionist software from GeneData Inc. (Daly City, CA/Basel, Switzerland). Gene expression analysis was performed using only experiments that meet certain quality criteria. The quality criteria that experiments must meet are a combination of evaluations performed
15 by the Expressionist software and evaluations performed manually using raw and normalized data. To evaluate raw data quality, detection limits (the mean signal for a replicated negative control ± 2 Standard Deviations (SD)) for each channel were calculated. The detection limit is a measure of non-specific hybridization. Arrays with poor detection limits were not analyzed and the experiments were repeated. To evaluate
20 normalized data quality, positive control elements included in the array were utilized. These array features should have a mean ratio of 1 (no differential expression). If these features have a mean ratio of greater than 1.5-fold up or down, the experiments were not analyzed further and were repeated. In addition to traditional scatter plots demonstrating the distribution of signal in each experiment, the Expressionist software also has
25 minimum thresholding criteria that employs user defined parameters to identify quality data. Only those features that meet the threshold criteria were included in the filtering and analyses carried out by Expressionist. The thresholding settings employed require a minimum area percentage of 60% [(% pixels > background $\pm 2SD$)-(% pixels saturated)], and a minimum signal to noise ratio of 2.0 in both channels. By these criteria, very low
30 expressors and saturated features were not included in analysis.

Relative expression data was collected from Expressionist based on meeting the quality parameters described above. Sensitivity data was calculated using an analysis tool. Up- and down- regulated genes were identified using criteria for percentage of valid values obtained, and the percentage of experiments in which the gene is up- or

-115-

down-regulated. These criteria were set independently for each data set, depending on the size and the nature of the data set. Results for several BSNA's are shown in the following table. The first three columns of the table contain information about the sequence itself (Oligo ID, Parent ID, and SEQ ID NO), the next 3 columns show the results obtained. '%valid' indicates the percentage of 36 unique experiments total in which a valid expression value was obtained, '%up' indicates the percentage of 20 experiments in which up-regulation of at least 2.5-fold was observed, and '%down' indicates the percentage of the 36 experiments in which down-regulation of at least 2.5-fold was observed. The last column in Table 1 describes the location of the microarray probe (oligo) relative to the sequence.

OligoID	Parent ID	Patent # SEQ ID NO	Sensitivity of up and down regulation			Oligo Seq location in original seq.	Oligo Seq location in FLEX seq
			% valid	% up	% down		
16052	8056	DEX0157_74, DEX0131_52 SEQ ID NO: 127/128	100	11.1	33.3	75-134	1928-1987
24688	5998	DEX0167_22, DEX0157_95, DEX0133_22, DEX0131_78 SEQ ID NO: 160/161	94.4	2.8	58.3	437-496	1093-1152
24689	5998	DEX0157_95, DEX0131_78 SEQ ID NO: 160/161	97.2	2.8	61.1	397-456	
27873	8713	DEX0157_74, DEX0131_52 SEQ ID NO: 127/128	100	13.9	30.6	101-160	1954-2013
33090	5973	DEX0157_73, DEX0131_56 SEQ ID NO: 125/126	97.2	2.8	44.4	408-466	2142-2200
33091	5973	DEX0157_73, DEX0131_56 SEQ ID NO: 125/126	100	2.8	41.7	368-427	1221-1280

Example 3: Protein Expression

The BSNA is amplified by polymerase chain reaction (PCR) and the amplified DNA fragment encoding the BSNA is subcloned in pET-21d for expression in *E. coli*. In addition to the BSNA coding sequence, codons for two amino acids, Met-Ala, flanking the NH₂-terminus of the coding sequence of BSNA, and six histidines, flanking the

-116-

COOH-terminus of the coding sequence of BSNA, are incorporated to serve as initiating Met/restriction site and purification tag, respectively.

An over-expressed protein band of the appropriate molecular weight may be observed on a Coomassie blue stained polyacrylamide gel. This protein band is
5 confirmed by Western blot analysis using monoclonal antibody against 6X Histidine tag.

Large-scale purification of BSP was achieved using cell paste generated from 6-liter bacterial cultures, and purified using immobilized metal affinity chromatography (IMAC). Soluble fractions that had been separated from total cell lysate were incubated with a nickle chelating resin. The column was packed and washed with five column
10 volumes of wash buffer. BSP was eluted stepwise with various concentration imidazole buffers.

Example 4: Protein Fusions

Briefly, the human Fc portion of the IgG molecule can be PCR amplified, using primers that span the 5' and 3' ends of the sequence described below. These primers also
15 should have convenient restriction enzyme sites that will facilitate cloning into an expression vector, preferably a mammalian expression vector. For example, if pC4 (Accession No. 209646) is used, the human Fc portion can be ligated into the BamHI cloning site. Note that the 3' BamHI site should be destroyed. Next, the vector containing the human Fc portion is re-restricted with BamHI, linearizing the vector, and a
20 polynucleotide of the present invention, isolated by the PCR protocol described in Example 2, is ligated into this BamHI site. Note that the polynucleotide is cloned without a stop codon, otherwise a fusion protein will not be produced. If the naturally occurring signal sequence is used to produce the secreted protein, pC4 does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the
25 vector can be modified to include a heterologous signal sequence. *See, e. g.*, WO 96/34891.

Example 5: Production of an Antibody from a Polypeptide

In general, such procedures involve immunizing an animal (preferably a mouse) with polypeptide or, more preferably, with a secreted polypeptide-expressing cell. Such
30 cells may be cultured in any suitable tissue culture medium; however, it is preferable to culture cells in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at about 56°C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100, µg/ml of streptomycin. The

-117-

splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line (SP20), available from the ATCC. After fusion, the resulting hybridoma cells are selectively maintained in HAT
 5 medium, and then cloned by limiting dilution as described by Wands *et al.*,
Gastroenterology 80: 225-232 (1981).

The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the polypeptide. Alternatively, additional antibodies capable of binding to the polypeptide can be
 10 produced in a two-step procedure using anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the
 15 hybridoma cells are screened to identify clones which produce an antibody whose ability to bind to the protein-specific antibody can be blocked by the polypeptide. Such antibodies comprise anti-idiotypic antibodies to the protein specific antibody and can be used to immunize an animal to induce formation of further protein-specific antibodies. Using the Jameson-Wolf methods the following epitopes were predicted. (Jameson and
 20 Wolf, CABIOS, 4(1), 181-186, 1988, the contents of which are incorporated by reference).

The predicted antigenicity for the amino acid sequences is as follows:

DEX ID	ANTIGENICITY Position, AI Ave, Length	TRANSMEMBRANE Predicted Helix, Topology	PTM PTM	SIGNAL PEPTIDE Position, Max Score, Mean Score
DEX0306_ 172			Myristyl 28-33; 53-58; 60- 65; Pkc_Phospho_ Site 67-69;	26, .882, .574
DEX0306_ 173			Myristyl 13-18; Pkc_Phospho_Site 19-21;	
DEX0306_ 174		1, i20-42o		
DEX0306_ 175	11-21, 1.07, 11		Pkc_Phospho_Site 4-6; 12-14;	
DEX0306_ 176	52-69, 1.16, 18 9-18, 1.16, 10		Asn_Glycosylation 82-85; Ck2_Phospho_Site 7-10; Myristyl 79-84;	

-118-

DEX ID	ANTIGENICITY Position, AI Ave, Length	TRANSMEMBRANE Predicted Helix, Topology	PTM PTM	SIGNAL PEPTIDE Position, Max Score, Mean Score
			Pkc_Phospho_Site 4-6;	
DEX0306_177			Asn_Glycosylation 7-10;55-58; Ck2_Phospho_Site 22-25;57-60; Pkc_Phospho_Site 57-59; Tyr_Phospho_Site 46-52;	
DEX0306_178	10-47, 1.07, 38 80-141, 1.03, 62		Myristyl 33- 38;129-134; Pkc_Phospho_Site 116-118;147-149;	
DEX0306_179			Myristyl 3-8;	
DEX0306_180	59-74, 1.04, 16		Ck2_Phospho_Site 4-7;49-52; Myristyl 45- 50;50-55;80- 85;86-91;95-100; Pkc_Phospho_Site 60-62;65-67;69- 71;	
DEX0306_182			Myristyl 22-27;	
DEX0306_184	12-36, 1.22, 25		Asn_Glycosylation 32-35; Camp_Phospho_Site 26-29; Ck2_Phospho_Site 9-12; Pkc_Phospho_Site 25-27;	
DEX0306_185	6-39, 1.13, 34		Asn_Glycosylation 64-67; Ck2_Phospho_Site 37-40;65-68; Glycosaminoglycan 48-51; Myristyl 14-19;49-54;51- 56; Pkc_Phospho_Site 18-20;42-44;	
DEX0306_187		1,o25-47i	Ck2_Phospho_Site 70-73; Myristyl 7-12; Pkc_Phospho_Site 42-44;	
DEX0306_188		3,i5-22o32- 54i61-83o	Myristyl 27- 32;141-146;144- 149; Pkc_Phospho_Site 17-19;55-57;90- 92;111-113;	17, .989, .91
DEX0306			Ck2_Phospho_Site	

DEX ID	ANTIGENICITY Position, AI Ave, Length	TRANSMEMBRANE Predicted Helix, Topology	PTM PTM	SIGNAL PEPTIDE Position, Max Score, Mean Score
190			73-76; Myristyl 12-17;17-22;66- 71; Pkc_Phospho_Site 91-93;	
DEX0306_ 192			Pkc_Phospho_Site 6-8;	
DEX0306_ 193			Myristyl 4-9;	
DEX0306_ 194	415-439, 1.14,25 242-251, 1.13,10 459-528, 1.11,70 159-197, 1.09,39 777-810, 1.09,34 632-669, 1.07,38 1034-1044, 1.04,11 1077-1103, 1.03,27		Asn_Glycosylation 12-15;19-22;23- 26;151-154;513- 516;873-876;886- 889; Camp_Phospho_ Site 107-110; Ck2_Phospho_Site 72-75;260- 263;283-286;319- 322;463-466;807- 810;975-978; Glycosaminoglycan 125-128;905- 908;913-916; Myristyl 13- 18;28-33;30- 35;52-57;53- 58;58-63;61- 66;62-67;126- 131;179-184;372- 377;529-534;699- 704;716-721;717- 722;721-726;837- 842;845-850;889- 894;906-911;910- 915; Pkc_Phospho_Site 129-131;160- 162;188-190;189- 191;356-358;613- 615;822-824;825- 827; Prokar_ Lipoprotein 44- 54;	
DEX0306_ 195			Pkc_Phospho_Site 6-8;	
DEX0306_ 196			Pkc_Phospho_Site 24-26;33-35;	
DEX0306_ 197			Pkc_Phospho_Site 7-9;	
DEX0306_ 198	39-55,1.09,17 25-34,1.05,10		Ck2_Phospho_Site 92-95; Pkc_Phospho_Site 107-109;	
DEX0306_ 199	97-113, 1.09,17		Ck2_Phospho_Site 150-153;193-	

-120-

DEX ID	ANTIGENICITY Position, AI Ave, Length	TRANSMEMBRANE Predicted Helix, Topology	PTM PTM	SIGNAL PEPTIDE Position, Max Score, Mean Score
	83-92, 1.05, 10		196; 200-203; Myristyl 11- 16; 178-183; Pkc_Phospho_Site 165-167; Tyr_Phospho_Site 53-61;	
DEX0306_ 200		1, i12-34o	Asn_Glycosylation 20-23; Myristyl 18-23;	
DEX0306_ 201			Myristyl 16-21; Pkc_Phospho_Site 24-26;	24, .944, .7 79
DEX0306_ 202	25-37, 1.17, 13		Ck2_Phospho_Site 12-15; Myristyl 27-32; 31-36; 53- 58;	
DEX0306_ 203			Asn_Glycosylation 28-31; Myristyl 8-13; 62-67; 63- 68; 64-69;	
DEX0306_ 204			Pkc_Phospho_Site 2-4;	
DEX0306_ 205			Ck2_Phospho_Site 60-63; 77-80; Myristyl 14-19; Pkc_Phospho_Site 57-59;	
DEX0306_ 206		1, o5-24i	Myristyl 4-9;	
DEX0306_ 207			Ck2_Phospho_Site 64-67; 75-78; Myristyl 71- 76; 81-86; 85-90;	
DEX0306_ 208			Asn_Glycosylation 53-56; 62-65; Myristyl 72-77; Pkc_Phospho_Site 63-65; 64-66;	
DEX0306_ 209			Asn_Glycosylation 47-50; Pkc_Phospho_Site 28-30; 38-40; Tyr_Phospho_Site 29-36; 30-36;	
DEX0306_ 211			Asn_Glycosylation 33-36; Ck2_Phospho_Site 17-20; Pkc_Phospho_Site 26-28;	
DEX0306_ 212	30-39, 1.06, 10		Ck2_Phospho_Site 76-79; Myristyl 19-24; 31-36; 92- 97; Pkc_Phospho_Site	17, .97, .82 9

-121-

DEX ID	ANTIGENICITY Position, AI Ave, Length	TRANSMEMBRANE Predicted Helix, Topology	PTM PTM	SIGNAL PEPTIDE Position, Max Score, Mean Score
DEX0306_213			12-14;76-78; Pkc_Phospho_Site 29-31;	
DEX0306_214			Myristyl 43- 48;48-53;	
DEX0306_215	104-118, 1.16,15		Myristyl 90- 95;101-106;104- 109;	21,.973,.8 2
DEX0306_216			Ck2_Phospho_Site 5-8;	
DEX0306_217		1,i11-33o	Myristyl 42- 47;54-59;67-72; Pkc_Phospho_Site 4-6;37-39;	33,.982,.8 23
DEX0306_218			Asn_Glycosylation 12-15; Ck2_Phospho_Site 8-11; Myristyl 3- 8; Pkc_Phospho_Site 23-25;	
DEX0306_219			Asn_Glycosylation 21-24; Ck2_Phospho_Site 43-46; Pkc_Phospho_Site 23-25;	
DEX0306_220	14-32,1.13,19		Amidation 19-22; Pkc_Phospho_Site 23-25;	
DEX0306_221			Pkc_Phospho_Site 18-20;	
DEX0306_223			Pkc_Phospho_Site 2-4;	
DEX0306_224			Ck2_Phospho_Site 31-34;38-41;57- 60;79-82;85-88; Pkc_Phospho_Site 7-9;	
DEX0306_225		1,i17-26o	Asn_Glycosylation 34-37; Ck2_Phospho_Site 36-39;	
DEX0306_226			Pkc_Phospho_Site 34-36;	15,.918,.7 44
DEX0306_227	52-72,1.19,21	1,i173-95o	Amidation 66-69; Ck2_Phospho_Site 6-9; Myristyl 74- 79;78-83;	
DEX0306_228		1,i20-42o		
DEX0306_230		1,o22-44i	Prokar_ Lipoprotein 23- 33;	
DEX0306_231			Camp_Phospho_Site 3-6; Myristyl 31-	

DEX ID	ANTIGENICITY Position, AI Ave, Length	TRANSMEMBRANE Predicted Helix, Topology	PTM PTM	SIGNAL PEPTIDE Position, Max Score, Mean Score
DEX0306_232		1,015-32i	36;90-95; Myristyl 47-52; Pkc_Phospho_Site 2-4;	
DEX0306_233			Asn_Glycosylation 4-7;	
DEX0306_234	24-39,1.2,16		Myristyl 8-13; Pkc_Phospho_Site 65-67;	
DEX0306_235	560-572, 1.27,13 509-519, 1.23,11 1126-1153, 1.19,28 861-873, 1.18,13 794-804, 1.16,11 964-976, 1.16,13 880-901, 1.16,22 812-828, 1.11,17 588-612, 1.09,25 41-77, 1.07,37 461-489, 1.07,29 735-751, 1.07,17 978-1011, 1.06,34 535-558, 1.04,24 1081-1.04,17 620-644, 1.03,25 654-671, 1.01,18 354-382,1,29		Amidation 281- 284;403-406;721- 724; Asn_Glycosylation 633-636;655-658; Atp_Gtp_A 507- 514; Camp_Phospho_Site 54-57;479-482; Ck2_Phospho_Site 132-135;144- 147;181-184;209- 212;217-220;244- 247;310-313;332- 335;345-348;546- 549;558-561;560- 563;593-596;617- 620;622-625;635- 638;651-654;656- 659;697-700;739- 742;740-743;745- 748;969-972; Glycosaminoglycan 482-485;719-722; Myristyl 110- 115;130-135;142- 147;159-164;230- 235;254-259;277- 282;341-346;400- 405;510-515;572- 577;582-587;645- 650;721-726;823- 828;842-847;843- 848;846-851;872- 877;922-927;940- 945;954-959; Pkc_Phospho_Site 72-74;83-85;148- 150;155-157;156- 158;209-211;627- 629;635-637;656- 658;660-662;661- 663;736-738;739- 741;745-747;766- 768;802-804;813- 815;913-915;965- 967;973-975;	

-123-

DEX ID	ANTIGENICITY Position, AI Ave, Length	TRANSMEMBRANE Predicted Helix, Topology	PTM PTM	SIGNAL PEPTIDE Position, Max Score, Mean Score
			Tyr_Phospho_Site 55-62;426-433; Zinc_Finger_C2h2 36-56;176- 197;250-270;278- 298;337-357;517- 537;	
DEX0306_ 236	11-29, 1, 19	1,032-54i		
DEX0306_ 237			Glycosaminoglycan 80-83; Myristyl 14-19;54-59;58- 63; Pkc_Phospho_Site 68-70;80-82;	
DEX0306_ 238		1,062-84i	Asn_Glycosylation 30-33; Pkc_Phospho_Site 31-33;	
DEX0306_ 239	42-63, 1.12, 22		Asn_Glycosylation 145-148; Ck2_Phospho_Site 4-7;63-66;151- 154; Euk_Co2_Anhydrase 126-142; Myristyl 25-30;33-38;125- 130; Pkc_Phospho_Site 280-282;	
DEX0306_ 240	20-34, 1.08, 15		Asn_Glycosylation 53-56; Camp_Phospho_Site 41-44; Pkc_Phospho_Site 39-41;	
DEX0306_ 242			Myristyl 49-54; Pkc_Phospho_Site 33-35;	
DEX0306_ 243			Ck2_Phospho_Site 23-26;24-27; Pkc_Phospho_Site 9-11;23-25;	
DEX0306_ 244			Asn_Glycosylation 4-7;	
DEX0306_ 245	45-55, 1.15, 11		Camp_Phospho_Site 51-54; Ck2_Phospho_Site 60-63; Pkc_Phospho_Site 22-24;	
DEX0306_ 246			Pkc_Phospho_Site 7-9;35-37;	
DEX0306_ 247			Myristyl 86-91; Pkc_Phospho_Site 17-19;	22,.929,.6 52

-124-

DEX ID	ANTIGENICITY Position, AI Ave, Length	TRANSMEMBRANE Predicted Helix, Topology	PTM PTM	SIGNAL PEPTIDE Position, Max Score, Mean Score
DEX0306_248				18, .993, .9 14
DEX0306_249			Asn_Glycosylation 2-5; Ck2_Phospho_Site 54-57; Pkc_Phospho_Site 54-56;	28, .911, .7 4
DEX0306_250	142-180, 1.03,39 9-21,1,13		Asn_Glycosylation 13-16;132-135; Ck2_Phospho_Site 97-100; Pkc_Phospho_Site 17-19;55-57;113- 115;134-136;153- 155;	
DEX0306_251	113-123, 1.14,11 37-60,1.09,24		Camp_Phospho_Site 50-53; Ck2_Phospho_Site 88-91; Pkc_Phospho_Site 39-41;49-51;88- 90; Prokar_Lipoprotei n 59-69; Tyr_Phospho_Site 87-95;	
DEX0306_252			Pkc_Phospho_Site 10-12;	
DEX0306_253		1,i12-43o	Myristyl 30-35; Prokar_Lipoprotei n 12-22;	30, .996, .8 62
DEX0306_254			Ck2_Phospho_Site 16-19; Myristyl 31-36;36-41; Pkc_Phospho_Site 32-34; Rgd 25-27;	
DEX0306_255			Asn_Glycosylation 386-389;516- 519;536-539;626- 629;638-641;883- 886; Camp_Phospho_Site 61-64; Ck2_Phospho_Site 147-150;201- 204;205-208;252- 255;394-397;435- 438;462-465;491- 494;511-514;524- 527;552-555;632- 635;646-649;756- 759;839-842;867- 870;887-890; Myristyl 25- 30;263-268;751-	

DEX ID	ANTIGENICITY Position, AI Ave, Length	TRANSMEMBRANE Predicted Helix, Topology	PTM PTM	SIGNAL PEPTIDE Position, Max Score, Mean Score
			756;879-884; Pkc_Phospho_Site 29-31;107- 109;147-149;201- 203;506-508; Tyr_Phospho_Site 467-473;	
DEX0306_256	65-75, 1.02, 11 25-50, 1.02, 26		Asn_Glycosylation 56-59; Myristyl 14-19; Prokar_ Lipoprotein 8-18;	
DEX0306_257	179-203, 1.18, 25 527-569, 1.15, 43 422-464, 1.11, 43 20-39, 1.06, 20 335-367, 1.06, 33 43-117, 1.01, 75		Amidation 267- 270; Asn_Glycosylation 176-179; Camp_Phospho_Site 71-74;324-327; Ck2_Phospho_Site 42-45;54-57;75- 78;99-102;109- 112;161-164;197- 200;206-209;223- 226;228-231;273- 276;283-286;336- 339;447-450;482- 485;497-500;567- 570; Glycosaminoglycan 246-249; Myristyl 24-29;38-43;86- 91;124-129;249- 254;262-267;278- 283;290-295;332- 337;410-415;430- 435; Pkc_Phospho_Site 12-14;18-20;28- 30;35-37;54- 56;69-71;296- 298;336-338;411- 413;434-436; Tyr_Phospho_Site 23-29;137- 144;310-318;	
DEX0306_258			Ck2_Phospho_Site 34-37;	
DEX0306_259			Asn_Glycosylation 31-34;	
DEX0306_260			Camp_Phospho_Site 6-9; Myristyl 54- 59;	
DEX0306_261	96-105, 1.19, 10		Ck2_Phospho_Site 71-74;101-104; Glycosaminoglycan 55-58; Myristyl 52-57;54-59;58-	

-126-

DEX ID	ANTIGENICITY Position, AI Ave, Length	TRANSMEMBRANE Predicted Helix, Topology	PTM PTM	SIGNAL PEPTIDE Position, Max Score, Mean Score
			63;67-72; Pkc_Phospho_Site 17-19;137- 139;146-148;197- 199;215-217; Prokar_Lipoprotei n 164-174;	
DEX0306_ 262	30-41,1.02,12		Asn_Glycosylation 86-89; Ck2_Phospho_Site 21-24; Myristyl 96-101; Pkc_Phospho_Site 18-20;	
DEX0306_ 263	239-249, 1.13,11		Amidation 72-75; Asn_Glycosylation 119-122;120-123; Camp_Phospho_Site 107-110;216-219; Ck2_Phospho_Site 28-31;43-46;63- 66;160-163;169- 172;187-190; Myristyl 69- 74;158-163; Pkc_Phospho_Site 17-19;24-26;35- 37;52-54;59- 61;106-108;122- 124;184-186; Prokar_ Lipoprotein 248- 258;	
DEX0306_ 264			Myristyl 35-40; Pkc_Phospho_Site 21-23;22-24;	
DEX0306_ 265		1,i7-29o	Camp_Phospho_Site 47-50; Ck2_Phospho_Site 54-57; Myristyl 37-42; Pkc_Phospho_Site 72-74;	
DEX0306_ 266			Asn_Glycosylation 7-10;17-20; Pkc_Phospho_Site 2-4;	
DEX0306_ 267			Amidation 43-46; Ck2_Phospho_Site 79-82; Pkc_Phospho_Site 11-13;89-91;	
DEX0306_ 268			Pkc_Phospho_Site 8-10;45-47; Prokar_Lipoprotei n 32-42;	

-127-

DEX ID	ANTIGENICITY Position, AI Ave, Length	TRANSMEMBRANE Predicted Helix, Topology	PTM PTM	SIGNAL PEPTIDE Position, Max Score, Mean Score
DEX0306_ 269			Camp_Phospho_Site 66-69; Ck2_Phospho_Site 12-15;34-37;56- 59; Myristyl 30- 35; Pkc_Phospho_Site 34-36;56-58;	
DEX0306_ 270	49-134,1,86		Asn_Glycosylation 46-49; Ck2_Phospho_Site 65-68;84-87;93- 96;109-112; Myristyl 4-9;59- 64; Pkc_Phospho_Site 60-62;89-91;104- 106;115-117;116- 118; Tyr_Phospho_Site 92-99;117- 124;118-124;	
DEX0306_ 272	235-299,1.1, 65 369-406, 1.07,38 99-109, 1.01,11		Asn_Glycosylation 37-40;69-72;284- 287; Ck2_Phospho_Site 85-88;141- 144;149-152;192- 195;204-207; Glycosaminoglycan 433-436; Myristyl 43-48;44-49;96- 101;118-123;402- 407;406-411;432- 437;438-443; Pkc_Phospho_Site 48-50;433-435; Rgd 278-280; Tyr_Phospho_Site 50-56;	
DEX0306_ 273			Pkc_Phospho_Site 6-8;15-17;	
DEX0306_ 274			Asn_Glycosylation 44-47;	
DEX0306_ 275			Asn_Glycosylation 78-81; Ck2_Phospho_Site 17-20; Myristyl 13-18;	
DEX0306_ 276			Ck2_Phospho_Site 58-61; Glycosaminoglycan 93-96; Myristyl 28-33;48-53;50- 55;67-72;71-76; Pkc_Phospho_Site	

-128-

DEX ID	ANTIGENICITY Position, AI Ave, Length	TRANSMEMBRANE Predicted Helix, Topology	PTM PTM	SIGNAL PEPTIDE Position, Max Score, Mean Score
			5-7;18-20;44-46;57-59; Rgd 59-61;	
DEX0306_277		1,037-59i	Ck2_Phospho_Site 22-25; Myristyl 71-76; Pkc_Phospho_Site 12-14;	
DEX0306_279			Myristyl 15-20;	
DEX0306_280			Ck2_Phospho_Site 76-79; Pkc_Phospho_Site 16-18;	27,.985,.682
DEX0306_281	17-29,1.07,13		Myristyl 5-10;9-14; Pkc_Phospho_Site 24-26;	
DEX0306_282		1,015-32i		
DEX0306_283			Asn_Glycosylation 35-38; Ck2_Phospho_Site 37-40; Myristyl 3-8; Pkc_Phospho_Site 57-59;	
DEX0306_284	28-37,1.09,10		Ck2_Phospho_Site 46-49; Pkc_Phospho_Site 32-34;	21,.958,.821
DEX0306_285	226-245, 1.37,20 489-501, 1.22,13 1271-1284, 1.21,14 1192-1203, 1.11,12 745-755, 1.09,11 929-940, 1.08,12 1039-1051, 1.08,13 1133-1150, 1.05,18 547-576, 1.05,30 89-98,1.04,10 22-53,1.03,32 1073-1086, 1.03,14 1243-1253, 1.03,11 1418-1461,		Amidation 473-476; Asn_Glycosylation 512-515;726-729; Camp_Phospho_Site 475-478;571-574;646-649; Ck2_Phospho_Site 29-32;143-146;176-179;228-231;230-233;232-235;263-266;294-297;388-391;447-450;493-496;506-509;517-520;581-584;664-667;890-893;929-932; Gram_Pos_Anchorin g 670-675; Myristyl 49-54;56-61;125-130;152-157;185-190;214-219;677-682;708-713;840-845;921-926;	

DEX ID	ANTIGENICITY Position, AI Ave, Length	TRANSMEMBRANE Predicted Helix, Topology	PTM PTM	SIGNAL PEPTIDE Position, Max Score, Mean Score
	1.01, 44		Pkc_Phospho_Site 21-23;29-31;143- 145;388-390;415- 417;443-445;530- 532;539-541;552- 554;565-567;581- 583;748-750;802- 804;925-927;931- 933;987-989;996- 998; Tyr_Phospho_Site 867-874;Amidation 473-476; Asn_Glycosylation 512-515;726-729; Camp_Phospho_Site 475-478;571- 574;646-649; Ck2_Phospho_Site 29-32;143- 146;176-179;228- 231;230-233;232- 235;263-266;294- 297;388-391;447- 450;493-496;506- 509;517-520;581- 584;664-667;890- 893;929-932; Gram_Pos_ Anchoring 670- 675; Myristyl 49- 54;56-61;125- 130;152-157;185- 190;214-219;677- 682;708-713;840- 845;921-926; Pkc_Phospho_Site 21-23;29-31;143- 145;388-390;415- 417;443-445;530- 532;539-541;552- 554;565-567;581- 583;748-750;802- 804;925-927;931- 933;987-989;996- 998; Tyr_Phospho_Site 867-874;	
DEX0306_286		2, i13-30o35-54i	Asn_Glycosylation 15-18; Ck2_Phospho_Site 41-44; Myristyl 2-7; Pkc_Phospho_Site 6-8;	
DEX0306			Asn_Glycosylation	

-130-

DEX ID	ANTIGENICITY	TRANSMEMBRANE	PTM	SIGNAL PEPTIDE
	Position, AI Ave, Length	Predicted Helix, Topology	PTM	Position, Max Score, Mean Score
287			43-46;51-54; Ck2_Phospho_Site 34-37; Pkc_Phospho_Site 70-72;	
DEX0306_ 288			Asn_Glycosylation 42-45; Camp_Phospho_Site 12-15; Myristyl 4-9;	
DEX0306_ 290	20-31, 1. 14, 12		Pkc_Phospho_Site 6-8; 21-23;	
DEX0306_ 291			Glycosaminoglycan 31-34; Myristyl 30-35;	
DEX0306_ 292			Camp_Phospho_Site 8-11; Ck2_Phospho_Site 11-14;	
DEX0306_ 293			Ck2_Phospho_Site 36-39; Myristyl 2-7; 94-99;	
DEX0306_ 294	31-52, 1. 01, 22		Pkc_Phospho_Site 47-49;	
DEX0306_ 295			Myristyl 56-61;	

Example 6: Method of Determining Alterations in a Gene Corresponding to a Polynucleotide

- 5 RNA is isolated from individual patients or from a family of individuals that have a phenotype of interest. cDNA is then generated from these RNA samples using protocols known in the art. *See*, Sambrook (2001), *supra*. The cDNA is then used as a template for PCR, employing primers surrounding regions of interest in SEQ ID NO: 1 through 171. Suggested PCR conditions consist of 35 cycles at 95°C for 30 seconds;
- 10 60-120 seconds at 52-58°C; and 60-120 seconds at 70°C, using buffer solutions described in Sidransky *et al.*, *Science* 252(5006): 706-9 (1991). *See also* Sidransky *et al.*, *Science* 278(5340): 1054-9 (1997).

- PCR products are then sequenced using primers labeled at their 5' end with T4 polynucleotide kinase, employing SequiTherm Polymerase. (Epicentre Technologies).
- 15 The intron-exon borders of selected exons is also determined and genomic PCR products analyzed to confirm the results. PCR products harboring suspected mutations are then cloned and sequenced to validate the results of the direct sequencing. PCR products is

-131-

cloned into T-tailed vectors as described in Holton *et al.*, *Nucleic Acids Res.*, 19: 1156 (1991) and sequenced with T7 polymerase (United States Biochemical). Affected individuals are identified by mutations not present in unaffected individuals.

Genomic rearrangements may also be determined. Genomic clones are
5 nick-translated with digoxigenin deoxyuridine 5' triphosphate (Boehringer Mannheim), and FISH is performed as described in Johnson *et al.*, *Methods Cell Biol.* 35: 73-99 (1991). Hybridization with the labeled probe is carried out using a vast excess of human cot-1 DNA for specific hybridization to the corresponding genomic locus.

Chromosomes are counterstained with 4,6-diamino-2-phenylidole and propidium
10 iodide, producing a combination of C-and R-bands. Aligned images for precise mapping are obtained using a triple-band filter set (Chroma Technology, Brattleboro, VT) in combination with a cooled charge-coupled device camera (Photometrics, Tucson, AZ) and variable excitation wavelength filters. *Id.* Image collection, analysis and chromosomal fractional length measurements are performed using the ISee Graphical
15 Program System. (Inovision Corporation, Durham, NC.) Chromosome alterations of the genomic region hybridized by the probe are identified as insertions, deletions, and translocations. These alterations are used as a diagnostic marker for an associated disease.

**Example 7: Method of Detecting Abnormal Levels of a Polypeptide in a Biological
20 Sample**

Antibody-sandwich ELISAs are used to detect polypeptides in a sample, preferably a biological sample. Wells of a microtiter plate are coated with specific antibodies, at a final concentration of 0.2 to 10 µg/ml. The antibodies are either
25 monoclonal or polyclonal and are produced by the method described above. The wells are blocked so that non-specific binding of the polypeptide to the well is reduced. The coated wells are then incubated for > 2 hours at RT with a sample containing the polypeptide. Preferably, serial dilutions of the sample should be used to validate results. The plates are then washed three times with deionized or distilled water to remove unbound polypeptide. Next, 50 µl of specific antibody-alkaline phosphatase conjugate,
30 at a concentration of 25-400 ng, is added and incubated for 2 hours at room temperature. The plates are again washed three times with deionized or distilled water to remove unbound conjugate. 75 µl of 4-methylumbelliferyl phosphate (MUP) or p-nitrophenyl

-132-

phosphate (NPP) substrate solution are added to each well and incubated 1 hour at room temperature.

The reaction is measured by a microtiter plate reader. A standard curve is prepared, using serial dilutions of a control sample, and polypeptide concentrations are plotted on the X-axis (log scale) and fluorescence or absorbance on the Y-axis (linear scale). The concentration of the polypeptide in the sample is calculated using the standard curve.

Example 8: Formulating a Polypeptide

The secreted polypeptide composition will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the secreted polypeptide alone), the site of delivery, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

As a general proposition, the total pharmaceutically effective amount of secreted polypeptide administered parenterally per dose will be in the range of about 1 $\mu\text{g}/\text{kg}/\text{day}$ to 10 $\text{mg}/\text{kg}/\text{day}$ of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01 $\text{mg}/\text{kg}/\text{day}$, and most preferably for humans between about 0.01 and 1 $\text{mg}/\text{kg}/\text{day}$ for the hormone. If given continuously, the secreted polypeptide is typically administered at a dose rate of about 1 $\mu\text{g}/\text{kg}/\text{hour}$ to about 50 $\text{mg}/\text{kg}/\text{hour}$, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect.

Pharmaceutical compositions containing the secreted protein of the invention are administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

The secreted polypeptide is also suitably administered by sustained-release systems. Suitable examples of sustained-release compositions include semipermeable polymer matrices in the form of shaped articles, e. g., films, or microcapsules. Sustained-release matrices include polylactides (U. S. Pat. No.3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman, U. et al., Biopolymers 22: 547-556 (1983)), poly (2-hydroxyethyl methacrylate) (R. Langer et al., J. Biomed. Mater. Res. 15: 167-277 (1981), and R. Langer, Chem. Tech. 12: 98-105 (1982)), ethylene vinyl acetate (R. Langer et al.) or poly-D- (-)-3-hydroxybutyric acid (EP 133,988). Sustained-release compositions also include liposomally entrapped polypeptides. Liposomes containing the secreted polypeptide are prepared by methods known per se: DE Epstein et al., Proc. Natl. Acad. Sci. USA 82: 3688-3692 (1985); Hwang et al., Proc. Natl. Acad. Sci. USA 77: 4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U. S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal secreted polypeptide therapy.

For parenteral administration, in one embodiment, the secreted polypeptide is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, I. e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation.

For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to polypeptides. Generally, the formulations are prepared by contacting the polypeptide uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate,

succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten residues) polypeptides, e. g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, 5 aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.

The secreted polypeptide is typically formulated in such vehicles at a 10 concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of polypeptide salts.

Any polypeptide to be used for therapeutic administration can be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e. g., 0.2 15 micron membranes). Therapeutic polypeptide compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

Polypeptides ordinarily will be stored in unit or multi-dose containers, for example, sealed ampules or vials, as an aqueous solution or as a lyophilized formulation 20 for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1 % (w/v) aqueous polypeptide solution, and the resulting mixture is lyophilized. The infusion solution is prepared by reconstituting the lyophilized polypeptide using bacteriostatic Water-for-Injection.

The invention also provides a pharmaceutical pack or kit comprising one or more 25 containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Associated with such container (s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the polypeptides of the present invention may be employed in conjunction with other therapeutic compounds. 30

Example 9: Method of Treating Decreased Levels of the Polypeptide

It will be appreciated that conditions caused by a decrease in the standard or normal expression level of a secreted protein in an individual can be treated by administering the polypeptide of the present invention, preferably in the secreted form.

Thus, the invention also provides a method of treatment of an individual in need of an increased level of the polypeptide comprising administering to such an individual a pharmaceutical composition comprising an amount of the polypeptide to increase the activity level of the polypeptide in such an individual.

5 For example, a patient with decreased levels of a polypeptide receives a daily dose 0.1-100 $\mu\text{g}/\text{kg}$ of the polypeptide for six consecutive days. Preferably, the polypeptide is in the secreted form. The exact details of the dosing scheme, based on administration and formulation, are provided above.

Example 10: Method of Treating Increased Levels of the Polypeptide

10 Antisense technology is used to inhibit production of a polypeptide of the present invention. This technology is one example of a method of decreasing levels of a polypeptide, preferably a secreted form, due to a variety of etiologies, such as cancer.

For example, a patient diagnosed with abnormally increased levels of a polypeptide is administered intravenously antisense polynucleotides at 0.5, 1.0, 1.5, 2.0
15 and 3.0 mg/kg day for 21 days. This treatment is repeated after a 7-day rest period if the treatment was well tolerated. The formulation of the antisense polynucleotide is provided above.

Example 11: Method of Treatment Using Gene Therapy

One method of gene therapy transplants fibroblasts, which are capable of
20 expressing a polypeptide, onto a patient. Generally, fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at room temperature over night. After 24 hours at room
25 temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e. g., Ham's F12 media, with 10% FBS, penicillin and streptomycin) is added. The flasks are then incubated at 37°C for approximately one week.

At this time, fresh media is added and subsequently changed every several days.
30 After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks. pMV-7 (Kirschmeier, P. T. et al., DNA, 7: 219-25 (1988)), flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf

-136-

intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads.

The cDNA encoding a polypeptide of the present invention can be amplified using PCR primers which correspond to the 5' and 3' end sequences respectively as set forth in Example 1. Preferably, the 5' primer contains an EcoRI site and the 3' primer includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the amplified EcoRI and HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is then used to transform bacteria HB 101, which are then plated onto agar containing kanamycin for the purpose of confirming that the vector has the gene of interest properly inserted.

The amphotropic pA317 or GP+aml2 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the gene is then added to the media and the packaging cells transduced with the vector. The packaging cells now produce infectious viral particles containing the gene (the packaging cells are now referred to as producer cells).

Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media, containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media.

If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his. Once the fibroblasts have been efficiently infected, the fibroblasts are analyzed to determine whether protein is produced.

The engineered fibroblasts are then transplanted onto the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads.

30 Example 12: Method of Treatment Using Gene Therapy-*In Vivo*

Another aspect of the present invention is using *in vivo* gene therapy methods to treat disorders, diseases and conditions. The gene therapy method relates to the introduction of naked nucleic acid (DNA, RNA, and antisense DNA or RNA) sequences into an animal to increase or decrease the expression of the polypeptide.

-137-

The polynucleotide of the present invention may be operatively linked to a promoter or any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques and methods are known in the art, see, for example, W0 90/11092, W0 98/11779; U. S. Patent 5,693,622; 5 5,705,151; 5,580,859; Tabata H. et al. (1997) *Cardiovasc. Res.* 35 (3): 470-479, Chao J et al. (1997) *Pharmacol. Res.* 35 (6): 517-522, Wolff J. A. (1997) *Neuromuscul. Disord.* 7 (5): 314-318, Schwartz B. et al. (1996) *Gene Ther.* 3 (5): 405-411, Tsurumi Y. et al. (1996) *Circulation* 94 (12): 3281-3290 (incorporated herein by reference).

The polynucleotide constructs may be delivered by any method that delivers 10 injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, intestine and the like). The polynucleotide constructs can be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

The term "naked" polynucleotide, DNA or RNA, refers to sequences that are free from any delivery vehicle that acts to assist, promote, or facilitate entry into the cell, 15 including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the polynucleotides of the present invention may also be delivered in liposome formulations (such as those taught in Felgner P. L. et al. (1995) *Ann. NY Acad. Sci.* 772: 126-139 and Abdallah B. et al. (1995) *Biol. Cell* 85 (1): 1-7) which can be prepared by methods well known to those skilled in the art.

20 The polynucleotide vector constructs used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Any strong promoter known to those skilled in the art can be used for driving the expression of DNA. Unlike other gene therapies techniques, one major advantage of introducing naked nucleic acid sequences into target 25 cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The polynucleotide construct can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, 30 thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue

-138-

ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

10 For the naked polynucleotide injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 $\mu\text{g}/\text{kg}$ body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg . Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration. The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked polynucleotide constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The dose response effects of injected polynucleotide in muscle *in vivo* is determined as follows. Suitable template DNA for production of mRNA coding for polypeptide of the present invention is prepared in accordance with a standard recombinant DNA methodology. The template DNA, which may be either circular or linear, is either used as naked DNA or complexed with liposomes. The quadriceps muscles of mice are then injected with various amounts of the template DNA.

Five to six week old female and male Balb/C mice are anesthetized by intraperitoneal injection with 0.3 ml of 2.5% Avertin. A 1.5 cm incision is made on the anterior thigh, and the quadriceps muscle is directly visualized. The template DNA is injected in 0.1 ml of carrier in a 1 cc syringe through a 27 gauge needle over one minute, approximately 0.5 cm from the distal insertion site of the muscle into the knee and about

-139-

0.2 cm deep. A suture is placed over the injection site for future localization, and the skin is closed with stainless steel clips.

After an appropriate incubation time (e. g., 7 days) muscle extracts are prepared by excising the entire quadriceps. Every fifth 15 um cross-section of the individual
5 quadriceps muscles is histochemically stained for protein expression. A time course for protein expression may be done in a similar fashion except that quadriceps from different mice are harvested at different times. Persistence of DNA in muscle following injection may be determined by Southern blot analysis after preparing total cellular DNA and HIRT supernatants from injected and control mice.

10 The results of the above experimentation in mice can be use to extrapolate proper dosages and other treatment parameters in humans and other animals using naked DNA.

Example 13: Transgenic Animals

The polypeptides of the invention can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, hamsters, guinea
15 pigs, pigs, micro-pigs, goats, sheep, cows and non-human primates, e. g., baboons, monkeys, and chimpanzees may be used to generate transgenic animals. In a specific embodiment, techniques described herein or otherwise known in the art, are used to express polypeptides of the invention in humans, as part of a gene therapy protocol.

Any technique known in the art may be used to introduce the transgene (i. e.,
20 polynucleotides of the invention) into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection (Paterson et al., Appl. Microbiol. Biotechnol. 40: 691-698 (1994); Carver et al., Biotechnology (NY) 11: 1263-1270 (1993); Wright et al., Biotechnology (NY) 9: 830-834 (1991); and Hoppe et al., U. S. Patent 4,873,191 (1989)); retrovirus mediated gene
25 transfer into germ lines (Van der Putten et al., Proc. Natl. Acad. Sci., USA 82: 6148-6152 (1985)), blastocysts or embryos; gene targeting in embryonic stem cells (Thompson et al., Cell 56: 313-321 (1989)); electroporation of cells or embryos (Lo, 1983, Mol Cell. Biol. 3: 1803-1814 (1983)); introduction of the polynucleotides of the invention using a gene gun (see, e. g., Ulmer et al., Science 259: 1745 (1993)); introducing nucleic acid
30 constructs into embryonic pluripotent stem cells and transferring the stem cells back into the blastocyst; and sperm mediated gene transfer (Lavitano et al., Cell 57: 717-723 (1989); etc. For a review of such techniques, see Gordon, "Transgenic Animals," Intl. Rev. Cytol. 115: 171-229 (1989), which is incorporated by reference herein in its entirety.

-140-

Any technique known in the art may be used to produce transgenic clones containing polynucleotides of the invention, for example, nuclear transfer into enucleated oocytes of nuclei from cultured embryonic, fetal, or adult cells induced to quiescence (Campell et al., Nature 380: 64-66 (1996); Wilmut et al., Nature 385: 810813 (1997)).

5 The present invention provides for transgenic animals that carry the transgene in all their cells, as well as animals which carry the transgene in some, but not all their cells, I. e., mosaic animals or chimeric. The transgene may be integrated as a single transgene or as multiple copies such as in concatamers, e. g., head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a
10 particular cell type by following, for example, the teaching of Lasko et al. (Lasko et al., Proc. Natl. Acad. Sci. USA 89: 6232-6236 (1992)). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. When it is desired that the polynucleotide transgene be integrated into the chromosomal site of the endogenous
15 gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous gene. The transgene may also be selectively introduced into a particular cell type, thus
20 inactivating the endogenous gene in only that cell type, by following, for example, the teaching of Gu et al. (Gu et al., Science 265: 103-106 (1994)). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant
25 gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to verify that integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques which include, but are not limited to, Northern blot analysis of tissue samples
30 obtained from the animal, in situ hybridization analysis, and reverse transcriptase-PCR (rt-PCR). Samples of transgenic gene-expressing tissue may also be evaluated immunocytochemically or immunohistochemically using antibodies specific for the transgene product.

-141-

Once the founder animals are produced, they may be bred, inbred, outbred, or crossbred to produce colonies of the particular animal. Examples of such breeding strategies include, but are not limited to: outbreeding of founder animals with more than one integration site in order to establish separate lines; inbreeding of separate lines in order to produce compound transgenics that express the transgene at higher levels because of the effects of additive expression of each transgene; crossing of heterozygous transgenic animals to produce animals homozygous for a given integration site in order to both augment expression and eliminate the need for screening of animals by DNA analysis; crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; and breeding to place the transgene on a distinct background that is appropriate for an experimental model of interest.

Transgenic animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

Example 14: Knock-Out Animals

Endogenous gene expression can also be reduced by inactivating or "knocking out" the gene and/or its promoter using targeted homologous recombination. (E. g., see Smithies et al., *Nature* 317: 230-234 (1985); Thomas & Capecchi, *Cell* 51: 503-512 (1987); Thompson et al., *Cell* 5: 313-321 (1989); each of which is incorporated by reference herein in its entirety). For example, a mutant, non-functional polynucleotide of the invention (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous polynucleotide sequence (either the coding regions or regulatory regions of the gene) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express polypeptides of the invention *in vivo*. In another embodiment, techniques known in the art are used to generate knockouts in cells that contain, but do not express the gene of interest. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the targeted gene. Such approaches are particularly suited in research and agricultural fields where modifications to embryonic stem cells can be used to generate animal offspring with an inactive targeted gene (e. g., see Thomas & Capecchi 1987 and Thompson 1989, *supra*). However this approach can be routinely adapted for use in humans provided the

-142-

recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors that will be apparent to those of skill in the art.

In further embodiments of the invention, cells that are genetically engineered to express the polypeptides of the invention, or alternatively, that are genetically engineered
5 not to express the polypeptides of the invention (e. g., knockouts) are administered to a patient *in vivo*. Such cells may be obtained from the patient (I. e., animal, including human) or an MHC compatible donor and can include, but are not limited to fibroblasts, bone marrow cells, blood cells (e. g., lymphocytes), adipocytes, muscle cells, endothelial cells etc. The cells are genetically engineered *in vitro* using recombinant DNA techniques
10 to introduce the coding sequence of polypeptides of the invention into the cells, or alternatively, to disrupt the coding sequence and/or endogenous regulatory sequence associated with the polypeptides of the invention, e. g., by transduction (using viral vectors, and preferably vectors that integrate the transgene into the cell genome) or transfection procedures, including, but not limited to, the use of plasmids, cosmids,
15 YACs, naked DNA, electroporation, liposomes, etc.

The coding sequence of the polypeptides of the invention can be placed under the control of a strong constitutive or inducible promoter or promoter/enhancer to achieve expression, and preferably secretion, of the polypeptides of the invention. The engineered cells which express and preferably secrete the polypeptides of the invention can be
20 introduced into the patient systemically, e. g., in the circulation, or intraperitoneally.

Alternatively, the cells can be incorporated into a matrix and implanted in the body, e. g., genetically engineered fibroblasts can be implanted as part of a skin graft; genetically engineered endothelial cells can be implanted as part of a lymphatic or vascular graft. (See, for example, Anderson et al. U. S. Patent 5,399,349; and Mulligan &
25 Wilson, U. S. Patent 5,460,959 each of which is incorporated by reference herein in its entirety).

When the cells to be administered are non-autologous or non-MHC compatible cells, they can be administered using well known techniques which prevent the development of a host immune response against the introduced cells. For example, the
30 cells may be introduced in an encapsulated form which, while allowing for an exchange of components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

Transgenic and "knock-out" animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function

-143-

of polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

All patents, patent publications, and other published references mentioned herein
5 are hereby incorporated by reference in their entireties as if each had been individually
and specifically incorporated by reference herein. While preferred illustrative
embodiments of the present invention are described, one skilled in the art will appreciate
that the present invention can be practiced by other than the described embodiments,
which are presented for purposes of illustration only and not by way of limitation. The
10 present invention is limited only by the claims that follow.

-144-

CLAIMS

We claim:

1. An isolated nucleic acid molecule comprising
 - (a) a nucleic acid molecule comprising a nucleic acid sequence that encodes
5 an amino acid sequence of SEQ ID NO: 172 through 295;
 - (b) a nucleic acid molecule comprising a nucleic acid sequence of SEQ ID
NO: 1 through 171;
 - (c) a nucleic acid molecule that selectively hybridizes to the nucleic acid
molecule of (a) or (b); or
 - 10 (d) a nucleic acid molecule having at least 60% sequence identity to the nucleic
acid molecule of (a) or (b).
2. The nucleic acid molecule according to claim 1, wherein the nucleic acid
molecule is a cDNA.
15
3. The nucleic acid molecule according to claim 1, wherein the nucleic acid
molecule is genomic DNA.
4. The nucleic acid molecule according to claim 1, wherein the nucleic acid
20 molecule is a mammalian nucleic acid molecule.
5. The nucleic acid molecule according to claim 4, wherein the nucleic acid
molecule is a human nucleic acid molecule.
- 25 6. A method for determining the presence of a breast specific nucleic acid
(BSNA) in a sample, comprising the steps of:
 - (a) contacting the sample with the nucleic acid molecule according to claim 1
under conditions in which the nucleic acid molecule will selectively hybridize to a breast
specific nucleic acid; and
 - 30 (b) detecting hybridization of the nucleic acid molecule to a BSNA in the
sample, wherein the detection of the hybridization indicates the presence of a BSNA in
the sample.
7. A vector comprising the nucleic acid molecule of claim 1.

-145-

8. A host cell comprising the vector according to claim 7.
9. A method for producing a polypeptide encoded by the nucleic acid molecule
5 according to claim 1, comprising the steps of (a) providing a host cell comprising the
nucleic acid molecule operably linked to one or more expression control sequences, and
(b) incubating the host cell under conditions in which the polypeptide is produced.
10. A polypeptide encoded by the nucleic acid molecule according to claim 1.
10
11. An isolated polypeptide selected from the group consisting of:
(a) a polypeptide comprising an amino acid sequence with at least 60%
sequence identity to of SEQ ID NO: 172 through 295; or
(b) a polypeptide comprising an amino acid sequence encoded by a nucleic
15 acid molecule comprising a nucleic acid sequence of SEQ ID NO: 1 through 171.
12. An antibody or fragment thereof that specifically binds to the polypeptide
according to claim 11.
- 20 13. A method for determining the presence of a breast specific protein in a
sample, comprising the steps of:
(a) contacting the sample with the antibody according to claim 12 under
conditions in which the antibody will selectively bind to the breast specific protein; and
(b) detecting binding of the antibody to a breast specific protein in the sample,
25 wherein the detection of binding indicates the presence of a breast specific protein in the
sample.
14. A method for diagnosing and monitoring the presence and metastases of
breast cancer in a patient, comprising the steps of:
30 (a) determining an amount of the nucleic acid molecule of claim 1 or a
polypeptide of claim 11 in a sample of a patient; and
(b) comparing the amount of the determined nucleic acid molecule or the
polypeptide in the sample of the patient to the amount of the breast specific marker in a
normal control; wherein a difference in the amount of the nucleic acid molecule or the

-146-

polypeptide in the sample compared to the amount of the nucleic acid molecule or the polypeptide in the normal control is associated with the presence of breast cancer.

15. A kit for detecting a risk of cancer or presence of cancer in a patient, said
5 kit comprising a means for determining the presence the nucleic acid molecule of claim 1
or a polypeptide of claim 11 in a sample of a patient.

16. A method of treating a patient with breast cancer, comprising the step of
administering a composition according to claim 12 to a patient in need thereof, wherein
10 said administration induces an immune response against the breast cancer cell expressing
the nucleic acid molecule or polypeptide.

17. A vaccine comprising the polypeptide or the nucleic acid encoding the
polypeptide of claim 11.

15

SEQUENCE LISTING

<110> diaDexus, Inc.
 Salceda, Susana
 Macina, Roberto
 Hu, Ping
 Recipon, Herve
 Karra, Kalpana
 Cafferkey, Robert
 Sun, Yongming
 Liu, Chenghua

<120> Compositions and Methods Relating to Breast Specific Genes and Proteins

<130> DEX-0306

<150> 60/268,292
 <151> 2001-02-13

<160> 295

<170> PatentIn version 3.1

<210> 1
 <211> 591
 <212> DNA
 <213> Homo sapien

<400> 1
 gctcttctctg tctacaaagg ggactgctca cagtggcctc agcttgggtgg ttttgagggg 60
 ccgccccccg gccctccata agggatcctt gggcctgaga attctgcac tgcattgga 120
 tggatgtaca gcctcaaatg gaagtgagtc ccacgggaga tgggtccgag gtccaggctg 180
 tggccatcca gccccctgtg gcttgtccag cctctgtgca cccctgggtgt cttcactcca 240
 ggggcagaca gtagccactg cagttccttt ctctgtgaga taacagtagt gatagcagct 300
 ggggctaaca ggctaggctt agtgtcctgc gcatttggtc agcttctcac tccatcctcc 360
 ctaaagcaat ggggaggccc ccactagccc agttttcagg aagtcaactg ggaggttaga 420
 tggggggccag aggtcccaca gctactgatg gcccgagcca ggttgagctt tcctggatgt 480
 ccagtccgga tcccacttgc agatctcatg ctctcagata ggtgggacaa gttcttttgt 540
 cacagtgtctg gctctgtcct gaggcctcat tgctggctgg tgtgctctgc t 591

<210> 2
 <211> 2754
 <212> DNA
 <213> Homo sapien

<400> 2
 gccagaagca gcctcagctt ggcaagggtg ggagatgact gctgttcct tgcatttgg 60

2

ggaaaaacagg ctccctcggg agctcgatga tcctcttttg atcttgtgtg acctcctgga 120
 gagtggatga cgctgggtggc cttagctttt ctagacagtg taaattgcac tgggcatgt 180
 cccagagca gggcaaggtc tctagagcgg gtctcccaca tgactggctt cacacaggca 240
 cttccgctcg ggttgcacgc tctgtgtcat cttaccggtc cagggttgca ggtaggaaat 300
 gtttgtaccc tcttctgatt gccacctcct tcccatcgcc ccttagggac agggcttgag 360
 ggccagttag gcgctgggtca ggcaccccag gcctccttgg gacctgcca ggggcaccct 420
 gagagctcct gaaacccccca cttagcttcc agacctttct gcaaaagctc ctctggtctt 480
 tcctccctcc cccaatctat gggtcacagc taacagatct gagggcaact gctgtgctag 540
 tggccagggc tgcaacctgc atccccggct ctgccacttt agggccttct agaggcagtg 600
 tccttaggaa gtagctctga ggcattgggtt ttctgctcct gtgcagggca gctgatggga 660
 taaggtgggg aaggacggtc agtgcttggg cccagctgg ccagcctggc gatggggaaa 720
 ccaaacctatg tccccagcg aaggccaga gtgggaacct gtctcatgc ccttcgctct 780
 gaggagccct gaggtgggca gcaggggcca ggggaagtct tcaggccttc atcaaagaga 840
 acaacatcct cagctccgca cccctcatcc tgtatcagca cttaccgggtg tgtgactgcc 900
 cttgtcagct agcatacggg gggcccacct ggcccactgg ctgtttatgc cactgattta 960
 tgatagggaa tattatcttt gaaccaatg aagtgttttc tccccatca caaaaaaaaa 1020
 aattcttatt ttagtagac atgtatttac caaaaatag tactcaatta ttgtattttg 1080
 gattttatca atttaaaaat tgtggaatt tgtttgctct tacgccaaca taatattgat 1140
 tttgcctctt ggctctgaaa gcccaaaaata tttaccgtct agcccgttac agaaaaagtc 1200
 tgctgactac tgagccagac ctccattacc tccatccctg ttggattatt taaagaaagc 1260
 ctgagacagt aagggtcttt ttaaaagaat aaaatgactt ggtttgcgct tggagcagg 1320
 ggaagcattc agatgagcgg tttctgcatt aacctgcct atcacgcac tcgtgtcctg 1380
 tgtggctggc gagccccct tggaaaggtc tgggtgctca gctggctcct gcagagtcca 1440
 ccccgctcgt tgggtgggaat gcagagccct ttgctttcct tcttgccgcc tgcttctgt 1500
 tcctggggac ccgctgggccc tttggtctgc atccccggc caggtcctc agggttgatg 1560
 cgtggagaag gactttgagc agtgggtggc agcagtggcc tcctggccag ctcacactct 1620
 tgtcctggga ggggcagcct gatctcacct ccacctagta ccttggggac tgaggacctt 1680
 ttgcttctc tggagcctgc aagcctctc ccatgtgtcc agctgctctt cctgctacaa 1740
 aggggactgc tcacagtggc ctgagcttgg tggttttgag gggccgcccc cggccctcc 1800
 ataagggtat cctgggcctg agaattctgc atctgccatt ggaggatgga cagcctcaaa 1860

tggaggagt cccacgggag atgggtccga ggtccggctg tggccatcca gccccctgtg 1920
 gcttgtccag cctctgtgca cccctgggtg cttcactcca ggggcagaca gcagccactg 1980
 cagttccttt cttcgtgagt aacagtagtg atagcagctg gggctaacag gctaggcttt 2040
 gtgttctgcg catttgggtca gcttctcact cgatcctccc taaagcaatg gggaggcccc 2100
 cactagccca gttttcagga agtcaactgg gaggttagat gggggccagg gtccccacagc 2160
 tactgatggc ccgagccagg ttgagcttcc tgggtgtccag tccggatccc acttgcagat 2220
 ctcatgctct cagataggtg ggacaagttc ttttgcaca gtgctggctc tgtcctgagg 2280
 cctcattgct ggctgggtgt gctctgctgg gaaaagcttt gcggggcttg cttggttaac 2340
 cacagaagag aaggggactg tttgggggtc ctctctgcag cctccccgtg ctgggtggaa 2400
 gcacggttac tgtgttctct aatgttcatt tatttaaaat gatttctttc taaagatgta 2460
 acctccacac ctttctccag attgggtgac tcttttctaa aggtgggtggg agtatctgtc 2520
 ggggtgggtg ggccttggga tgggtcaggt ggggtgtgaga ggtcctgggg aggtgggctg 2580
 tgagctcaaa gttgtcctac tgccatgttt ttgtacctga aataaagcat attttgcact 2640
 tgttactgta ccatagtgcg gacgagaagt ctgtatgtgg gatctgtgct tgggttagaa 2700
 tgcaataaaa actcacattt gtaagaaaa aaaaaaaaaat aaaaagatgc ggcc 2754

<210> 3
 <211> 856
 <212> DNA
 <213> Homo sapien

<400> 3
 acgttaaaat taagaactta ggctttgggt taaaaaaca taaatgaagt gaaaaaaca 60
 agccacagag taaaagaaga tacttgcagc aagtataaa ggattagtat ccaggatata 120
 taaagactgt tattgagtca atgtgaaagg gagaaaaaca cctgaagcaa agaatggatg 180
 ccggcattaa ataggcactt caaagaggaa ccatgaacga ccaaaatcaa gtgagttagt 240
 gaccagttcc cattagtaat taggaaatag caaattaaga ccacaaagag ggcagtgagg 300
 gtggctcaca cacctctaat ctgagcggct tgggagtcca ggcccagagg atcccttgag 360
 gccaggaggt ggagtctagc ctgggaaaca tagcaagacc ctgtctctac aaaaaataa 420
 ataaataaaa taagaaaaaa gtaaaccaca aggagatgac ttaccaccag gcaaaaatat 480
 taaagtatgc taataccaag tatcaagaag aatgaagcaa gatagctcaa atatgctttt 540
 gaaggaaata tactgggctt ccattcattc tgaataccc cttatttaag atactctatt 600
 atattaaata cagttccaaa acaaaagaaa tccaaagaac aaaaaactaa cccaatactt 660

4

ttatcacttg taattgtata ttacaccata ttgaaagata tattttacga cttattagag 720
 aacgattttt aaattggata tcaactctgtg catacaaata aaataaagtg attaaggttc 780
 taacaaaaaa acaaaccaca acaccaaagg ctttttaagg gggggaggaa taaggaaagg 840
 ggcccaaaaa agggac 856

<210> 4
 <211> 1580
 <212> DNA
 <213> Homo sapien

<400> 4
 gtcccttttt tgggccctt tccttattcc tccccctt aaaaagcctt tgggtttgtg 60
 gtttgttttt ttgttagaac cttaatcact ttattttatt ttgtatgac aagagtgata 120
 tccaattaa aatcgttctc taataagtct aaaatatatc tttcaatatg tgtaataac 180
 aaattacaag tgataaaagt attgggtag tttttgttc tttgatttct tttgttttg 240
 aactgtattt aatataatag agtatcttaa ataaggggta tttcagaatg aatgaagccc 300
 agtatatttc cttcaaaagc atatttgagc tatcttgctt cattcttctt gatacttgg 360
 attagcatac tttaatattt ttgcttggg gtaagtcatc tccttgggtt ttactttttt 420
 cttattttat ttatttattt tttttagag acaggtctt gctatgttcc ccaggctaga 480
 ctccacctcc tggcctcaag ggatcctctg ggcctggact cccaagccgc tgagattaga 540
 ggtgtgtgag ccaccctcac tgcctcttt gtggtcttaa tttgctattt cctaattact 600
 aatgggaact ggtcacctac tcacttgatt ttggtcgttc atggttcctc tttgaagtgc 660
 ctatttaatg ccggcatcca ttctttgctt caggtgtttt tctcccttcc acattgactc 720
 aataacagtc tttatatatc ctggatacta atcctttatc acttgctgca agtatcttct 780
 tttactctgt ggcttgtttt tttcacttca tttattgttt tttaaaccaa agcctaagtt 840
 cttaatttta acgtacttga actgacattt tctaccctgg cccctccca cccttagttc 900
 ccagacacct cttatgatct ggggtcgagg agcccgcctt cctggcggtc tcagctgggg 960
 cctggggagc gaaggcggcg ggcgctcgcg ggaggagctg cgcgattcgg atgctgggga 1020
 ggtgaagctc gcggggccgc caggccgccg gggtaggaa ggccgggagg ccgcgggggt 1080
 ccacggcgcg gaggagaccg caggcaccgg gcacagccct cgcccatcgc cgagaccgg 1140
 caggcccagg agccagaggg cggcggtg agagggaaacc gcctccaaag gacgcccctg 1200
 ccctcccga ggcatagtcg caggcgccag tcccggctcg agccagctgg ggggtgctcc 1260
 ggggagctga gccgggggag ggccgggccg cccaacggat caataggggg gtttctcca 1320

5

ggttgccggt tctctggccc gcgacgccct acccgccgga gccgcccaac cgcaagcccc 1380
 gccccgaagg cgggtgcagc caggaaggcg gggcctgggt gcctctgggc gctggcgcca 1440
 agttcagagc cgcgccctgg gctgggcggt ggcgcccgcg tctgcacttc cccccctgcg 1500
 cgcctctgga gagcccgga gagacgcacc ctcaggtcgg ccaagaccga gaacaagcgg 1560
 gcgcgggcag cggagcccat 1580

<210> 5
 <211> 800
 <212> DNA
 <213> Homo sapien

<400> 5
 tggtcgcggc cgaggtacaa aggctttgag gtccatggac tatacttgtc ccatttatca 60
 tcccaggtgg tgctttgacc ctagggatac cctggctatt aagataaaaa gatttgtgga 120
 cattaataat atgaatatgt cagtaataat ccagcacaca ttgaaatatt gacacagatt 180
 accataatth gtgcaacatc ttataaacia tgcatttcc acagtagtct aaggcttcac 240
 cagcctggcc cactgtatct agactttagg ttcattttaa ttaattatgc tttccttctc 300
 tgtatcattt gggaagttga taaatatcac ttccttagat accttcattc agtgatata 360
 ctggctttta caattaaatt ggaaaaggta agtttctctt tggggggttg agagttggac 420
 catcaattct aatctacaaa aggaaattca tgatttctact ctgacgccta ggatctagcc 480
 aaggctggtc tgcagtatca gatgtccaaa ctcacttact attagccata ttttgtgagt 540
 cgtttgtcta aactttgtca aaaatgcctt tgccatgatt ttgttgctat ctggatttca 600
 aacatggaca gttaggaaga tgtgcattga agtaggaaaa ttttggcag catctgctgt 660
 tatttatttt ttaccacttc aaaaatggcc actgtctttt taacaaacac caacgacaac 720
 aacacacaaa acaaaaaaaaa acaccctgcg gcttaccctg gccctccttt tcctgttga 780
 attgtttccc cccaatcac 800

<210> 6
 <211> 956
 <212> DNA
 <213> Homo sapien

<400> 6
 atttataagg cccttcaaat ttgtggcttc ctttctcata cttctcaagt ataataaag 60
 ggggagaaaa accccaccat caacacaaaa gaaggctata aagactgtgc accttttaac 120
 aagtcaattt gtagtcagtc cctgggectg tctttttttt tttttaattt tgaagctacc 180

6

tgaggtttag aattccttca gccctagctg cttttattct gctttttatt taaacaaaa 240
 gagggggagg atctgaagga aactagtttt ctgtacaaag gctttgaggt ccatggacta 300
 tacttgccc atttatcatc ccaggtggtg ctttgaccct gccataccct ggctattaag 360
 ataaaaagat ttgtggacat taaaattatg aatatgtcag taataatcca gcacacattg 420
 aaatattgac acagattacc ataatttgtg caacatctta taaacaatgt catttccata 480
 gtagtctaag gcttcaccag cctggcccac tgtatctaga ctttaggttc attttaataa 540
 ttatgctttc cttctctgta tcatttggga agttgataaa tatcacttcc ttagatacct 600
 tcattcagtg atatatctgg cttttacaat taaattggaa aaggtaagtt tctctttggt 660
 gggttgagag ttggaccatc aattctaate tacaacagga aattcatgat ttcactctga 720
 cgcctaggat ctagccaagg ctggtctgca gtatcagatg tccaaactca tctactatta 780
 gccatatttt gtgagtcggt tgtctaaact ttgtcaaaaa tgcctttgcc atgattttgt 840
 tgctatctgg atttcaaaca tggacagtta ggaagatgtg cattgaagta ggaaaatttt 900
 gttcagattt gctgttattt attttttaaa taaaaatgg aatgtaaaa aaaaaa 956

<210> 7
 <211> 489
 <212> DNA
 <213> Homo sapien

<400> 7
 actatgtggt aacataatcc caccttctta gagctttggt cttctgaag gtgtatagat 60
 acagcttgtc ttgaaatgtc tttctccaca taatgaagca tgctgaatgc tgggaatctg 120
 gagcagcagc cctgggagcc ctgagttttg aagtgttttg gtttgcttca aaggtagaa 180
 gaacttgata tgtatggcaa acaactttag aatactagtt actcactaac atgaggcggg 240
 taatgttgct ctagattcta tattccagta aagccagctt ttcttattat tggagtaggc 300
 aaatgaatgg cattagaatt agtgggtggc ttgtaagttg tagttatagg cactttacca 360
 cttcctgccca ttagcaggca tccttgtttt ttcttctttt ccctctttgt tccttctttt 420
 cccttctccc ttatacattt tcttctctta ctttaattct ccttcctcct tactgtagat 480
 cccaagctt 489

<210> 8
 <211> 3190
 <212> DNA
 <213> Homo sapien

<400> 8
 ctctcattag cctgttcaga gtcttggggg aaattgagat ttttgagatt ttttttaaaa 60

actcaaatat ttactagtt tgctgccat tttatttctt ttacaaagca gaagcatata 120
 ccaatttatc acagtatfff agtaaatact gcaacattca tccttaaatg ttcaccaaga 180
 aaagcatctt tgtagtagtg ctggaaaact attcagaata tacagataaa aatgctgttc 240
 ttttaattgct tacattgctt cttcccataa aaagcaaaaa ggaatcagtg cttgctattg 300
 ctcttttctt tgaagttgta acaattgata catatattat gagttgactg gtcgattctg 360
 tacctggccc atcctttaga atgttcttgt catgtagcag tcctacgtac tcttttcatg 420
 agcagtcctgt gatctcactc tgtgagttca gctattactc gctcgtggga gcttaatcct 480
 ttcaaaatga agttgattta aaaagtcttc aggcagagta atcatgtag aggtggattt 540
 cgatggaaga aagtttagag agttaggagt ggggtagaa ttctagaatt tataaaggtc 600
 caggaagcat agcagtcagg ggcaaaaatt agcgtaatat ggagtaggca atagaggagc 660
 tactggagtc agaagtcact gcagagtgca acataggaag atggactcct agcttacatg 720
 agattccctg cagctgtaat atagacaatt cccacatggc tgttctacac agaattacct 780
 gctaagattt tttgtttatt tttgtttgag tggatatttc actccaattg tataatggaa 840
 atcagtgagg aaatagggtt taccttatat tcatgagttc tagtttctac tgttctgcta 900
 tgtgtttcta agcaagagca aaggatactt catacttttt togttatatg attgatcttc 960
 aaattgggat ttaccttttt caatatgttt taaagtagtc ttattcctct tttgatttgt 1020
 taaacaagca ttttagttca gctattgaat agccttccaa aaaattaatt cagccttgca 1080
 ggtaagtacc atactaagac ttaacccaa tagtttttaa tcattctgcc tttattccaa 1140
 actgtaaate tgtacacata agataaaaca tactaagtat tgcataaatt gttaacgtta 1200
 cagtaaattg ttatctgcag ggctgacaga cataatgttg gtgggcaact gtgatcctat 1260
 acatacatat atgcaaaagg ggattttaaa agtgcagatt atagagtaga ttgacaaatt 1320
 ttattttata ttcagttgct ctctctgctt ccatctgtgt tgctctctta gttgagagag 1380
 agttagccat ttgacgattt taagtcagtg ggaacttatt tttagttact caataaaatt 1440
 aatattttat ttgtatttta acttacagag taggttggtta ataacagctg aactgtgtaa 1500
 cattgttgct tcaaattgaa gtttatatta tgaacattca gaatcaatgc tcatgtagca 1560
 gcatattatt gagctatfff gagtttgaaa tgtggagaaa cgctaaacca tgtactatgt 1620
 gttacataa tcccaccttc ttagagcttt gttccttctg aagggtgata gatacagctt 1680
 gtcttgaat gtctttctcc acataatgaa gcatgctgaa tgctgggaat ctggagcagc 1740
 agccctggga gccctgagtt ttgaagtgtt ttggtttgct tcaaaggta gaagaacttg 1800

8

atatgatgg caaacaactt tagaatacta gttactcact aacatgaggc gggtaatggt 1860
 gctctagatt ctatattcca gtaaagccag cttttottat tattggagta ggcaaatgaa 1920
 tggcattaga attagtgggt ggcttgtaag ttgtagttat aggcacttta ccacttctg 1980
 ccattagcag gcatccttgt tttttcttct tttccctctt tgttccttct tttcccttct 2040
 tccttataca ttttcttctct ctactttaat tctccttctt ccttactgta gatcccaagc 2100
 ttctagctta ggtttgcaag tcatattgct tggccctcca cattcactga gaggtgaaga 2160
 taggctgacc ccctgtcctc ttacatttga gggatcatag actgctgtgt gaattctgga 2220
 aagtctcagg tccctaccag ggcactgaat ggcttctcaa tggctgtaga gacagtacag 2280
 ttttccaaag cagcctaatt catctggaca gctaccaggc actttgaaa gttggttcag 2340
 ttactactat gaggccataa tatatttgct ggtattaaaa ttcttcagaa ttggaattac 2400
 tatttgaaat aatattttgg ttgacttaag ttttgagaga caattctaaa attgatctag 2460
 agactcattc aatagcaatg tgacctttta aatacttaca ttaagtaaaa ctgccagtag 2520
 attaaatcat atatatatat atatatatat atatatatgt aagagcttcc tctatttact 2580
 actggtgaac ttcagtaatt tttagaggct aaataatggc cagaatgttt ttaagtgtgc 2640
 tcttttatta catgcttggt caggttttgt aattcagtag agaaaagttt aacctgtac 2700
 atttttgat gtaaaaagtc ttttaagtag tcttatcctt atttaaataa acagaataaa 2760
 attaccttga gtaggtctgt tattcttatt aaaatggaaa aatgctctgt aatgacttga 2820
 tctgttttta tttgagtga caattttgga agtattctt tatagtacaa ctttctatac 2880
 ctggattgat taagatcaga tgtgattcga gtagtccagc catatcttgt agcccttctt 2940
 tgaatgagag ggtggctgga gtggtctggt gctgggatat cacgggtgcta cagagcctga 3000
 catggtgact gtcactacat gttgagggat ggaaatagaa gtctctgaac ttcccatgta 3060
 atattaaagc tcttaacaaa atgagacaaa ctagagattc agttgagaga ttttatgtta 3120
 gagtgatctg aaaaaaagtt aatttctaaa ctgctatctt aatattatta tatttgagga 3180
 ctgatgctgt 3190

<210> 9
 <211> 672
 <212> DNA
 <213> Homo sapien

<400> 9
 ggtcggggcc gaggtactat tgctctggct cctggcctc tccttgetat gggcttacc 60
 ctcaagtcgc tctgtgattc aaagatgaac tgccaatcaa atgttctctt aatgaaagat 120

9

ccaatcactc tacagcatgt gtgtattcaa agaacctatc taagactttc ttttggtcat 180
 ggtgggaggc tgttgctgaa aacataccag agcccattgt ggaggtcagc tgacaggccg 240
 catgaccttg gcaatggact actggtcatc tgggactgct taggactgtg caatggaact 300
 tgggggcaaa actgatggag acagccaatg ggccttaaat ccagcaggca aagacagagt 360
 aagttcttat ttgtgtagcc cagggcttat caaagtgtgg ttcttggacc acgtgcatca 420
 gtatcagctg taagtatttg gcaaaatgca gattcccggg ccctgcacca aacagattga 480
 ctttgaatct ctgggggttg ggctaaaaaa aaagaaaaaa aaaccctaca ttttaacaa 540
 gctcttcaga tgacccttgt gtaagtttga gagcatctgc tggaaaacca ctagaatttg 600
 caaacggcac tcaaaatact ccagccagtc cactagccaa agaccagatc tgagaccgga 660
 tgggaaatta tc 672

<210> 10
 <211> 997
 <212> DNA
 <213> Homo sapien

<400> 10
 ggtcggggcc gaggtactat tgctctggct cctggccctc tccttgetat gggcttacc 60
 ctcaagtgcg tctgtgattc aaagatgaac tgccaatcaa atgttcctct aatgaaagat 120
 ccaatcactc tacagcatgt gtgtattcaa agaacctatc taagactttc ttttggtcat 180
 ggtgggaggc tgttgctgaa aacataccag agcccattgt ggaggtcagc tgacaggccg 240
 catgaccttg gcaatggact actggtcatc tgggactgct taggactgtg caatggaact 300
 tgggggcaaa actgatggag acagccaatg ggccttaaat ccagcaggca aagacagagt 360
 aagttcttat ttgtgtagcc cagggcttat caaagtgtgg ttcttggacc acgtgcatca 420
 gtatcagctg taagtatttg gcaaaatgca gattcccggg ccctgcacca aacagattga 480
 ctttgaatct ctgggggttg ggctaaaaaa aaagaaaaaa aaaccctaca ttttaacaa 540
 gctcttcaga tgacccttgt gtaagtttga gagcatctgc tggaaaacca ctagaatttg 600
 caaacggcca cctcaaaata ctccagccag tcccactaag ccaaagactt tcttttggtc 660
 atggtgggag gctgttctg aaaacatacc agagcccatt gtggagggtca gctgacaggc 720
 cgcatgacct tggcaatgga ctactggcca tctgggactg cttaggactg tgcaatggaa 780
 cttgggggca aaactgatgg agacagccaa tgggccttaa atccagcagg caaagacaga 840
 gtaagttctt atttgtgtag cccagggctt atcaaagtgt ggttcttggga ccacgtgcat 900
 cagtatcagc tgtaagtatt tggcaaatg cagattcccg ggcctgcac caaacagatt 960

10

gactttgaat ctctgggggt tgggctaaaa aaaaaa 997

<210> 11
 <211> 696
 <212> DNA
 <213> Homo sapien

<400> 11
 gccgcccggg caggtacaaa tggtgcccat gccattcatt tgactgtggg tggccctcta 60
 gtctagggct ctcttagtga atggttgtgg aaatatgatt tttctaagtt ccttcctttt 120
 ccttttgata gatgagtttg agatgatgga gtaggagtga ggccctcagg cacttctggg 180
 aaagacattc cacctgcaag cagcattttg agtaaagcac tgctgtgggt tgccgattta 240
 tgggccattt aatgttaggc taaagcacct ttaatcattt ttgtgtttt aagataatgt 300
 atttgtgaag tggataaaca ctggaaatag ggtgcttctt ctggaaagtt cagtgtaaaa 360
 cactaacaag gctttggcgg gtttatctgg ctttataaac aagtctgaaa aatggatgaa 420
 agctaaatat ataaagcagt tggttgtcta tcttttatca ttttttactc agatctgtat 480
 ttaacactta tttatttgtt agtttttaca ttcaaaagaa actacacttg gaactttggc 540
 taacattgta ggatattttt taattgttcc tacattttta agcatgattc atcattttgg 600
 taacttagat catttttaat ggtcttttct ttcaataacc agttacatca tgttttggga 660
 actccttggg tccatataag gtgaattggg gcaaaa 696

<210> 12
 <211> 3233
 <212> DNA
 <213> Homo sapien

<400> 12
 aacggctccta aggtagcgag agaatactac caggtgctag tttttccagt attgacttct 60
 gattactatt tccttttctc atctttagtt tttcaagatt tgctttacca aaatagtaaa 120
 gcctttatca tcagcttata ttgaataatg ttgtaattgg tttcaatcaa agtttctcct 180
 caggtacttg ggggcccta gccttctaag gaactcccag gcacctactt aacaaggcca 240
 gctacacact cagtatgtga taagcccat gatggatgca ggtagaatt caaagacctg 300
 gttggagtcc tagatgtgga gacaggatca tcaggtcaca cttgttagat gactaacact 360
 atcagtagaa gctcttgaga gattttccta acgcagcaag atttctgtga gtagaggat 420
 cctgggaggt atcctgggag gcagcctatt gacttgacca agtaagctga tcagggtgcc 480
 tcctctaccc actaaagaaa tgtgtaaaca ctagcaataa ttgctttatc ttaaactcct 540
 ggacatactc agttcctcca ttccactggt ctattgcaa tacctttggt gttttcttca 600

cactcctctt	ggcagcaaat	gtctgaaagt	atttcaattg	tgtaatgta	aggagttttt	660
tcatagcttc	agaaaagagg	gcagcaata	tgaagcctta	agttcaaat	aagtcattct	720
acctagaaat	acagaccca	gagcacattg	catgaaaata	cctgtactct	gcagttcctc	780
aaagcagtat	tcttcctgaa	aagccaaaca	ccacacctat	tttcctattt	gctaagaatc	840
agaataagca	cgttgtaaat	agtatccaaa	gcagattcta	aatgacata	gtaagaagcc	900
agattcaaat	tgtaaccaa	gaagacaata	gaaatccac	tttaccac	tgatcatcagt	960
tagaacaccc	ttgcaaaaac	tgtaaccact	taagcaattc	atctgatccc	agaagatcat	1020
accttctttt	gaaagtatag	gacagatata	agtgggaaac	gtcggcggtc	tgagcaacac	1080
aggataaatg	taggagggcc	ttaaaaaata	aatctcaatt	catacactgg	agcagcaaaa	1140
aactgagcag	gaaaggaaac	agaatccaaa	gtcatttttc	atatagctgt	tgtcaaatag	1200
tataaccttg	gtgtcttctt	tgagttgcct	ggacagtatt	tatgaaaca	aaaactaat	1260
gccccccatt	tggggacggg	gggaggggtt	cagacctcta	acctggattc	agagccttag	1320
aggccgagag	ggaatctgga	atctggtatt	actgagatcc	taggtaaaag	aaccagcctg	1380
gcagtctttc	ccacctcatt	ggtccgtgct	tttattttta	aacccaaaaa	aaaaaaaaaa	1440
acacaccctc	ttatgtagga	atttcccttt	tacaaataat	ttgacctggt	agaaataaac	1500
ttgcctgcct	gctcttaaat	gccagacagt	tggaagcaa	tgccgagggg	aagggtgcca	1560
gagccatgct	tgataggact	ttgaatattt	tctccttaat	taaagtacgt	tgcttgattt	1620
tagactataa	gatgctatgg	aggtctatcc	catgccatgc	caatgtgaat	tgctttgctt	1680
cagtaacaat	cagaagacca	gtccaacaga	aaataacttg	tcataattcc	accttagatt	1740
ctagacctct	catacctgca	gtgtacagaa	tatgtacatg	ttccaatgga	attcactatt	1800
tttggcttta	gtgtcaaaga	gattggttct	acaaggttca	tctgatttcc	cataacaagt	1860
aaatttata	atcctatgat	tctaaattca	atccccata	tagattctaa	gcatcaaatc	1920
aaaatcacag	acaaggggga	actggtcgag	aggggtctta	gttatttcaa	atccatgacc	1980
aaagtgtcca	aagacatgaa	actcttatac	ctgctgagca	tttcacttta	ctatacaaaa	2040
tgtcagctac	ccagttgcat	cctgtgacat	gatcagactg	tcaatgtgga	ccagtgcca	2100
ggagcatatt	tatgggcat	ttctgttcat	cattctttac	agagcattga	ggtttccac	2160
tgaaacagct	tctttagtca	gacgtctata	gattttacat	aaatttcat	ttaaatgcat	2220
taagttagat	ggccaattg	agcatctgaa	tgaatatagt	gggggttgg	ggtggtgcaa	2280
attctgctgg	ctttatgta	tggtttctt	cgtgttttt	cttggtttt	tctgcttct	2340

12

tctggcaagt gccctaaaag actggaacac tgtataaagt catagacata gaaccatatg 2400
 ggaaagccca gatgaaaaaa tggagaata aatcaagtt gtcaaagttc cagcaacagc 2460
 cctgacttct tcaggaatcc aagcaaattg aaagccaaga caaaatgtac aaatggtgcc 2520
 catgccattc atttgactgt ggggtgccct ctagtctagg gctctcttag tgaatggttg 2580
 tggaaatatz atttttctaa gttccttctt tttccttttg atagatgagt ttgagatgat 2640
 ggagtaggag tggggccctc aggcacttct ggtaaagaca ttccacctgc aagcagcatt 2700
 ttgagtaaag cactgctgtg gtttgccgat ttatggcca tttaatgta ggctaaagca 2760
 cctttaatca tttttggtgt ttttaagataa tgtatttgtg aagtggataa aactggaaa 2820
 tagggtgctt cttctggaaa gttcagtgtg aaacacaaac aaggctttgg cgggtttatc 2880
 tggctttata aacaagtctg aaaaatggat gaaagctaaa tatataaagc agttggttgt 2940
 ctatctttta tcatttttac tcagatctgt atttaacact tatttatttg ttagttttta 3000
 cattcaaaag aaatacactt tgaactttgg ctaacattgt aggatatttt ttaattgttt 3060
 ctacattttt aaagcatgat tcatcatttt tgtaaactta gatcattttt taattgtctt 3120
 ttcttttcca atagaccagt taccactcat gtgtctgcag aacctttta ttgtattcct 3180
 ataataaatg taaaatattt gtagcaaaaa aaaaaaaaaa aaaaaactcg gtc 3233

<210> 13
 <211> 847
 <212> DNA
 <213> Homo sapien

<400> 13
 actagactat gatatggact taccctcttg gtgccagccc tacaagctgc atgaccgtaa 60
 tcagcctgtg aactacgac atgcgccact cagcctgtgc cactcttggc ctgagccttc 120
 ggcctcttat gactgaggcg gacaactcac gctcaacaat ggcaaagact gctgcacat 180
 tgctagatca catcaatggt gccaccaact actctcttct tcacctacca acagtggact 240
 gactggcttc tatgactcct atccacctca tctgctcccc tagtcacgaa ctacaagaca 300
 ccacacacc ccagccgcag cgcaatgcc aaaggttcag cacacacggt gcgcaaaca 360
 cccaatgcgc gaccatcat catccataca tatctggcgc tgctacgagg acctacctac 420
 gccatgtcgc tcctgactac tctgctcac tgatggctcc tcctaccaac acgcgcttgg 480
 ctctgccag cctgcagccc acacacctgc gccctcctt ggacgacac cactcaccg 540
 ccgactgccc aacgcaccag ctgacggaca cagctccact ccaccgccc ccaatcacct 600
 cgcgacgccc ccagccctg ccctccata cccacggcct acaccacacg cgccacctc 660

13

aactgcgac cggctgcca tatctctcca cgtcccgcc tctgcccc ctcacacac 720
 gcagcatcca cccagacaac ccacactgca cgaccctca tcacagcccc tcaaagcct 780
 ccaccaccac acaccagcag tccccgcc caacacctaa taaacccac ccccgccgag 840
 cctcaca 847

<210> 14
 <211> 267
 <212> DNA
 <213> Homo sapien

<400> 14
 actgtagcag tgagctcaag tgttgggtgt atcagctcaa aacaccatgt gatgccaatc 60
 atctccacag gagcaatttg tttaccaaga atctaagaat taaatcttag aatgtattaa 120
 tgttaaattt ctgtgagatt atattgtagt cacgtagaat gtcctgactt gtaggaatac 180
 ccactaagga aatcagaaat cacggtagag cgtcagcaat ttactctcaa atggttcaga 240
 gaaagaaagt tctttgtagt aaagctt 267

<210> 15
 <211> 824
 <212> DNA
 <213> Homo sapien

<400> 15
 tggtcgcggc cgaggtagc tgggtggaaa gggcatttgg agctcattag aatgagacat 60
 agttaagagt cccattctca tcagtgtatt ccagactgag gaagaaatgg ggcagcagtc 120
 aggagagctc gggatthtga gtatagcaga atttaagtga aatggaaact acactcttta 180
 atttgttctt ccatggaatt gctthtctca tgcaagggtg gagccccag gagagccctt 240
 gtgaagggaa tgctgatttg tgtgaatata tgtaggtag taggtatcta gtgaggatga 300
 gttgggagga tgagttgggt aaggcgtgcc cctctgacac tgttctgggt ataaaagaca 360
 acatcatgat gagatcttca tctgaataaa actatgccct ggcctthtca gaaactgagg 420
 gcactgcagg tcccacagtg tgatggagtc caagctggga tcaactgcgag atgaggagtc 480
 agaggagtgg ctccggcagg catgggagct tcaggccctg agagagaaga cagaaattca 540
 gaaaacggag tggaaaagaa aaactgaag gaactgcatg aagagcacat ggctgagaag 600
 aaagagctac aggaggagaa ccagaggctc cagggcctcc ctgtctcagg atcagaagaa 660
 ggcaggctgc cagttccaag tgccagatca agcaccctcc gtgccagctg caggaacgag 720
 ctaggatcat tgcttccagg aggagagacc agccttggtc tcaaggaagg gcaccggacc 780
 aaaggggcaa gggggggaca cagagagat ccacaggaaa aatg 824

<210> 16
 <211> 1998
 <212> DNA
 <213> Homo sapien

<400> 16
 tttactttta ttaaagtata ggaatcaaac tggataccaa attctcagtg cagttgggta 60
 gtcattttgt taatgtattt ttaaaaaatt ttaagggtaa aaaccagcaa gattccattt 120
 agaatgattg tgaaaaaaaa actgtaagac gtccattttc aaaatgcaaa aatgattct 180
 tcctgatggt aggaaggcca atgaaaacta tatgtatatt gaaaatattt tttcctcaaa 240
 actttttccc tgatacagaa gtctgagagc ttactttggc tacattacct gactaaagag 300
 agaacttttag attagacctg gggtaaattg agatgccaaag ggagtgtcta gctaaatgga 360
 aataccacga aggtttgtaa tgccaagaaa gtcagctctg tgggtgtgtca taagcagcat 420
 atggaaacca ggagtgacac attagaacct gggagtgtg catacatctg atcaagcatt 480
 tgactctgaa aatattcagg gagtttagaa attgttaacc tttggaacca gtattgttta 540
 gcaatagttg agaagtgta gcaagaatga tatcaagtta aacttaggca cttggagtta 600
 catccttaaa gccttaatag ggcttatgag ttttatacag tcatacagat agaaatatgt 660
 tgcttttggt actacgacag tcatatatta taagaaataa tcaaagggtg gtggaaaggc 720
 atcctctctt tgatccaatt ttctgtacct ttttctcag gtcacacaca ctgctagccc 780
 aggaatcact aggtattgat gactctactt caagctgtgc aaagcccttt ctggagacag 840
 ccaggatggt ttgtagggag agaggcagga gtccctcagg agtggcctgg ggtgagaccc 900
 tcccataggc tctaagagtc tcattctcat cagtgtatc cagactgagg aagaaatggg 960
 gcagcagtca ggagagctcg ggatttatga gtatagcaga atttaagtga aatggaaact 1020
 acactcttta atttgttctt ccatggaatt gctttttcta tgcaagggtg gagccccag 1080
 gagagccctt gtagaaggga atgctgattt gtgtgaatat ctgtagggtga gtaggtatct 1140
 agtgaggatg agttgggagg atgagttggg taaggcgtgc ccctctgaca ctgattctgg 1200
 gataataaaa gacaacatca tgatgagatc ttcacatga aataaaaacta tgccctggcc 1260
 ttttcagaaa ctgccccac tgacaggtccc acagtgtgat ggagtccaag ctgggatcac 1320
 tgcgagatga ggagtccag gagtggcttc ggcaggcatg ggagcttcag gccctgagag 1380
 agaagacaga aattcagaaa acggagtgga aaagaaaaac gtgaagggaac tgcatgaaga 1440
 gcacatggct gagaagaaag agctacagga ggagaaccag aggtccagg gcctccctgt 1500
 ctcaggatca gaagaaggca ggctgcccag tcccagtgcc agatcagcac cctccgtgcc 1560

cagctgcagg aacaagctag gatcattgcc tcccaggagg agatgatcca gtccttgtct 1620
 ctcaggaagg tggaagggat ccacaagggtg ccaaaggctg tggacacaga ggaggactct 1680
 ccagaggaag agatggagga ctcccaggat gaacagcaca aggtgctggc agctctgagg 1740
 cgtaaccca ctttgtgtaa gcacttcaga ccaatcctgg aggacaccct ggaagagaag 1800
 ctcgaaagca tggggataag gaaggatgca aagggatct cgattcagac tctcagacac 1860
 ctggaatccc tgctgagagt ccagcgggag cagaaggccc ggaagtttct tgaatttctg 1920
 agtctgaggg gaaagcttgt caaggaagtc accagcagag cgaaggagag acaggagaat 1980
 ggcgctgtgg tgtcccag 1998

<210> 17
 <211> 653
 <212> DNA
 <213> Homo sapien

<400> 17
 gcgtggtcgc cggcgaggta catggccgca agcagactaa cgcgttgacg ctaatttaat 60
 gtattttacc tcacactaag gtcattgcttg ataaagacgt taaactcaac ttgtaaaatg 120
 gtagcccagt gctatgcaca gagtgggtgc tcattagtgt tgaatgaaca catttgtaat 180
 actacatgta attccatctg actgctttgt taaattttca gttagaacgt agatactgta 240
 aagtccacac acacattaaa tcttgttttc ctgaaagtat ggcatacaaaa atactttag 300
 aaaaaccttg tcacaactga tttgaaatgtt cctattttct ttgactttga tattggcttg 360
 taatgtctct tttcatcata tgtaatatca gtggaacagg cagcgtact caagtcctaa 420
 ggattcctca gtgatcagtg atccagggcc gttcatgaac cactgggctg gatttgactg 480
 ttgagtgtgg cagttaatgc ccctcaagaa atcaaaggat gtcttataag tgtcttccaa 540
 aaaaaagcaa atgctgaaat cctattggca aagtaaactg aaattggctg ctatatttta 600
 tataatcatt tctgcaaate ccattttttg aataactaata tttgacatgg tta 653

<210> 18
 <211> 1498
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (29)..(29)
 <223> a, c, g or t

<400> 18

16

ttattcagtg catagcttta agccagtnt ggattcacta agtggacagc cagtctccca 60
 gctctctgcc ttccccaaa gggtcgtagt aggtcacctt tctacagcag ctaactagag 120
 tcctaactaa tgggatccag cagggccatt tctccagagg gccagtatcc tattaggaga 180
 ctcttggaaat tcttaggttc tactcaagag tggaaggacc aatcacctct gatattctgt 240
 ggaaggtttg gggtaaatt ctgccctctg cattctgtgc aacttgata aaagtcaagt 300
 tagtattaca tgaattgggg tagggttagt gctttgaaa aatggtgaac cggctgggcg 360
 cggtggtca cgtctgtaat cccagcactt tgggaggccg aggcgggtgg atcatgaggt 420
 caggagtctg agaccagcct ggccaacata tagttgcttt ggacctcatt tggaaaaata 480
 atctgccttt ctaattgttc tgcataaggtt aaaatgataa atttacattc tttgaacctc 540
 taccagattg tgggtgtccga gtgaccggca cactgtctga cacacagtca gtgtgcacgt 600
 atttgtctga gtgaatgagg agacctgaga aaccgggtgac gtggcacagg gaagccagct 660
 ggcccaggat tccgtacatg gccgcaagca gactaacgcg ttgacgctaa tttaatgtat 720
 tttacctcac actaaggta tgcttgataa agacgttaa ctcaacttgt aaaatggtag 780
 cccagtgcta tgccaggagt ggggtgctcat tagtggtgaa tgaacacatt tgtaatacta 840
 catgtaattc catctgactg ctttgttaaa ttttcagtta gaacgtagat actgtaaagt 900
 ccacacacac attaaatctt gttttcctga aagtatggca tcaaaaatac ttgtagaaaa 960
 accttgtcac aactgatttg aatgttccca ttttcttga cttagatatt ggcttgaat 1020
 gtctcttttc atcatatgta atatcagtgg aacaggcagc gctactcaag tcctaaggat 1080
 tcctcagtga tcagtgatcc agggccggtc atgaaccact gggctggatt tgactgttga 1140
 gtgtggcagt taatgccct caagaaatca aaggatgtct tataagtgtc ttccaaaaa 1200
 aagcaaatgc tgaatccta ttggcaaagt aaactgaaat tggctgctat attttatata 1260
 atcatttctg caaatccat tttttgaata ctaatatttg acatggtaa ttcttattaa 1320
 tttgttgaa ttgtttattg ttaataatgc aaatagataa tttttaatta tccacaactg 1380
 atttgaatgt tcctattttc tttgactttg atattggctt gtaatgtctc ttttcatcat 1440
 atgtaatatc agtggaacag gcagcgtac tcaagtccta aggattcctc agtgatca 1498

<210> 19
 <211> 171
 <212> DNA
 <213> Homo sapien

<400> 19
 gccgccggg caggactaa atgaaacata atatttattt ataaaagtgt gtagattggt 60

17

aatcacaaa aagagtgcta tgaccattat gtatgaggaa acaggccttt gacctcctgg 120
 aaagcactgc tcaaaagtca ttagtgccca ttttgaatt ccccaaacag a 171

<210> 20
 <211> 1820
 <212> DNA
 <213> Homo sapien

<400> 20
 gaatttcgta atccttgaaa ttgaaaaaaaa aaaaattgtg tttttaaga gtgaaaacag 60
 ttaggaaaca agtagaactg taatcagaac gctgcttcaa ttgatattaa aaataacctc 120
 aataataatg taaaggttcc tttctcttgt gtcagttata ttcttaggga tagcctagaa 180
 ggaatatatg gttagaacta agtgtgacta atcatctgag ccttgaagag aaacttcagt 240
 gcctctaaac agatcatcta caaaacaaca ggtaaacatt tatgccagtt aagtgggtca 300
 tgtttttggt tcttggggtt ttcctaaatt taagtgaggt tgggcttacc ttgtagataa 360
 aattatggtt tctttttggt aaatacttga atgtggataa cgtcaaatca gaatattttg 420
 tgaggaggtg atgatttgaa attaagctag atttctaggg aggtgttggg tccaatgaag 480
 gatgggaaga aattaaata gtcttcaaac tcttctctta ttatatttgg ttgctttgga 540
 aaagattggt cctatcctca atctaattta ttcactatta atattttaaa aacattcctg 600
 agatacttaa aaagaccac ttagcgatta tagttgctca atgaaacaag aatttattta 660
 tgcatagatt tttctctgta tcttaccaaa atccacttta cttagataac actaaattgt 720
 tcttaagac tactcatttc ccaataatcc tttatgattt caaaatttct agtggctcag 780
 aagtgaattt tattttattt gtctttcact tgaataaatg agaaccaga aattaataat 840
 gttgtttatt gcttactgtc aggactattht caaagactaa gaagagtttc ttctaaccoc 900
 tccctctcaa aggaatccta aattattagt tgtagataa gttttgtatg ctaagatatt 960
 caggtttata gtttatgtat gtgtgtatat atataaatat atatgtatat ataaatatta 1020
 tgttcagttt ggagtctggc acaactccat tatgtggatt agagagtaag atattatgga 1080
 tgataaagta ctaaataaaa cataatattt atttataaaa gtgtgtagat tgttaaatca 1140
 caaaaagagt gctatgacca ttatgtatga ggaaacaggc ctttgacctc ctggaaagca 1200
 ctgctcaaaa gtcattagtg cccatttttg aattcccaa acagaaagct tcttagaaaa 1260
 cacgctgaga ttttatttac aggaattct ttgacacatt tcaattggtg tgtagtcaag 1320
 tatagcaagt acttaataat gactgaattt catgttccta cagtcataca tattcattag 1380
 aagttttatg ttgttggctc gatctgattc ttctttggtt gtgggtggaa cggcactgag 1440

18

agaagtatag ttttttaaac ttgaacatgt tcagtagtta cattgcctta gaaaaccag 1500
 acacatagca gtggaaatga aagaaatggc atcagaagtg acttaattta gcaattgtga 1560
 ttcctcttgt aaaacaaaac aaaaaaaca tgccatattt tttggagaaa agttggcaat 1620
 ataggggttt cgttgtctgt ttcacaagaa gactcatttg ttcttttggg ggaaccagtg 1680
 ccttacagat tttgtatata ctgtaattat tcaggactag ggaacaaaca attgtattgt 1740
 atttgttaca gattgtatat ggctttgttt taacattccc ctaaataaaa tggcttcatt 1800
 ctccccttgg aaaaaaaaaa 1820

<210> 21
 <211> 611
 <212> DNA
 <213> Homo sapien

<400> 21
 acccaagaca ggttctgaac catgcttatg cagagctttt agtattaaag agggagagta 60
 aaagaagtgt cagagtccag atttatcact gaaccaata ctttcttact ccctggggca 120
 tctcctaata ctgatcctaa aatgctcctg tttctgagaa gctagggcaa gacctgcctt 180
 acaaagacca gccatttgcc ttattcatag gatcataagc aagagaactg cattccagga 240
 agaatgaagg aagaaggaag gctgctcaca gtagcagaag ggaggcaggg gccgagctgt 300
 tcaagtcaca taaactctaa gaagcccagt cagcaaaata agtctatctt caattctagt 360
 tgagtcagg actctgagga gctgtgatc acccagtttt tcctgcaaaa gccacagctg 420
 ctaaaactaaa ttggtgcaat tcacttctc ttgcctctct gggtcattcc accaattgtg 480
 gttgagaaac acatcttagg gaagaaacag tatctaagca ttaaagagaa aatatccac 540
 tttgctctc ttctcccta aaccggaact gctcttacct acaagataat ttttaaatca 600
 taagattggt a 611

<210> 22
 <211> 1885
 <212> DNA
 <213> Homo sapien

<400> 22
 catgaacatt tgaggctgat tccctgtggg aaaaatcatt caaatctatt cactcatctg 60
 atggctgttg cttgttttat tttttgtcca agagaggtgg tgttggaccg aggtagagaa 120
 gacagtggta caccagaaat aacccaaagg attgcccctt ctgtagaagg cccttagact 180
 ccatgatgcc tttcagctgg gtgctatact tgcacctaac tctgggggct tcactttcta 240
 tccctacaat tactcaaaca gataaaaggc tggatgtaa catgtagtta taagggcgct 300

gatctaatag taaggaatat cacttccac aagtccttca aacaagattt gtgaggagct 360
 ggatttgtca gcatgtcaga tctttttgaa aaccagagag tagaatgtaa gcaataccct 420
 tgtcgttaatt aaagaccaga ctccatcctt ataccactga tgcctctggg accttaatcc 480
 ttaaaatatt tagtgaccct tgccttctaa ttcttgacac aaatatataa tgaccatttt 540
 agatcgggga actcccttct tttgaaggca gtttagggat tccacagatg ggctttgaac 600
 ctgctaaatg tgtatggaaa actgagttaa ttacaaatgt ctttttctca aaagtgcgtt 660
 tctggtttct gtcagattca acaggtctgt acccaagaca ggttctgaac cactgcttat 720
 gcagagcttt tagtattaaa gagggagagt aaaagaagtg tcagagtcca gatttatcac 780
 tgaaccaat actttcttac tccctggggc atctcctaact actgatccta aaatgctcct 840
 gtttctgaga agctagggca agacctgcct tacaagact agccattttg ccttattcat 900
 aggatcataa gcaagagaac tgcattccag gaagaatgaa ggaagaagga aggctgctca 960
 cagtagcaga agggaggcag gggccaagct gttcaagtca cataaactct aagaagccca 1020
 gtcagcaaaa taagtctatc ttcaattcta gttgagtcca ggactctgag gagctgtgat 1080
 tcaccagtt tttcctgcaa aaggcacagt cgctaaacta aattggtgca attcacttcc 1140
 tcttgcctct ctggttcatt ccaccaatg tgggtgagaa acacatctta ggaagaagaa 1200
 agtatctaag cattaagag aaaatatccc actttgctcc tcttctccc taaccocgaa 1260
 ctgctcttac atacaagata atttttaaata tataagattg gtattaacac aattattgat 1320
 aaagagaaac aatgaccaac tcattagcta acgatgctag aatacttatg caagccctag 1380
 agttaagggt cttagtgtgg acaccttcc agaattggaa ggaaaacca ccagaaagct 1440
 tattaccctg catcagctga aaagctaagc cacagccatt tccctaag ttctgtttct 1500
 gggagaatga gatcttcaag aataactctt gcccttgat gaggcagtca aattcaaacc 1560
 agtgatggca acaacttgca aacacgtaat tcttgcctta atttccagc acttaaaaca 1620
 aatccccac tcaatacaaa gtttctatgt gcctcttgcc tgaaatcaac aagaacagc 1680
 tcacctgcc aaagactcct ctttctctgc cagggcaaaa gcaatctgca gccagagat 1740
 tcaaactag acatacacat ccacaattgt cttaatctca gcagtactgg gaaagctttg 1800
 tactcaactt aacctgtcat ttaaccctt ccactagttc tcccttaacc agactgcttc 1860
 ctgtcttgaa acaaagaaaa aacc 1885

<210> 23
 <211> 494
 <212> DNA

<213> Homo sapien

<400> 23

```

aagcgcgcgc attgtgatgg atctatattt taccctgtgc ttttctatag ctgtcctcaa      60
agcgtaaacc attccaaatt atttccaac gtagtgttat atgtgtgcag cagagctatt      120
tctgcctggg cattgccagt ccctgagcag gagggtctca cagtgaggtc tgcaggactg      180
taagtttggg gtctgactcc ctggccacc tgtgtgggct gtgactgtct ctcagagcta      240
taccgcctct ttctctgctg gcagcccagc agagctggct caaccatcgg aggtcgcagg      300
ccaccagcca cgtggcacca ccatggcagc cttccagggt aaggtgagac acacaaggca      360
tgacctgggg gccgaccgga tccccatcac aaacgccaca aacaccataa acacaacca      420
ccctgatcag agactaagca gagaaagcag ggagaggacc tagagttact cagtaatgac      480
tcaggaagga gacc                                     494

```

<210> 24

<211> 1692

<212> DNA

<213> Homo sapien

<400> 24

```

gtccccacc atggaagagg cggggccac ccaactgcaag tcttctctga gccacgttct      60
caagtcttct ctgagccgcg ttctccagg tgtgtctctg gagtcagttg gcatttcctc      120
caagcctgaa agtgtagtca gattcagaat gggcttttct agattcccct gtaagatctt      180
tcccctgctc ctggcaggag caccacacca tgggaacccc agggcccacg cagctgcccg      240
ggactggggg accaggacgt ggcacttctc acatgggtgg aaagatgggt ttacagaatg      300
gtggcatgga gacgctgtgg cctggcaagg atcaatgggg tggcatctgg cattagccat      360
caggaagact taaggctgaa gggacattgg gcagggagct ctcagggctg ctccaccgac      420
cccagggtg acagcccata gtatcactta ggggtgggact gagagtcacc tgggggagag      480
gagagaaggg gcccaacttc cccagcccct agtatcactt aggggtgggac tgagagtcac      540
ctgggggaga ggagagaagg gacccaactt cccagcccc tggcaccttc cctgcctttc      600
ccagtctttt accagagtca taagatggtc cttggctctg ggcaggcatg tggccctggg      660
gagctctggg gtcagaggtc aagggtcttt gcatgtcagg caggcttgac ttttgctgtg      720
agaaagacta tagaaagatg gcaagctagg cctcttttct ggaaaagtgc caacagctga      780
taattttagg aaataatgtt ttgaatgtga agtgtgactt tttagaataa aaagacagga      840
agctcttaga aactgcaaga ttctaaatct aagcaaaagg ctatatatta ccctgtgctt      900
ttctatagct gtcctcaaag cgtaaaccat tccaaattat tttcaactag tgttatatgt      960

```

gttcagcaga gctatctctg cctgggcatt gccagtcctt gagcaggagg gtctcacagt 1020
 gaggtctgca ggactgtaag tttggggctt gactccctgg ccaccctgtg tgggctgtga 1080
 ctgtctctca gagctatacc cgccctttct ctgtggcag cccgacagag ctggctcaac 1140
 catcggaggt cgcaggccac cagcccgtgg cccacctggc agccttcag gtgaagggtga 1200
 gacaaacaag gcatgacctg ggggccgccc ggctcccat cacaaacgcc acaaacacca 1260
 caaacacaac ccaccctgat cagagactaa gcagagaaag cagggagagg acctagagtt 1320
 actcagtaat gactcaggaa ggagacccta agcttctacc acatgccaga ctctgtgccc 1380
 agtgcagcat aaacgtcctc agaaccagcc tggctccagc ctggccgagc cggacgttcc 1440
 tgggaaaggt tacaggagga gcagggccag gccacagca cttttagaag cccatgaaaa 1500
 tgtcttcatt tctcttcaaa tcacaaacaa aacgtgcaaa acccattctg gagtgcattc 1560
 tttcactggc gaccaacca gtccaaagat aaccttctta atagttctat ggaggaagct 1620
 gaaaggcag aagtgactac aaccacaaa agtcatgatg gagccctgac gtgtgtgtac 1680
 acacacacta ca 1692

<210> 25
 <211> 430
 <212> DNA
 <213> Homo sapien

<400> 25
 acccagcgtc ccctggccag agccaccaga ggacagagct cccaatgagc ccagctgcta 60
 gaaaagaagg tggagtccca ggcagaagag ttcttcaggc tgaacggaaa tgattccaga 120
 gggaaatgca gatatgaaga aggagataaa gagctccaga aatggcaaat agcaggggtga 180
 gcctacgcga cttctctaac ggaagaaatt accttataaa cacacgtgca ggcttagagc 240
 aaaagaaacc gtgccataag gtgtgagtaa gtgaagtgcc tgtgacacct acagatcaga 300
 gaagcagagg cctccgggat ggcaaggcaa ggttgccgca tttcatatga agtgcacaat 360
 catcataaaa gaatgcatta aatatacata tgtatgcatt caaattacac taacatcaca 420
 tatatccatt 430

<210> 26
 <211> 2603
 <212> DNA
 <213> Homo sapien

<400> 26
 tgtttggtcc agtgaatctg cccaccaaca ccccgctct caccatccac cagcccttgg 60

acccctagca ctgagctcac agtgaaaggg aatatttgct tgtaaataga aatagacgct 120
 ggtagaaac caactggaaa gaatctttcg ttgaatagga gttaaaaaac aaggaaatta 180
 acccactcct gggatattct gaaactggca atctatgctt gtctaggacg gccagacta 240
 accctaactg ccccgctcgc aacacagcag cacagcggtt cctccaggag aaacaccaag 300
 atctcacgtc ccatccacag gctgaggctg ctgctcctgc aggaacctgg tgcagtgtag 360
 caattccaca tcctgaaatt gctcatcaaa actcctatta aagtgtcaaa cagtgaatag 420
 ctaaaatacc actttgcttg aacagtgaag aggttgggaag gaaaacgta actgtatcag 480
 agaatatgga ctccatacat acagggagtc aggttcattt tgaagtcaact cttcttccaa 540
 cagattcaact aaggctcttt gtcaacacaaa attgaaaacc gttaaaaaaa aaagtaatta 600
 tgatgcttcc tgcctccat gaaaggacca catacagaca ccacgctcat atctgaggcc 660
 ctggggtagc ctttaatggc ccagcagaat ggccagaacc gttagaggaa acatttaata 720
 aagtctggag tcagagccct gcgggtctag ctggattcct ggaggtgctg cccagaagcc 780
 agcgggaggg aattggaggc cggaggctca aactgtcccc acttccacca agggccccctc 840
 ctccaacagc ttccaggctg ccaaagcccc tgcatacct ccagggtccc ctgggtccag 900
 cctcatgctt ccataatga gtttttaaac cacaacgctg catcagggtga catctcttct 960
 gcaggctgtg cgcgtctcca gggggaaggg ggctgtgtct ttgggacagt ttgtgctctc 1020
 aatcacttga ctgctgacag gcacctcagc tgaatgggtg gatattatgca aagattgtgc 1080
 tgaattattt aaagcattct ctatttaaag aacagagaat atttaattag cattctgctg 1140
 tgcttaattg aagactcaca aatcaattaa aactgcttac cttttggcag ttcagtaact 1200
 tcacagaaac ctcccaggaa atgcataccta ttcacagctg ggttcatcct ataccagcgc 1260
 acctgtggcc agtgtggcgc tgtgattaga ggcggtcag cgccttcaga ggagcggcct 1320
 ggctgtgctc acattagaga aaggcttcca tcgtcgttgg tcctctttct cacagggact 1380
 ctggggtctt ggtgccggga gatgcaaccg cctctggcag cccggcttca ttttagggac 1440
 agtgactatg gagaaacca ggtctgacct atttctcca gaggggaggg agccacaggg 1500
 aaaggcccct tgttctctg ttggccctga gtgcctcca ggaaaggtca gagcacagac 1560
 tcagccctgg gagggccgag agatcccgcg ggacctgcc ctctcgaca ctctggacaa 1620
 gatgcagaga gtgggtcctt ggcagcaaga tcccgaggga gtggggcctt ggagctcagg 1680
 gccagaccga ggggtgctc attgctggct ctggctaca gacacgttga cattggcacc 1740
 acacgggcca actgaaacct taagagaaaa cccagcgtcc cctggccaga gccaccagag 1800
 gacagagctc ccaatgagcc cagctgctag aaaagaaggt ggagtcccag gcagaagagt 1860

tcttcaggct gaatggaaat gattccagag ggaaatgcag atatgaagaa ggagataaag 1920
 agctccagaa atggcaaata gcagggtgag cctacgcgac ttctctaacg gaagaaatta 1980
 cctttaaaac acacgtgcag gcttagagca aaagaaaccg tgccataagg tgtgagtaag 2040
 tgaagtgcct gtgacacca cagatcagag aagcagaggc ctccgggatg gcaaggcaag 2100
 gttgccgcat ttcatatgaa gtgcacaatc atcataaaag aatgcattaa atatacatat 2160
 gtagtcattc aaattacact aacatcacat atatccatta gactttatca aaattaaaaat 2220
 cttctgttca tccacataaa acgatgtcac ttactgcaa aaatattctc aaatatttat 2280
 ccaagtgctg agatccagaa taagtaacc ctaaaatttc ataataaaac aacttggtga 2340
 aacaacggtc aaaggatttg aacacttgcg caaatgatgg caaataaaca caagaaaaag 2400
 tgctcgacag actcgagcac caggaagatg cgctgtaaac accaacaaaa accaccacac 2460
 acaccacag tagccaaaat ctataaaact ggtggcacca aacgtgaggg aggatgtggc 2520
 ccaccagca ctggtgctgt gcattcttgg tgagaacacc taagacgtcc cctcaatggg 2580
 attagaaaac cacaaggcag gca 2603

<210> 27
 <211> 614
 <212> DNA
 <213> Homo sapien

<400> 27
 acatatat t aaaggaaga tggatacaat ttgtttttat tatataaatc taggtaaggt 60
 gaaatgcttt tgtcaacaaa aatacagtg agtgaat ttt atatttgctg cttgattagg 120
 taaactgaaa actaacaata gaaatattat ttactgcat tgaatacca tgaactttca 180
 gacttgtag ttctacaagc agttgtgcta ccttaat ttt gtgtttccag aaataaaaat 240
 taaccttagt tatgctgtca tttttaacta ataaaaaaag tataattcat aaaacttttg 300
 gctttataag ataattataa aattatata tttttctgt tttgtggg ttgggaaaac 360
 attttcttat ttctattcac tcttcaaag caggtctcat aatattgtgc aatgatataa 420
 gatgatggaa gacttctgta ataaaaacat atgtcattat cttcaatttg ttcaataaat 480
 aatttaactg tgaacaaca aaaaaaaaa ccaaaaaaaaa aaaaaaaaa acaaaaaaacg 540
 ggggggggccc accggggcaa agggggcccc gggggaaggg ttccgggca aatcccata 600
 agagcaaaaa acat 614

<210> 28
 <211> 1134

24

<212> DNA
 <213> Homo sapien

 <400> 28
 gcacgaggat tgggtcaaagt agtattctct tgaagttcta gtcaatttaa tttgatccaa 60
 taagtttttc tgaatctcct ttttaagttc caagaaattc tattataaat aagtgtactt 120
 ttaccaattc cattgtataa gcaaacagac accttttaga aaaggataag taatcatcaa 180
 tttgtttttt ttaaaaaaaaa acaatttcca gactactaaa tttggcataa gaataattct 240
 tttaaaatgc aacatacttt aattagtttt tttggtatat gcataagatg tgaactttcc 300
 tattgatatc actttatatt aatagagatg tacatttctt tctatgccgt ggctagagca 360
 aaagttaata atgattatct acacaattga ttaatttct taggatatgt ataataattgg 420
 atattatatc tgatttaaaa atactattcc atacattttt tttttcagga gataaaacat 480
 agggaaaggt tttcatgtga attctttgta tcaacttgaa gtacatatat ttaaagggaa 540
 gatggataca atttgttttt attatataaa tctaggttaag gtgaaatgct tttgtcaaca 600
 aaaatacagt gtagtgaatt ttatatttgt cacttgatta ggtaaaactga aactaacia 660
 tagaaatatt attttactgc attgaaatac catgaacttt cagacttggt agttctacaa 720
 gcagttgtgc taccttaatt ttgtgtttcc agaataaaa attaacctta gttatgctgt 780
 catttttaac taataaaaaa agtataattc ataaaacttt tggctttata agataattat 840
 aaaattatat atttttttct gtttttgggg ggttgggaaa acattttctt atttctattc 900
 actcttcaaa tgcaggtctc ataataatgt tcaatgatat aagatgatgg aagactttgt 960
 aataaaaaaca tatgtcatta tcttcaattt gttcaataaa taatttaatg tgaaaaaaaa 1020
 aaaaaaaaaa ccaaaaaaaaa aaaaaaaaaa acaaaaaaacg ggggggggccc accgggggcaa 1080
 agggggcccc gggggaaggg ttcccgggca aatccccata agagcaaaaa acat 1134

<210> 29
 <211> 1139
 <212> DNA
 <213> Homo sapien

 <400> 29
 cgaggtaccc attataatta ctaaactgtg aagtcactat tattagtatc tgaccagcta 60
 taaaaaacat catcaatttt acttttgaca caaaaggtag taaaaatcgc aaacgataaa 120
 gaagacacta ctcatataaa gtcattgtta ctaatccagc accataattc cagtctcaga 180
 acctcccatg cagattggaa agggattatg ggaacgaggt gagtatgtag gacatgtcgg 240
 cgctagtaac atcaaattga cggccccata tttgctcgtc tcacaagaca aaaaacacag 300

25

ggtcctccca aagtaagcag aagatgacat gacggcatgg agacgaaaa caaaacgcta 360
 gcgcgctaaa tcaatggtca atagctgcaa aaccatctga tgacaactag ggtaacttcc 420
 cgtgtcaacc aaaaattcac aaacaagtaa gcactacctg tagaacagac acgaagtcac 480
 gcaaacctac actttgagca cgcctgacca gagatccgag cacactcccc gaccaccaa 540
 cacacagcag gccacgcggt agagagaaca agaatacaaa ggacaagcga gtagctgtag 600
 aagcgatgag agagagcgta cgtagagatg ggggaggaac accacgtagg agcagaactg 660
 ctgcactgcg tgcacacgcg acgcgaacag acgaaactac acgaagacaa aaggaaaagg 720
 aaaggatggg accagagggg agagccaagc atgagagaca caccaaaagg caccgcacg 780
 ctgcatggcg aagcgagaag aacagcagat aaccacaaaa aaaagcacac acggtgggac 840
 atacacacca gagggggagc atcagacaca gggacaaacc actaaagcag gagaacatgg 900
 cgcgaaagga ctgaactaaa cagcacaaac acgcaacgag cagcgaacag ccgatcatag 960
 gcgtgacacc cgactacagc aaaagaaacg gagaagttat cgacacaagg gatgacaagg 1020
 aaacaggcta atggcccaag gagaggaaca ataagatgga tgagcacagt agggcgaaca 1080
 agggataacc caagtgaaga aacagtgaag aagaggaatg cacacaataa gaacgcaaa 1139

<210> 30
 <211> 235
 <212> DNA
 <213> Homo sapien

<400> 30
 agtgtttgca acagcaccat ttgtcaaatt caaagatgct caaaaggtgt tccctacttt 60
 gcatgagagg gagagctttg taacaggaaa ttgtataagg caaactctct attcattcct 120
 aaggcctctg ttcattccta atgtttacat ggttctctac tctgaagggc accaacatgg 180
 acctcacctt cttaacatgg aaaatcaaaa tctaaatgaa ttaccattaa aagga 235

<210> 31
 <211> 2171
 <212> DNA
 <213> Homo sapien

<400> 31
 ctgcattttt ctgtcattct cttcatttgt ttaaggtgg aaaattttct tacagttgat 60
 gcaaagtatc aackacttta ccctaccttc tcccctttta gatgggttct tctgagttt 120
 tggagtcttg tatgattatc agtattcccc tgtcaaaatc aaatctatc aggtttcttc 180
 actgttgaga acacctaagt gtttttattt ttgagaagtg gggacagagt ctcaactatgt 240
 caccaggct ggagtgcaat ggcattgatc cagctcactg caaccttcgc ctctgggkt 300

caagcgattc tcctgcctcc gcctcctgag tagctgggat tataggcacg caccaccaag 360
cccagytwwt tttttgtatt tagtagagac agagtttcac catgttggcc aggctggtct 420
tgaactcctg accttgat ccaccacct cggcctcccg agggtgctgg gattacaggc 480
atgagccacc acgcttggt aagaacacct aaattttat gtttcttggc tcaaaaacca 540
gttccatttc taatggtgtc ctcaagaag ggctaattgg tggtagagaca gcaggggagg 600
aggaagagct gtggtttga acttgttcaa ctgaggcaat aagcgatttt agctttat 660
aaagtcttct gtccagcttt aagcacttg taagacatgg ctgaaagtag cttttctatc 720
agaattgcag atagtcatgt tgggctaaca gtcaattgga tatattcctt tacctcacat 780
gacccagca actgtggtgg tatctagagg tgaacaggc aagtgaatg gacacctctg 840
ctgtgaatgt tttagagaag gaaattcaaa aaatggttga actgaaagca ctggtgaata 900
tgggtatcgg ctttctttt cactttgact cttaacatta tcagtcaact tccacattaa 960
tgaaagtga ccatagttat ttccaataa aaagaaacca actcttacca ggtcttggac 1020
tgtgatgtca tattattcag ttttatgctt gttcctgagc agaactcata agagtgacat 1080
agtcagctgc tgacggcacc tcagccacgc cactcttact cagttcagtg ggtgtgcttg 1140
cgtgtagga tgtggtgcag ccctctctac gctcttctat ttttggata tttcctatct 1200
aacctcaaa tagcttcaa ttctttttt cttggactgg cttcattctg aatttgtgct 1260
aaaataatct ttcataaaga gacctcagtt tatagcgtaa cagactacac aatgcactga 1320
tgttttcata atgtttaagg gacccaactgc aagaagcttg ctgcctcctt ttaattgtat 1380
tcatttagat tttgattttc catgttaaga aggtgaggtc catgttggtg cccttcagag 1440
tagagaacca tgtaaacatt agaatgaac agaggcctta ggaatgaata gagagtgtgc 1500
cttatacaat ttctgttac aaagctctcc ctctcatgca aagtagggaa caccttttga 1560
gcatcttga atttgacaaa tgggtctggt gcaaacactt ttttttgag atgaagtctc 1620
gcggttgc caacgggctgg agtgcagtgg cgtgatctcg gctcactgca acttccacct 1680
cctgggttcc agccagttct cctgcctcag cctccaagt agctgagatt acaggcgct 1740
gccacccac ctggctgatt tttgtaatt tagtagagac ggggtttcac catgttggcc 1800
aggctgatta actcctgacc tcaggtgatc cacctttctc ggctccaaa gtgcttggga 1860
ttacgggtgt gagccaccgt gccggcctg caaacacatt ttaattgaca aacttagggc 1920
tgttgtacaa aatagtaatg atagccatgg aagtttacc ttattctgtg agaagtgtc 1980
ttaacttat taaagtgtct aaaactaagg ttagtgcttc taaaggaagt ggccggttct 2040

27

cctaagaagc aattatcact gtcctgact ttgtctggtt ggtttggttc ccctgtccc 2100
 cgattggctc tggtagcctg ctttgccgcg gttcttttaa gccagcgcgg gttatTTTT 2160
 gaaaacctcg g 2171

<210> 32
 <211> 192
 <212> DNA
 <213> Homo sapien

<400> 32
 gcgtggcgcg gccgaggtac tgtctctaca gccattgaga agccattcag tgcctggta 60
 gggacctgag actttccaga attcacacag cagtctatga tccctcaaat gtaagaggac 120
 agggggtcag cctatcttca cctctcagtg aatgtggagg gccaaagcaat atgacttgca 180
 aacctaagct ag 192

<210> 33
 <211> 2641
 <212> DNA
 <213> Homo sapien

<400> 33
 tttttttttt ttcttttcca agttatTTAA tttacagcat cagtctccaa atataataat 60
 attaagatag cagtttagaa attaactttt tttcagatca ctctaacata aaatctctca 120
 actgaatctc tagtttgtct ctttttgTTA agagctTTAA tattacatgg gaagttcaga 180
 gacttctatt tccatccctc aacatgtagt gacagTCAAC atgtcaggct ctgtagcacc 240
 gtgatatocc agcaccagac cactccagcc accctctcat tcaaagaagg gctacaagat 300
 atggctggac tactcgaatc acatctgatc ttaatcaatc caggtataga aagttgtact 360
 ataaagaata ctttccaaaa ttgttcactc aaataaaaaac agatcaagtc attacagagc 420
 atttttccat ttaataaga ataacagacc tactcaaggt aatTTTattc tgtttattta 480
 aataaggata agactactta aaagactttt tacatacaaa aatgtacaag gTtaaacttt 540
 tctgtactga attacaaaac ctgcacaagc atgtaataaa agagcacact taaaaacatt 600
 ctgaccatta tttagcctct aaaaattact gaagttcaac agtagtaaat agaggaagct 660
 cttacatata tatatatata tatatatata tatatgattt aatctactgg cagttttact 720
 taatgtaagt atttaaaagg tcacattgct attgaatgag tctctagatc aatTTtagaa 780
 ttgtctctca aaacttaagt caacccaaaat attatttcaa atagtaattc caattctgaa 840
 gaattttaat accagcaaat atattatggc ctcatagtag taactgaacc aactttccaa 900
 agtgcctggT agctgtccag atgaattagg ctgctttgga aaactgtact gtctctacag 960

ccattgagaa gccattcagt gccctggtag ggacctgaga ctttccagaa ttcacacagc 1020
 agtctatgat ccctcaaatg taagaggaca gggggtcagc ctatcttcac ctctcagtga 1080
 atgtggaggg ccaagcaata tgacttgcaa acctaagcta gaagcttggg atctacagta 1140
 aggaggaagg agaattaaag tagagaaaga aaatgtataa ggagaaaggg aaaagaagga 1200
 acaaagaggg aaaagaagaa aaaacaagga tgcctgctaa tggcaggaag tggtaaagtg 1260
 cctataacta caacttacia gccaccact aattctaata ccattcattt gcctactcca 1320
 ataataagaa aagctggctt tactggaata tagaatctag agcaacatta cccgcctcat 1380
 gttagtgagt aactagtatt ctaaagttgt ttgccataca tatcaagttc ttctaacctt 1440
 tgaagcaaac caaaacactt caaaaactcag ggctcccagg gctgctgctc cagattccca 1500
 gcattcagca tgcttcatta tgtggagaaa gacatttcaa gacaagctgt atctatacac 1560
 cttcagaagg aacaaagctc taagaagggtg ggattatggt aacacatagt acatggttta 1620
 gcgtttctcc acatttcaaa ctcaaatag ctcaataata tgctgctaca tgagcattga 1680
 ttctgaatgt tcataatata aacttcaatt tgaagcaaca atggtacaca gttcagctgt 1740
 tattaccaac ctactctgta agttaaata caaataaaat attaatttta ttgagtaact 1800
 aaaaataagt tcccactgac ttaaaatcgt caaatggcta actctctctc aactaagaga 1860
 gcaacacaga tggagcaga gaggacaact gaatataaaa taaaatttgt caatctactc 1920
 tataatctgc acttttataaa tccccttttg catatatgta tgtataggat cacagttgcc 1980
 caccaacatt atgtctgtca gccctgcaga taacaattta ctgtaacgtt aacaatttat 2040
 gcaactacta gtatgtttta tcttatgtgt acagatttac agtttggat aaaggcagaa 2100
 tgattaaaaa ctattgggtt aaagtcttag tatggtaact acctgcaagg ctgaattaat 2160
 tttttggaag gctattcaat agctgaacta aaatgcttgt ttaacaaatc aaaagaggaa 2220
 taagactact ttaaacata ttgaaaaagg taaatcccaa tttgaagatc aatcatataa 2280
 cgaaaaaagt atgaagtatc ctttgctctt gcttagaac acatagcaga acagtagaaa 2340
 ctagaactca tgaatataag gtaaacctta ttttccact gatttccatt atacaattgg 2400
 agtgaaaata ccaactcaaac aaaaataaac aaaaaatctt agcaggtaat tctgtgtaga 2460
 acagccatgt ggggaattgtc tatattacag ctgcaggga tctcatgtaa gctaggagtc 2520
 catcttccta tgttgactc tgcaagtact tctgactccc agtagctcct ctattgccta 2580
 ctccatatta cgctaatttt tgccccctga ctgctatgct tcctgggact cttattaat 2640
 t 2641

<210> 34
 <211> 434
 <212> DNA
 <213> Homo sapien

<400> 34
 atttccttat acacacaccg aatcagaata tactttcagt tctacaattt gacaatacac 60
 atagctgatt tatagcaagt gtgccatgaa ctgagggttt gtttagttg tttttgcagg 120
 gctgccaata tgctgtcttc acgggacggt aaagaaagta tcacttgggc cgcactctaat 180
 atgaaatact gaagggtgggt gtagagaggg tgctagggct ttgaacagcg gcacttcctt 240
 tctgagagag agaaaacatc atgctcccc cgcgccgaac tcattttaca ggttgattgg 300
 gtgaacaatt cttggcaggc cctgagctag tctgggtatc ctgagtcaag agagaggccc 360
 tgcctctgag gtaaagtgtc tctcatctgc ctaagtttgc ttagaaactt tggcttatga 420
 aagattaacc taag 434

<210> 35
 <211> 197
 <212> DNA
 <213> Homo sapien

<400> 35
 tctgagacaa tagggcatgg gtcctctaata tcatctcgag cggcgcagt gatggatagc 60
 ggcgcccggg cagggaaacc cctactggac cctgtgtgtc tgccagcctg ggcctttgt 120
 ctccagccct gcctttatc ctccttgct ccacaccagc ctccccttgc ttctccttac 180
 agactatcca agaagtg 197

<210> 36
 <211> 3414
 <212> DNA
 <213> Homo sapien

<400> 36
 atgggggatt tgcagcccc cgctgctgcc gcgaatggca gtagtatttg catcaacagt 60
 agcctgaaca gcagcctcgg cggggccggg atcgggtgga ataatactcc caatagtact 120
 cccgctgctc cgagtagcaa tcaccggca gccgggtggat gcggcggtc cgggggcccc 180
 ggcggcggtt cggcgccgt tccaagcac agcacctgg tggagcggct ccgccagcgc 240
 atcgagggct gccgtcggca ccacgtcaac tgcgagaaca ggtaccagca ggctcagggtg 300
 gagcagctgg agctggagcg ccgggacacc gtgagcctct accagcggac cctggagcag 360
 agggccaaga aatcgggctc cggcaccggc aaacagcagc acccgagcaa accccagcaa 420

gatgcggagg	ctgcctcggc	ggagcagagg	aaccacacgc	tgatcatgct	acaagagact	480
gtgaaaagga	agttggaagg	agctcgatca	ccacttaatg	gagaccagca	gaatggtgct	540
tgtgatggga	atTTTTctcc	gactagcaaa	cgaattcgaa	aggacatttc	tgcggggatg	600
gaagccatca	acaatttgcc	cagtaacatg	ccactgcctt	cagcttctcc	tcttcaccaa	660
cttgacctga	aaccttcttt	gcccttgcat	aacagtgga	ctcacactcc	tgggcttcta	720
gaagatctaa	gtaagaatgg	taggtccctt	gagattaaac	ttcctgtcaa	cggttgcagt	780
gacctggagg	atagcttcac	catcttgcat	agcaaagacc	tcaaacaaga	acctctcgat	840
gacctactt	gcatagacac	atcagaaaca	tctctttcaa	atcagaacaa	gctgttctca	900
gacattaatc	tgaatgatca	ggagtggcaa	gaattaatag	atgaattggc	caacacgggt	960
cctgaggatg	acatacagga	cctgttcaac	gaagactttg	aagagaagaa	ggagccagaa	1020
ttctcgcagc	cagcaactga	gacctctctc	tcccaggaga	gtgcgagcgt	gaagagcgac	1080
ccctctcact	ctcccttcgc	acatgtctcc	atgggatctc	cccaggcgag	gccttcttct	1140
tctggtctc	ccttttctac	tgtctccacg	gccactagtt	taccttctgt	tgccagcact	1200
cccgcagctc	caaaccctgc	aagctcacca	gcaaactgtg	ctgtccagtc	ccctcaaact	1260
ccaaaccaag	cccacactcc	aggccaagct	ccacctcggc	ctggaaatgg	ttatctcctg	1320
aatccggcag	cagtgacagt	ggccggttca	gcgtcagggc	ctgtggctgt	gcccagctct	1380
gacatgtctc	cagcagaaca	gctcaaacag	atggctgcac	agcagcaaca	aagggccaaa	1440
ctcatgcagc	agaagcagca	acagcaacag	cagcagcagc	agcagcagca	gcagcagcag	1500
cagcaacagc	agcagcagca	gcagcaacag	cactcaaate	agacttcaa	ttggtctccc	1560
ttaggacctc	cctctagtcc	atatggagca	gcttttactg	cagaaaaacc	aaatagccca	1620
atgatgtacc	ccaagcctt	taacaaccaa	aacctatag	tgctccaat	ggcaacaac	1680
ctgcagaaga	caacaatgaa	taactacctc	cctcagaatc	acatgaatat	gatcaatcag	1740
cagccaaata	acttgggtac	aaactcotta	aacaaacagc	acaatattct	gacttatggc	1800
aacactaaac	ccctgaccca	cttcaatgca	gacctgagtc	agaggatgac	accaccagtg	1860
gccaaaccca	acaaaaacc	cttgatgccg	tatatccagc	agcagcaaca	gcagcagcaa	1920
cagcaacagc	agcagcagca	gcagcagcag	ccgccacctc	cacagctcca	ggcccccagg	1980
gcacacctga	gcgaagacca	gaaacgcctg	cttctcatga	agcagaaagg	agtgatgaat	2040
cagccccatg	cttacgctgc	acttccatcc	cacggtcagg	agcagcatcc	agttggactt	2100
ccccgaacca	caggccccat	gcagtcctcc	gtgccccag	gctcaggtgg	catggtctca	2160
ggagccagtc	ccgcaggccc	cggcttctctg	ggcagccagc	ccaagcagc	catcatgaag	2220

cagatgctca ttgatcagcg ggcccagttg atagagcagc agaagcaaca gttcctgctg 2280
 gagcaaaggc agcagcagca gcagcagcag cagcagatth tggcggaaca gcagttgcag 2340
 caatcacatc taccceggca gcacctccag ccacagcggga atccataccc agtgcagcag 2400
 gtcaatcagt ttcaaggttc tccccaggat atagcagccg taagaagcca agcagccctc 2460
 cagagcatgc gaacgtcacg gctgatggca cagaacgcag gcatgatggg aataggacct 2520
 tcccagaacc ctgggacgat ggccaccgca gctgcgcagt cggagatggg actggccctc 2580
 tatagcacca cgcctaccag ccaaccagga atgtacaata tgagcacagg catgacccaa 2640
 atgttgccag atccaaacca aagtggcatg agcatcacac ataaccaagc ccagggacct 2700
 aggcaacctg cctctgggca gggggttggga atgggtgagtg gctttggtca gagcatgctg 2760
 gtgaactcag ccattaccca gcaacatcca cagatgaaag ggccagtagg ccaggccttg 2820
 cctaggcccc aagcccctcc aaggctgcag agccttatgg gaacagtcca gcaaggagca 2880
 caaagctggc aacagaggag cttgcagggc atgcctggga ggactagtgg agaattggga 2940
 ccattcaaca atggcgccag ctaccctctt caagctgggc agccgagact gaccaagcag 3000
 cacttcccac agggactgag ccagtcagtc gtggatgcta acacgggcac agtgaggacc 3060
 ctcaaccag ctgccatggg tggcagatg atgccatcgc tcccggggca gcaaggcacc 3120
 agccaggcga ggccaatggt catgtctggc ctgagccagg gagtcccagg catgccagcg 3180
 ttcagccagc cccagcaca gcagcagata cccagtggca gctttgctcc aagcagccag 3240
 agccaagcct atgagcggaa tgcccctcag gacgtgtcat acaattacag tggcgacgga 3300
 gctgggggtt ccttccctgg cctcccggac ggtgcagacc ttgtggactc catcatcaaa 3360
 ggcgggccag gggacgagtg gatgcaggag cttgatgaat tgtttggtaa cccc 3414

<210> 37
 <211> 678
 <212> DNA
 <213> Homo sapien

 <220>
 <221> misc_feature
 <222> (310)..(611)
 <223> a, c, g, or t

<400> 37
 tcataatgct gtcgagcggc ccgagtggt gatggatcgg ccgcccggca ggtacctgct 60
 gtgtggcagg ctctgggctg ggggctttat tcagcttct cagcctgctt cgacttcccg 120
 attagagagc taatgtgaat caccaaccct gtgatgcctc ttgagatgag agttcagatt 180

tccaagaag atctaagcag ttggtccaaa ttgtagttca ctagcaaatg acccagtgtc 240
 gtccctgtgg tgtgtttatg acatgatgga agatgctgcc ttcaaaagtg tccacttgta 300
 agaagatgtn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 360
 nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 420
 nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 480
 nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 540
 nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 600
 nnnnnnnnnn nacacacaaa aaaaagggtg ggaaccagg gcaagcggc ccgggggaaa 660
 tggtttccgc acattcca 678

<210> 38
 <211> 461
 <212> DNA
 <213> Homo sapien

<400> 38
 gtgcgccga ggtaccaact gacatttcag tttttctggt tgaagtccaa tgtattagtg 60
 actctgtggc tgetctcttc acctgccct tgtggcctgt ctacaattct aatggattt 120
 tgaactcaat gtogtcgctt ctggtttctt gcataacca atagcattac ctatgacttt 180
 tttttctctg agctattttc actgagctga gctaataaac taaaactgag ttatgtttaa 240
 tatttgtatc aaatacataa aaggaatact gctttttctt tttgtggctc aaaggtagct 300
 gcattttaaa atatttgtga aaataaaaac ttttgttatt agaaaaaaaa aaaaaaaaaa 360
 aaaaaaaaaa ggcttggggg aaaccgggg ccaaaagcgg tgtccgggg ggggaattggt 420
 ttctccggtc caaattcccc aaaaaaatcg agaagaaaag t 461

<210> 39
 <211> 633
 <212> DNA
 <213> Homo sapien

<400> 39
 caacaccatc tttttttttt tttttttttt ttgagacaga ttcttactct gcactccagc 60
 ctggtgacag agcgagattc catctcaaaa aaaaaaaaaac agtatgcacg tacaatttc 120
 ttaacctgtt atcaatgtct gagctacata attatcttct tagttggagt ttgttttagg 180
 tgtgtaccaa ctgacatttc agtttttctg tttgaagtcc aatgtattag tgactctgtg 240
 gctgctctct tcacctgccc cttgtggcct gtctacaatt ctaaattgat tttgaactca 300

33

atgtcgtcgc ttctggtttc ctgcatatac caatagcatt acctatgact ttttttttcc 360
 tgagctatth tcaetgagct gagctaataa actaaaactg agttatgtht aatatttgta 420
 tcaaatacat aaaaggaata ctgctttttc cttttgtggc tcaaaggtag ctgcatttta 480
 aatattttgt gaaaataaaa acttttgtha tttagaaaaa aaaaaaaaaa aaaaaaaaaa 540
 aaggcttggg ggaaaccagg ggccaaaagc ggtgtccagg 'gggggaattg gtttctccgg 600
 tccaaattcc ccaaaaaaat cgagaagaaa agt 633

<210> 40
 <211> 536
 <212> DNA
 <213> Homo sapien

<400> 40
 ggggccgccc gggcaggtac ttgacagtgt tatctgtcac ttatttaaaa aaaaaacaca 60
 aaaggaatgc tccacatttg acgtgtagt ctataaaaca cagaatattt cattgtcttc 120
 attaggtgaa atcgcaaaaa atatttcttt agaaacataa gcagaatctt aaagtatatt 180
 ttcatataac ataatttgat attctgtatt actttcactg ttaaattctc agagtattat 240
 ttggaacggc atgaaaaatt aaaatttcga tcatgtttta gagacagtgg agtgtaaate 300
 tgtggctaate tctgttggtc gtttgtaata taaatgtaaa atagtattcc agctattgtg 360
 caatattgaa atagtgtaaa taaacacaag taataaatga agtgtttgtt ataaaaaaaa 420
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaagg gtggggggaa cccggggcca aaaggggttc 480
 cgggggggaa attggtttcc gggccaaaat ttccaacaat ttgggagaaa aaaggt 536

<210> 41
 <211> 1206
 <212> DNA
 <213> Homo sapien

<400> 41
 gtactctccc aatgcagcc taatcttagt aacctgaag tttatcattc tttaaaacta 60
 aatagaatac caatggttta gatattccaa caaagaatgc tagaaacaaa tgtctaate 120
 cgattattag ctttaccaac cctgtgaaca ctgaggttgc agaactgcca ggttaatecc 180
 tgtggcctag actactgagg attctgatag cacatgtaag actaagcact cttcaagctg 240
 taataaagca tccacatgta tctgtgatga ttttcattgc tttagcattg cagccatgta 300
 acaactgcag aaagaaggta tttttaaaaa tacaatagac tacacttttt ggatcacaga 360
 gaaatacaga tgcactctga gactgcctat gtttataaac atgtttgtgc ccctaactga 420
 agtgacaggt cttctggaat tgacattaag aagtgtggat agtcatatca cacgcaatgt 480

atttgttttc agcagtgagc agaccgtaca ggagcagcac accaggagcc atgagaagtg 540
 ccttggaac caacagggaa acagaactat ctttatacac atcccctcat ggacaagaga 600
 tttatTTTTg cagacagact cttccataag tcctttgagt tttgtatgtt gttgacagtt 660
 tgcatatata tattcgataa atcagtgtac ttgacagtgt tatctgtcac ttatttaaaa 720
 aaaaaacaca aaaggaatgc tccacatttg acgtgtagtg ctataaaaca cagaatattt 780
 cattgtcttc attaggtgaa atcgcaaaaa atatttcttt agaaacataa gcagaatctt 840
 aaagtatatt ttcatataac ataatttgat attctgtatt actttcactg ttaaattctc 900
 agagtattat ttggaacggc atgaaaaatt aaaatttcga tcatgtttta gagacagtgg 960
 agtgtaaate tgtggctaate tctgttggtc gtttgtatta taaatgtaaa atagtattcc 1020
 agctattgtg caatatgtaa atagtgtaaa taaacacaag taataaatga agtgtttgtt 1080
 ataaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaagg gtgggggggaa cccggggcca 1140
 aaaggggttc cgggggggaa attggtttcc gggccaaaat ttccaacaat ttgggagaaa 1200
 aaaggt 1206

<210> 42
 <211> 209
 <212> DNA
 <213> Homo sapien

<400> 42
 ccgggcaggt ggaacttagt gggcagcatt acgggcagcg ctaaggaacc atttaaagta 60
 agacaagtcc acacagctgt ggtgcttttc tacgagtctt gttccaactg ctgcataaca 120
 atagaatggt ggaagcagga attagtttta aagtaagact tcagaagtgg aaacaaattt 180
 gatatttatt tttataatga tataatagc 209

<210> 43
 <211> 706
 <212> DNA
 <213> Homo sapien

<400> 43
 gaaccctcca aaacatctga aaagcaaatt tgggggggatg aggaagtgag atgatgactt 60
 gattctcctt ctaggaagaa tagaggaacc cttctggcaa aatttcaagc atctacaaga 120
 ggaggttttc cagaaaataa agacactggc tcagctctca aaggatgttc aggatgtcat 180
 gttctacagt atcctggcca tgctcagaga cagaggggct ctacaggacc tgatgaacat 240
 gctggaattg gacagctcag gtcatttggg tggccctggg ggtgccatcc taaagaaact 300

35

tcaacaggat tcaaaccatg catggtttaa cccaaaggac cccattcttt atctccttga 360
agccataatg gtgctgagtg acttccaaca cgatttgctg gcctgttcca tggagaagag 420
gatcctgctt cagcaacagg agctggtaag gagcatcctg gagccaaact tcagataccc 480
ctggagcatt cccttcaccc tcaaacctga gctcctcgcc ccaactccaga gtgagggttt 540
ggcatcacct atggctgctg gaggagtgtg gccttaggac ggagctggat aaccccagg 600
caacctggga tgtagaagca aagatgccct gtctgtcctc tatgggactc tctcgttct 660
gagcagtggg tgaaggctaa gcctccctga tgggagcagt cagaaa 706

<210> 44
<211> 1298
<212> DNA
<213> Homo sapien

<400> 44
atatgaagtt aaaaccagag ctatttctga cacagcaatt tttgagcggg catttgccaa 60
aatacgaaca agttcacatc ctcccagtag gtgagtgtga gtttgctgga ggtgggggtg 120
gggatcccat cctgcacaca tggggtaagt agggcagatt gccctgcct cgcctttgcc 180
accaccgccc tagggcctgg cgtttggtca tgtggaatgg gaagggcca gaaagctgag 240
aacatggagg atgaatggga atggggcag gaagaagtg agtaagagg aggaggtggt 300
aggagagcag aaccctcaa aacatctgaa aagcaaattt ggggggatga ggaagtgaga 360
tgatgacttg atttctctc taggaagaat agaggaacc ttctggcaaa atttcaagca 420
tctacaagag gaggtttcc agaaaataaa gacactggct cagctctcaa aggatgttca 480
ggatgtcatg ttctacagta tcctggccat gctcagagac agaggggctc tacaggacct 540
gatgaacatg ctggaattgg acagctcagg tcatttggat ggccctgggt gtgccatcct 600
aaagaaactt caacaggatt caaacatgc atggtttaac ccaaaggacc ccattcttta 660
tctccttgaa gccataatgg tgctgagtga cttccaacac gatttgctgg cctgttccat 720
ggagaagagg atcctgcttc agcaacagga gctggttaagg agcatcctgg agccaaactt 780
cagatacccc tggagcattc ccttcaccct caaacctgag ctctcgcct cactccagag 840
tgagggtttg gccatcacct atggcctgct ggaggagtgt ggcccttagga cggagctgga 900
taaccccagg tcaacctggg atgtagaagc aaagatgccc ctgtctgccc tctatgggac 960
tctctcattg ctgcagcagc tggctgaggc ctaagccctc cctgatgggc agtcagtcca 1020
gagatgctgg cctcgccea gtctatgctg tgagtgtcct tatgggtgca agagataggg 1080
ctgtgcctct ctgcgtttcc aggtggagta gagacagtaa tgggtagaga ctttaggaaa 1140

36

tgttttgggg tgggtggaata ctctatatat tgacaagagt ttatatattg acaagagttt 1200
 atatatttgt caaaactcct caaatagtat gttaaagacg taagcgtttc actatgtata 1260
 aattttactt caaaataata aaaacaaata ctgactct 1298

<210> 45
 <211> 531
 <212> DNA
 <213> Homo sapien

<400> 45
 acaacattca aacaaccagt ggtgaggttg taaatcaaat gagagaggag gaactgatcc 60
 gggtagcagg aacacatttc caagtaaaat ttgcaacaga gcatggtgag atcatggttt 120
 taatttatga atggcattat tatctttaa ctattatttt ccaagctcat atatggcctt 180
 tttgaagggt ttccgaatgt tacatttgat ttttaagatct aatccaaaat gaaatataga 240
 atgtgcttag ttttctataa aaatgccaat gactatctct taaattagtc aaggaaagac 300
 aaattaccaa aattcaact tatttgaatt atttttaagt gattccaggc aataaataca 360
 tagaacccat ggaaagtttt agcttcaaat cacaaaattg caaaaaaaaa aaatggtaaa 420
 tggctaaca taaggggggt tatggaaaat attgggtcac ctttaattata ggtttaaatg 480
 ccacaaacaa tataataata gttttaactt actttttctg attactaagc a 531

<210> 46
 <211> 469
 <212> DNA
 <213> Homo sapien

<400> 46
 taacgccatc agctcgctgc ttaaagccgt gtttgcgtct ctttttctca aagaaatctg 60
 ctttagtttg agattacagt ttatcaaatg ttaaggcttt gacccccaaa tctggtccca 120
 gaaagacagg aaggccagct aagaggaggt tttcagagtg cgtagaaagg ctgctctgtg 180
 cttcggcatt tgttctggaa gtgcttcttc ggttgccaaa gattcctagc aaaacctttg 240
 actggaggct ttacagggcc atacacccaa tatcactaat gacagtgttg taaaatagct 300
 tttgtgcacc atgcttagga ttcaaggag ataaagtata tctttctaaa gttatacttt 360
 agaaaactgtc attccatgtt gaaatgataa acattccatg tttatctttt gtgtaagaag 420
 taaaaagca aaaattcatt gcatcaaagt aggtcaggca ctgctaaag 469

<210> 47
 <211> 483
 <212> DNA
 <213> Homo sapien

<400> 47
 aaaccgagtt ctggagaacg ccatcagctc gctgcttaa gccgtgtttg ctctcatttt 60
 ctcaaagaaa tctgttttag tttgagatta cagtttatca aatgtaagg ctttgacccc 120
 aaaatctggg cccagaaaaga caggaaggcc agctaagagg aggttttcag agtgcataga 180
 aaggctgctc tgtgcttcgg catttgttct ggaagtgctt cttcgggttg caaagattcc 240
 tagcaaaacc tttgactgga ggctttacag ggccatacac ccaatatcac taatgacagt 300
 gttgtaaaat agcttttgtg caccatgctt aggattcaag gaggataaag tatactcttc 360
 taaagttata ctttagaaac tgtcattcca tgttgaaatg ataaacattc catgtttatc 420
 ttttgtgtaa gaagtaaaaa agcaaaaatt cattgcatca aagtaggtca ggcactgcta 480
 aag 483

<210> 48
 <211> 600
 <212> DNA
 <213> Homo sapien

<400> 48
 tccatttctc atggcttgct catcttccgg cttcaggctc tgacttcac tcaggatggg 60
 atcgggtgtgt gtctgttttc atagatccac tacatcagaa gtatctttac atctctgtat 120
 ctttacatcc caaggtcaag gccctggcaa cctcagaggt tcccatagct tcagtcttcc 180
 ccaaaccatg ccaacttctc ccatttcttt gggtcaggaa tctggctttt gttttccata 240
 tttctttttc ccaagacatt gggaggcatc tggggaacaa caccaataaa acagttctct 300
 cccccaaaa aaaaaaaaa aaaaaaaaa aaaaaaaaa aaaaaaaaaac aaaaaacgaa 360
 gaacaaagaa cagagaaaaa aaaaaacaag aaacaccaa aaacaaaaaa gaaaacgcgg 420
 ccgccagcgc acgcgcgagg ggcgccgagc acaccctgtg gccagcccgc gagcgagaag 480
 ggagcgggcg gggcgggcg gaccggagac ccaaggaggg cgcagggagc aacgaacggg 540
 agccggagga gcgcgacact gcacgcagga gagcagacgg gaggggagac agcgcgggga 600

<210> 49
 <211> 1098
 <212> DNA
 <213> Homo sapien

<400> 49
 aacctcttca acaataaatt gctctttggg gacattttat gcacagaact gtgcaccctc 60
 ctcagaacag caggtcttta atggcccatg tgatgagaag ggccccatca aggcagcagg 120
 aatgggccac tctccacac cccatgggcc aggccactgc cactcctgct gccttgcac 180

cccaggttta tggctgcatg gtagaagtca cttctgtaag aaattcacct ttctaaaata 240
 aagtatgctc ttttttctga gacatctata gaataacttg tggcagagtg ttttaaaaac 300
 tgatttggat tttttttatc ctttaaccgt gtgaaaggat ggaagggatt ttaggtggaa 360
 gagaagttaa gaacagaaaag atagagcagg ttttttagagt gggagaatta atcccaaaga 420
 aaaagagggc atggaaacaa atgtggatgc catgggctct gtgccagact tgccagtgct 480
 gactggaaca ggccgggctc ctcaactcagc ggctcctgcc tcagctgtgg ttcccgcagc 540
 ctctgggtct cacggaacct ttccttggga gttccatttc tcatggcttt gctcatcttc 600
 cggcttcagg ctctgacttc atctcaggat gggatcggtg tgtgtctgtt ttcataagtc 660
 cactacatca gaagtatctt tacatctctg tatctttaca tcccaaggtc aaggccctgg 720
 caacctcaga ggttcccata gcttcagtct tccccaaacc atgccacttc ctcccatttc 780
 tttgggtcag gaatctggct tttgttttcc atatttcttt ttccaagac attgggagggc 840
 atctggtgaa caacaccaat aaaacagttc tctccccacg gtcattccagg tcaactctct 900
 aactcattcc tgcacacaca gcacacgtgg aatttgcttg tttagtctat gttcttgact 960
 tgatcacaga cgctgtaca ataaagcccc ttttcaacaa ggtgctgcag aatgataatg 1020
 ctttcccaa aatctgaaac tgatttgtat cattgaagtt tttttctgta ttaaaaataa 1080
 agcaaaaatta aaaataaa 1098

<210> 50
 <211> 540
 <212> DNA
 <213> Homo sapien

<400> 50
 ggtcgcggcc gaggtactcc cgcctcctgg agcggccgac cccacatgga ttctcaacag 60
 gtggccggca catcttctga gcctcgtctc ctcatctgaa agtggagtgt aagtccaaga 120
 agattcattt agacaaagaa ggtggaaaaa aaggactttc tgggccagca agtcggatga 180
 ccacctcca aggggcagag gagggcccat tttgtgaaga agaatcaac taccggaaa 240
 acgccacagg aggacatggt tctgcagatg tagttgccct agaaacagaa gagtatgggg 300
 gtgtgaatgt cttctctttt gggggcaaac actatgtcct tttcttttct tagatacagt 360
 taattcctgg aaatttttagc gagtttggtc ttgtggatat tttgaacaat aaagagtgaa 420
 aatcaaaaaa aaaaaaaaaa aaaaaaaaaa accctgggag gtacccatgg cgcaaagcct 480
 ggtcccctgg ggggacactg ggttaccgg ccccaattc cccacaattg cggagcaacg 540

39

<210> 51
 <211> 1028
 <212> DNA
 <213> Homo sapien

<400> 51
 cggccgcggc atgaaaggcg gcgaggagag gcagcactgc tgctcttgac ttctgagcag 60
 ggcttagaga gcctgccccg gcttaagccg agctgctggg gctgaccctg agcgcgcgagt 120
 ccgcgagctc tgagtccgga gcctcccagc cgtggagccg tgggatgagg ggggcgttgg 180
 gggacagggc aaagtcgatc ttggtgtac agccgccoga tcctagcgcg gagctgcgag 240
 cctgaccggc cgcgtctggc atggtcagag aaagaatctt cttttccca ctccggcttt 300
 tggttttgtg tgtccacctt gcgcaactcc ggagccagcc gacccacat ggattctcaa 360
 caggtggccg gcacatcttc tgagcctcgc tctctcatct gaaagtggag tgtaagtcca 420
 agaagattca tttagacaaa gaaggtggaa aaaaaggact ttctgggcca gcaagtccga 480
 tgaccaccct ccaaggggca gaggagggcc cattttgtga agaagaaatc aactaccgg 540
 aaaaacgccac aggaggacat gtttctgcag atgtagttgc cctagaaaca gaagagtatg 600
 ggggtgtgaa tgtcttctct tttgggggca aacactatgt ccttttcttt ttctagatac 660
 agttaattcc tggaaatctt agcgagtttg ttcttgggga tattttgaac aataaagagt 720
 gaaaatcact ttggagtcac ttaatcttcg ttagaagggc agtttcttcc agggccattt 780
 tctttcacca gatttgtttt tcctcgttcc caaatgaggt agttttaaaa atcaaagtcc 840
 acttgctaac tcacctggga aagagactgc gacagaagga agagaagtaa atagacatca 900
 ctctcaaact aaaagtgtaa ctttcattcc tggcagctga gattcagaac acaaagaac 960
 aaactcgttt acctttgagt atttccccg tatgggtaat ttatctagag ctttcccaac 1020
 aattaatc 1028

<210> 52
 <211> 541
 <212> DNA
 <213> Homo sapien

<400> 52
 acagattggg aaggtagacat tgtatcacia agctagtctt tgagtccaaa gttttgtggg 60
 tttatgttat gatatacttt tatcatggaa ttgtcttatt aaatgttttg ccagtgggtc 120
 ttaaagtgtg tttctgacac cagtagcatt gacttcactt agaaacctgt tagaaataca 180
 aattatgttg cccacccaa cacttgagtc acaaactttg cagatggggc tcaatctggt 240
 ttaacaagcg cttcatgtaa ttttgatgca ggcctaagtt tttgagccgc tgcagtatgc 300

40

atctctatctt ttaagcaaag atcttggctc ttctttttgg acattgtaga aataacatga 360
 acttgtcctt tgtttgtttt ggttttggtt tgttttaagc tcctgatcct tgttggttat 420
 gttgcaaaag attgtatcag gagaagcctc agcatggaca ttggcatcct gacataaccc 480
 ccattaatct agtattcctt ctgaaactca aatggattct caagtccaag agactatgga 540
 a 541

<210> 53
 <211> 261
 <212> DNA
 <213> Homo sapien

<400> 53
 atgccatcag tggcacaggg ccctgtgccc tggcatctgg gttcacgctc tgctgttct 60
 gtcttogaat tcctagtgat gtttgaacaa aggccctatg tttgcatttt gcactgggcc 120
 ccacaaatca catggcccat cctgagaaga ggagtctcac acctccagtc tctaaatca 180
 cctctggaag tttttctcaa cgaagaact gaagctttcc tcaaagttc cgtaggggag 240
 acagttcatc accatacca a 261

<210> 54
 <211> 325
 <212> DNA
 <213> Homo sapien

<400> 54
 gctctgtttt gtgttttgtt tggattgtgc tggttgtggt ttgtgtttgt ggaaggtgtg 60
 tgtgtgggtt tggcgagtac atgtcgcccg ggaccgctat ggctctgggt gcgcccacgc 120
 tttttttttt tttttttttt tttttttttt ataatcaacc tataagggat ttatcaataa 180
 ataaaccctt atttattata aggaattggc ttacacaata atggaggccg agaaggcccc 240
 aagtctgctg tccgaaggtc tgagaaccag gagcactgat ggtgtcagtc ccagttcaag 300
 ggcaggagaa gatgggtgtc ccagc 325

<210> 55
 <211> 2461
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (356)..(393)
 <223> a, c, g or t

<400> 55

gcctgaatag agctgtgcag cccaaggggt ggactgagcc agcagtggat atgcaccact 60
gagatctctt gctgtggaac gtaattgact ggggggggtcc ccgctactgc tctctgaatc 120
cattgataca gtcatgccaa ggctacattt cccatggggt gtttccataa gaataacaat 180
aactgaatga agaaggtata ctaataatgc aggcctatc ctgtgaggta gggggctcct 240
ccaatgggag actttgggtt gagtgttctt catcagctga ccttaaactt tattggaatt 300
gtgctacagc ctaagcttcc tgctactcaa cccgccttcc ttcctctct ccttcnnnnn 360
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnaaaccta tggctctatc aatcaattaa 420
taaggattta taatcaacct ataaggattt atcaataaat aaacccttat ttattataag 480
gaattggctt acacaataat ggaggccgag aaggcccaa gtctgctgtc cgaaggtctg 540
agaaccagga gcaactgatg tgtcagtccc agttcaaggg caggagaaga tgggtgtccc 600
agcggcacag tcaggcagaa aattcaagct tctccacct attttatttg ggtccttaga 660
agactggatc aagcccatcc aactgggga ctgcaatctg ctttattcag tccatcaatt 720
caaatgcaaa tttcctccag aaaaagcttc atgaacacaa ccagaaataa tgttcgatca 780
tatatggggc atcctgtggc ccagcgaagt tgacatgcat aaaattaacc atcacacctc 840
catggaggcc agaaatttta tgtattctga ggactctgtc tctctggctt cctctccatt 900
tcaaccctca ggcattcttt ttttcccaa ataatgtct tgtaaacta ttttttttt 960
acatgtgttt ttttggagt actggaacta ataaattgg actttgttct gcaggtagtt 1020
tagttactta cagatcagct ttatactatt gaggctgtt ttaagcttt ttgtgggggt 1080
gaggagcaag tgtagttag tttttagtca ggatagtta gccctacgta tgtaacacac 1140
aacctttctg ggatcttag cgaatgcctg gtttctagtc aggactctcc actgtggctg 1200
gatggaaatt agatatttcc caatctcacg ctaatttga gaattgtag gcttagtggt 1260
ccgcaatgta cacaatggg tctctacata gatttctgga tttttttgtc taccttctc 1320
cttgttgata ctttgcctg aaaattcaag ctatctcagt atcaactcca atcttctct 1380
cctcaactca gcaagactct gatctcctgc gatgggatcc agagtgtca cctcagaaag 1440
gcagggcgat cggagggtctc actttgttcc cctgcctcag cgatcaaatt gtataaactt 1500
tttccctggc ttctggaac actgatagcc aacctttcgg taaaatttct aacatagta 1560
cttagagctt taatagcta agtatttaa tgctaagtat tttctttaa attcatttca 1620
aaatattttt gcaattttgc cgtgatttat ttttggctc atgggtcatg aggtgtatgt 1680
tccttaactt aatgcatttt ggaaattttt tgttaccttt atgtacttga tttctgggtt 1740
aattcactat ctatatgatt tcaatctttt gaaatttctt gagacttgat ttgtgatcca 1800

gcctaacatg cgccccagaa cacatggtaa atgttttggg gtatacttta aaagaacatg 1860
tattctgatg tttttgaatg taatatctta tgtcttattc atatttatag taccaataca 1920
cagcacatag tagaaactta acatatattg agttaaataa ttcaaagggt ttatccgatt 1980
gggtggcaatt caagaccaa taagagagga tgatgatgac atcactattt ctgtaagac 2040
agggcctcat acaacataca ggaatgtcca gttgtcaagt catgcagttt ctcccctatt 2100
ctaaccaatg ttaatgcaa tactttgtga tgaaaattat cccagtattt ttcctcctat 2160
gtttccacca gttttccctt cattgattgt taagatttat atcactagag ctatttgaca 2220
gtaggaaaca attaccttag gaaaagttgg tgacattggg ctataaagggt cacggagaca 2280
taagaaatgg ttattttttc atttttcacc aaacaattca cgattgtttc taagattaca 2340
aaagattaga cgatagctaa tatttctatg caatggtaa atttttcaag tagaatcatt 2400
tttaaatttt ccaagttcca atgtcacttt ctcttgaac acgactcaag gtcaaaactt 2460
a 2461

<210> 56
<211> 643
<212> DNA
<213> Homo sapien

<400> 56
ccgcccgggc aggtacacat gagtgcgtgt atgccccag gctgggtcag ctcttctgtg 60
gattgcatgg cgtgtgatta aaagcccatg tgttcccaca catccacatc atgggaagggt 120
taatgtgtgc ctcttggaa ctgggtgtg gtgtccatgg aacttctct ctgtatctca 180
ggtcagtagg cgcagaaacg cctcatgatg aagattcttg agccccattt ccaagacccc 240
tcacatocaa tcctgtcctg taacatccat caaggatttc cataggggtg actggtgccc 300
accaagact gcaccagtgc ctgctcattg aggagagtaa ctgctggcca ggcagaaaga 360
atatgggctc tgcaatgaga cagacctgga ggggactctc ccgttgagca ctagcagctg 420
gaggagtgg gagttcatgg ctatcatggg tgtgttcaat cgattgtggg gatgacatgt 480
cattgtgtat ggaaggcggg gctcatggct gattggccaa taaaatggcg gctgccgttg 540
tcattgaaaa aacacaccac accacaacca aaaccgctgg ggcacaccg ggcacaaggc 600
ccccgggga aacgggttcc ccgccc aaat tctccaaatt aga 643

<210> 57
<211> 1611
<212> DNA
<213> Homo sapien

<400> 57
 ctcctcccga ggaaccagtg gtgacagctg aggccatgtg agtaggatcc tgaatgagggc 60
 tttatctctg gctgttcgtc ccatcgtcca ccgtggcacc agctccctca gccagccggg 120
 atgggaccag cgactgagag agccagagggc agagaggtga ggggtgacat atcctggact 180
 gtgagaggaa tgggactctg ggcctgtagc tgccaagcag gtggcaggtg ctccaggctg 240
 tgatctgcac cctctgacct ctgacattga cctcctacct tgaccctgc ctgaccaagc 300
 catgtctgaa caggaggctc aagccccagg gggccggggg ctgcccccg acatgctggc 360
 agagcaggtg gagctgtggt ggtcccagca gccgcggcgc tcggcgctct gcttcgtcgt 420
 ggccgtgggc ctcgtagcag gctgtggcgc gggcggcgtg gcaactgctg caaccaccag 480
 cagccgctca ggtgaatggc ggctagcaac gggcactgtg ctctgtttgc tggctctgct 540
 ggttctggtg aaacagctga tgagctcggc tgtgcaggac atgaactgca tccgccaggc 600
 ccaccatgtg gccctgctgc gcagtgggtg aggggcccgc gccctcgtgg tgctgctcag 660
 tggcctcgtg ctgctggtca ccggcctgac cctggccggg ctggccgccc cccctgcccc 720
 tgctcggccg ctggccgcca tgctgtctgt gggcattgct ctggctgcct tgggctcgtc 780
 tttgctgctg ggctgctgc tgtatcaagt ggggtgtgagc ggacactgcc cctccatctg 840
 tatggccact cctccacctc acagtggcca tggcggccat ggcagcatct tcagcatctc 900
 aggacagttg tctgctggcc ggcgtcacga gaccacatcc agcattgcca gcctcatctg 960
 acggagccag agccgtcctt cttctcacag cggcctcagc gtccccagag ccgagccagg 1020
 gtgtgagtgc atgtgaaagt tgagtacaca tgagtgcgtg tatgccccca ggctgggtca 1080
 gctcttctgt ggattgcatg gcgtgtgatt aaaagcccat gtgttcccac acatccacat 1140
 catgggaagg ttaatgtgtg cctccttggc actgggtgtt ggtgtccatg gaacttcctc 1200
 tctgtatctc aggtcagtag gcgcagaaac gcctcatgat gaagattctt gagccccatt 1260
 tccaagacct ctcacatcca atcctgtcct gtaacatcca tcaaggattt ccataggggt 1320
 gactggtgcc cacccaagac tgcaccagtg cctgctcatt gaggagagta actgctggcc 1380
 aggcagaaag aatatgggct ctgcaatgag acagacctgg aggggactct cccgttgagc 1440
 actagcagct ggaggagttg ggagttcatg gctatcatgg ttgtgttaat cgattgtggg 1500
 gatgaaatgt cattgtgtat ggaaggcggg gctcatggct gattggcaat aaaatggcgg 1560
 ctgccgttgt cattgtctcc aaaaaaaaaa aaaaaaaaaa aaaccgcgga c 1611

<210> 58

<211> 617

<212> DNA

<213> Homo sapien

<400> 58

```

actgtgaagt cttcaggctc ttāgaaggct ccagcctgag agagcccttt attattgcca      60
ttcctgtcct tcctcaaggc ctggtgacct gtgacctttc gctctgggca gggcccaggct    120
agatggggccg tcatccgggc ctgtaagccg tactatgatt tctgcattga tttacatatt    180
ttttactgtg atccttggtc caaacacaga atcgtcacc cttctccct tgaatgtgcc      240
ggatccttgt aaattctcat tcacctactt gttccttaggt gtgtatgtgt gtgcgaaact    300
ctatgttcaa gaaagaaatc atacaaagag taacgaacca tggttctgtt ggccattgga     360
cgaaacttgg tttttggact ttcttaccta acattaattt tgcctctgcc tgggtttaca    420
cacacacaca cactacaaca aacacaacac aaacaacggt ctgggccaac accacgcggc     480
gccagcgccg gctccctggg ttgaaacttg gatctcttcc cgcgccacaa ttctcccaac    540
aactataatg agcacaagga ccacaacat acacaagaac aacacaaacc agcgacacaa     600
cagagacaac acacaac                                     617

```

<210> 59

<211> 913

<212> DNA

<213> Homo sapien

<400> 59

```

caaaaccaca cccatgcaca cacataccct cagccccac acacaccccg ttgaaccogt      60
gagtctatca gggcatccta aaactccgtg agttgacatt tcagtaattt caggggaagg    120
tgttttccag ggatggggtc tcccagggtc agatagtgcc tttggctgca aatgctcctt    180
tagctaaact tttcctcagg aagaattcat tattctagac attatgtgat atatctgtta    240
ggaataaaag gtgcttaacc ttctccctg ggatgtggga gaagggtctg gaggtgtac     300
tgtgaagtct tcaggctctt agaaggctcc agcctgagag agccctttat tattgacatt    360
cctgtccttc ctcaaggcct ggtgacctgt gacctttcgc tctgggcagg gccaggtag     420
atggggccgc atccgggcct gtaagccgta cttgatttct gcattgattt acatattttt    480
tactgtgatc ttggttccaa acacagaatc gtcacccat tctcccttga atgtgccgga     540
tccttgtaaa ttctcattta cctacttgtt cttagtgtgt atgtgtgtgc gaaactctat    600
gttcaagaaa gaaatcatac aaagagtaac gaaccatggt tctggtggcc attggacgaa    660
acttggtttt tggactttct tacctaacat taattttgct cttgcctcgg tttacacaca    720
cacacacact acaacaaaca caacaaaac aacgttctgg gccaacacca cgcggcgcca     780

```

45

gcgccggctc cctgggttga aacttgatc tcttcccgcg ccacaattct cccaacaact 840
 ataatgagca caaggaccac aaccatacac aagaacaaca caaaccagcg acacaacaga 900
 gacaacacac aac 913

<210> 60
 <211> 554
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (304)..(430)
 <223> a, c, g or t

<400> 60
 tggaaaataa agtttaaac cagattgccc agagcaagac tctaattgtc ccaacgggtga 60
 tgacatctag ggcagaatgc tgccattttg aggggcaggg ggtcagctga tttctcatca 120
 agataataat gtatggtttt tactactaagc aactgataaa tggacaattt atcactggac 180
 aatctccctc tgcttcttta atggggccag ctttgcagcc ctgcagcctg ggtagtgcga 240
 cacatttcca tgcattcaag gccccctgtc ttggggagaat gatctgctag tgccatttta 300
 aatnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 360
 nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 420
 nnnnnnnnnn tcaactgtgtc cggcataaag tagaacattc ttacaagaaa taaatatttc 480
 gtagtcatgg agaagaacgc gaaaaaaaaa aaaacaaaaa aaaggctggg ggtaaccagg 540
 gcccaagcgg ttcc 554

<210> 61
 <211> 1401
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (803)..(929)
 <223> a, c, g or t

<400> 61
 aattatcttg ggtctctgtt caaatgagtt tggagaatgc ttgacttggt ggtctgtgtg 60
 aatgtgtata tatatatata cctgaataca ggaacatcgg agacctattc actcccacac 120
 actctgctat agtttgcgtg cttttgtgga caccctcat gaacaggctg gcgctctagg 180
 acgctctgtg ttcactgatg atgaagaac ctagaactcc aagcctgttt gtaaacacac 240

taaacacagt ggcctagata gaaactgtat cgtagtttaa aatctgcctc gcgggatggt 300
 actaaactcg ctaatagttt aaaggttact tacaatagag caagttggac aatthttgtgg 360
 tghtggggaa atggttagggc aaggcctaga ggttcatttt gaatcttggg ttgtgacttt 420
 agggtagtta gaaactttct acttaatgta cctttaaact agtccatttt ctatgttttg 480
 tataatctga aactgtacat ggaaaataaa gtttaaacc agattgccca gagcaagact 540
 ctaatgttcc caacgggat gacatctagg gcagaatgct gccattttga ggggcagggg 600
 gtcagctgat ttctcatcaa gataataatg tatggttttt aactaagca actgataaat 660
 ggacaattta tcaactggaca atctccctct gcttctttaa tggggccagc tttgcagccc 720
 tgcagcctgg gtagtcgcac acatttccat gcaccaagg ccccatgct tgggagaatg 780
 atctgctagt gccattttaa atnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 840
 nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 900
 nnnnnnnnnn nnnnnnnnnn nnnnnnnnt cactgtgtcc ggcataaagt agaacattct 960
 tacaagaaat aaatatttcg tagtcatgga gaagaacgct cctaaaatga tgaacgcacg 1020
 acggaaaaga gtagggaaca ttttgcttga tgagaaaatc cgccagcaag gatgttgggc 1080
 tctaagcaga actgaagctc tggaaattaag aacacagcca aggaagagct ctggactctg 1140
 agtttaaaga agctgactga cttgtaaggc aattccaggg aagattggtg aatcaagtta 1200
 agaatacaaa gcaactgaga tcaactgga ggcctggaag gtaaggcca tattttacct 1260
 agatactagc ttagagactt gctacattgg cactgtattt taagtatgtt atttagtagt 1320
 attgtgaaat caactggtt caacattgaa aaggataaaa atagcttatg aaaacaaac 1380
 ggthtttttt tttttttaa a 1401

<210> 62
 <211> 568
 <212> DNA
 <213> Homo sapien

<400> 62
 agatgctgcc gagcggcgca gtgtgatgga tagtcaaaa aaaaaaagta ttaaaatgtg 60
 attgatgtaa tttaccatgt ttactttatg catgcatttt attggggagg ggacgtgtca 120
 gaataataca cccaaatcta gtggctctaat ttcatagtgc taatctggtt tatattggca 180
 ttaaacgata ctgccaagga gctagatcat tttacaagag ttgtaggttt gtcttatggt 240
 ggaaaagcag tcctctatta atatcatgtg tgaagagtat ctgttcacaa gatttatgag 300
 attatgacgt gtttcagaga atgtctacta gtatatcttt acagtatttg cctgttgaac 360

47

tccctgcaca aactggaatt actttccaga agacttaggg aatgcaaata tggtactcat 420
 aagatgcatt ggagtatggg aaataaaaca aaccattttg gattggttta aattggetcg 480
 ttacagttct cttgtgggga gggactttgt cagtcatttt ggcatcttaa gctagactaa 540
 actttttggt gttgttttcc taaaacca 568

<210> 63
 <211> 791
 <212> DNA
 <213> Homo sapien

<400> 63
 tggctatgg taatTTTTta tagcagtc cc agccaagaca gtgcgctcat ttactacata 60
 ccatttatat tattatatag gctcctttca gaaaccatg ttcaaataag agataagata 120
 ctgaaacaca taacaccttc actagttttt agtatacaaa tattgagaaa tagttgttat 180
 taactatctc atccaagaaa tgcagattca tgttgtttct aaatTTTTta tatatatgga 240
 ccaaaatgaa gaaacttaac accatcctag attttagctg ccaaagaat gaaaagaatg 300
 aaaaaaaaaat cttgtgaaaa ccacaagtg atatggatct aatttatggg taaatagata 360
 tagataacaa acagaatacg cctgtttaa actgttaaaa tgacattggg tctaattata 420
 cttttattta aattgaaaga caaggcattt atatggtatc tctaaccatc acaactttgg 480
 tgtgacaaaa agaaattatc accaaaatac acctccttaa gtaagtgtct gatttcacac 540
 ttccagaaaa agtgctcttt ctggtcaagg ccagcaagaa ttgagaaaga ttaagaaagt 600
 gcttcaaaga tgtttattac aaagttgtca taaaaactgt gaagtagatg tagacatcaa 660
 gcataccaaa taaagtaaaa actgtcctcc ggcaaaaaca caacccaaaa aaaaaagcgg 720
 gggggggacc ggggccaaaa cgggtcccgg ggggaatggg tccgccaatc accccaacaa 780
 aaaaaaagg a 791

<210> 64
 <211> 1523
 <212> DNA
 <213> Homo sapien

<400> 64
 gggagatgct gccacctagg ttactttag gaccctatac ggcaacctcc tttgccagga 60
 actatttata aacatcctgc aggaaaatgc agtgaagtag aagagacagg gatatcccag 120
 aaggttatgc aaaacatcaa gagaagatga gaggagtcta tatgtcagaa tacacatttc 180
 ccaccttgcc caacagtaga aaaacataag aagagaaaaa cattaaaaaa tgacaaggaa 240

48

gttaatggaa gtcagcaatg tgatggtggt tggaggtgga gccttcagaa ggtaattaat 300
 gcccttgtaa gaagaggcca gagagcttgc gcaccttctt cctgccatgt gaggagccaa 360
 gaagccggct gtctgcaacc tgcaagagga ccctcactag aagctagcca tactggcatc 420
 ctcatcttgg ctttccaact tccagaactg tgagaagtat atggttggtt ttagtcaatg 480
 gtctatggta atttttttat agcagtccca gccaaagacag tgcctcattt actacatacc 540
 atttatatta ttatataggc tcctttcaga aacctatggt caaataagag ataagatact 600
 gaaacacata acaccttcac tagtttttag tatacaataa ttgagaaata gtttgttatt 660
 aactatctca tccaagaaat gcagattcat gttgtttcta attttttata tataattgac 720
 aaaaagaaga aacttaacac catcctagat tttagctgcc caaagaatga aaagaatgaa 780
 aaaaaaatct ttgaaaacc cacaagtgata tggatctaata ttatgggtaa atagatatag 840
 ataacaaaca gaatagcctt gtttaaaact gttaaaatga cattgggttct aattatactt 900
 ttattttaaat tgaaagacaa ggcatttata tggatctctt aaccatcaca acttttgggt 960
 gacaaaaaga aattatcacc aaaatacacc tccttaagta agtgtctgat ttcacacttc 1020
 cagaaaaagt gctctttctg gtcaagccag caagaattga gaaagattaa gaaagtgctt 1080
 caaagatggt tattaaaaag ttgtcataaa aatgtgaagt agatgtagca tcaagcatac 1140
 caaataaagt aaaactgtca tcaagaagat tcaacagcta tgaaaagagt tcttcaaaat 1200
 atgatatggt tttctagatg ataataaaat ttatcaattc caaatgtcca cattagtctt 1260
 tcataaagac accaatgagt cacaggaaaa aaattaaaaa taaaaaacct ctatctcagg 1320
 gaatcatgct aacaacctga tgtgttttct tccacatatt tatgtctgct tataagtatt 1380
 tacaacataa tattegcata tatgcatttt gaattttttc tgttgctgca cttaaatttt 1440
 tttcataata aaacaagact cctgcaattt gcttttttag gtagactatg tatccctgac 1500
 aaccatccag gtcagcttga tga 1523

<210> 65

<211> 377

<212> DNA

<213> Homo sapien

<400> 65

ggtcgcgcc gaggtacaaa agtgcaaaca aggttagtga ttaacaactt accatcaata 60
 taccacttca acatacttta cattcagcca aatactgaag gtttcaccgt ggaaaaacac 120
 ttttatcact tttaaagtaa cttgactatg ttcaccctga gtgctcttgc ctgagtatgg 180
 caactgatta tgagttcagg ttaagagcaa caccagggaa tacagaaacc cacgttaagt 240

49

tggccattct gacatgaatc tatacttgaa aatgaaaaca atcccaaaga aaacctgtat 300
 gtcaaaaaca gaactgttcc tgcctttcac cccaaaatat ttaaaactaa atctaagcca 360
 cttttaaaat gcatgct 377

<210> 66
 <211> 1703
 <212> DNA
 <213> Homo sapien

<400> 66
 ccaggctgga gtgcagtggg gtgatctcca ctactgcaa cctccacctc ccagcttcaa 60
 gtgattctcc tgcctcaacc ttccaagtag cttggattac aggcgtgccc caccacagct 120
 ggctaataatt tgtattgta gttagagacag ggtttcacca gtggtgtcca ggcttgctga 180
 acttctgacc tcacgtgatc cacctgcctc agcctcccaa agtgctagat tataggcgtg 240
 aacctctgag cccggccagc atgcatttta aaagtggctt agatttagtt ttaaataatt 300
 tggggtgaaa ggcaggaaca gttctgtttt tgaatacag gttttctttg ggattgtttt 360
 cttttcaag tatagattca tgcagaatg gccaaactaa cgtgggttcc tgtattccct 420
 ggtgtgtctc ttaactgaa ctcataatca gttgccatac tgaggcaaga gcaactcagg 480
 tgaacatagt caagttactt taaaagtgat aaaagtgttt ttccatggtg aaaccttcag 540
 tatttggtg aatgtaaagt atgttgaagt ggtatattga tggttaagtg ttaatcacta 600
 acctgtttg cacttttgta caccactgct tgcactagga tcttgggtgtg aattttcaat 660
 tgttttacag tgtatacaga ttattaagga taatttatat aaagatgttt ctgtttaact 720
 ttgtgtgttt tacaacaaag agctataata gatgggttaa cgtttttgaa ttgtgtttat 780
 atgttagttt gattagtatt ttatttttcc ctccctaaca ctcaaattca tggcagggtga 840
 aaagataata gaacataatc aaactaacat ataaacacaa ttcaaaaacg ttaaccatc 900
 tattatagct ctttgttgta aaacacacag agttaaacag aaacatcttt atataaatta 960
 tccttaataa tctagtatac actgtaaac aattgaaaat tcacaccaag atcctagtgc 1020
 aagcagtggg gtacaaaagt gcaacaagg ttagtgatta acaacttacc atcaatatac 1080
 cacttcaaca tactttacat tcagccaaat actgaagggt tcaccatgga aaaacacttt 1140
 tatcactttt aaagtaactt gactatgttc accctgagtg ctcttgcctc agtatggcaa 1200
 ctgattatga gttcaggtta agagcaacac caggaatac agaaaccac gttaagtgg 1260
 ccattctgac atgaatctat acttgaatat gaaaacaatc ccaaagaaa cctgtatgtc 1320
 aaaaacagaa ctgttcctgc ctttcacccc aaaatattta aaactaaatc taagccactt 1380

50

ttaaaatgca tgctggccgg ggcagtggt tcacgcctat aatctagcac tttgggaggc 1440
 tgaggcaggt ggatcacgtg aggtcagaag ttcgacaagc ctggacaaca tgggtgaaacc 1500
 ctgtctctac taacaatata aatattagcc agctgtggtg cgcacgcctg taatccaagc 1560
 tacttggag gttggtgagg cacgagaatc gcttgaacct ggggaagcaga ggttgacgtg 1620
 agtggagatc acaccactgc actccagcct gggtgacaaa gcaagactcc atctcaaaaa 1680
 aaaaaaaaaa aatgagcgg tcg 1703

<210> 67
 <211> 456
 <212> DNA
 <213> Homo sapien

<400> 67
 atctctttaa ataattagca agaagggaga caagatgcag gagttcactt ggctctttga 60
 aaaggaaaac tttaaagtca gtggttgac tgagtcccat gaagccagat cacttctgac 120
 tgcaaggagc ttggaaaagc aagtatctgg atcttttacc agctaaattg ggaggaacta 180
 taaaatgaga aaagattgat gaatattaag tagaagagtg agatgggtcat ctttgcatth 240
 aaaaaagatc atttgctgta gttgtatgga aaatgaattg gagcaggcga tgaggcttcc 300
 tctttgaaga tcacaggtga gaagattagg tgctttctca gaagcccagc aacctgatgg 360
 gagtgtggag tgagcaagac ccaaatcgga gcttcatccc tgcatggttc attttgctta 420
 tttggcaaac ttgcctgca gaatctactc aagctt 456

<210> 68
 <211> 380
 <212> DNA
 <213> Homo sapien

<400> 68
 ccgcccgggc aggtagaggt ttagtgagc cgggatcacg cactgcact ctaggcctgg 60
 gcaacagaga gagagactgt ctaaaaaagg aaaagaaaa aatttatacg ccaaaaaaga 120
 tattctgaga taacctgtag ttaccactaa ctttgtgaca aaattataaa aatccacagc 180
 catctatgaa tctgtaggca gacctgaagt ttgaacgact ggtgaagaca tctgcatttt 240
 ctttatagcc aagttaggat acaaaaaatg caaacaagtc attaataatt actatatgca 300
 agatacagaa acgatgaacg gaaggagtaa gaagttatcc ttcgtggaac tatttaaagc 360
 aaaaatgcaa aataccaggg 380

<210> 69
 <211> 2177

<212> DNA

<213> Homo sapien

<400> 69

ttccaacatc	tcatttctcc	catgaactat	ttggaaaaag	ctgcaggcgt	aatattggat	60
ccctaaatac	tttattctcc	ttataaccatt	atcagaccca	agtatcatct	aatagtcacat	120
aatcaaaactg	cctaaacagt	ttctacactg	tctttttaac	tatttcaaac	tatcaaggtc	180
cgcattttct	tccttagaac	ttttagtctt	tttcttcccc	aaaatatttg	agtccatgcc	240
agttgccttt	agttgtacco	aaataatggt	ttgtctattt	cctaaaagta	gtactcttaa	300
atthaaattt	agtgttatth	ttgttgcatt	cgttccttct	tcctcatgtg	gttgtgcagg	360
cagagcttga	gcatccagat	ttcaaaatta	aaaaataaaa	gataatctag	tttaatatat	420
agtagttgaa	tcaccttaag	tctagactgc	tgtatgagca	cccattatct	ttcactatat	480
tccatcatcc	ccctccccca	tgaactatth	ggaaaaagct	gcaggcgtaa	tattggatcc	540
ctaaataactt	tattctcctt	ataccattat	cagacccaag	tatcatctaa	tagtccataa	600
tcaaaactgcc	taagcagtht	ctacactgtc	tttttaacta	tttcaaacta	tcaaggttcg	660
cattttcttc	cttagaactt	ttagtcttht	tcttccccaa	aatatttgag	tccatgccag	720
ttgcctttag	ttgtacccaa	ataatgtht	gtctattht	taaaagtagt	actcttaaat	780
ttaaatttag	tgthatttht	gthgtcattg	ttccttcttc	ctcatgtggt	tgtgcaggca	840
gagcttgagc	atccagatth	caaaatthaa	aaataaaaaga	taatctagth	taatatatag	900
tagttgaatc	accttaagtc	tagactgctg	tatgagcacc	cattatctth	cactatattc	960
catcatcccc	caacatatcc	acagtagatg	aagggcagth	tgctcaaaca	ttgthttgat	1020
cctgtcatgt	ctgttcagaa	atgctctgtc	atcagaaac	ccacgtctaa	taacaaaatc	1080
ttggactggt	tactatcaaa	acccaacaac	atacagactc	ctcagctagg	ccctagggat	1140
atthttctac	cttgattthc	aatgtthcat	tgaagaatg	cttaattcta	atthggaaaa	1200
aagthtttg	cttcccactt	ctgctttaca	cgthcatctt	tcttgaatc	aatccaatc	1260
caatctatat	tctaagaacc	tgctcaaatc	ttggttcttc	aaagcttht	ctggtattht	1320
gcattthttg	tttgaatagt	tccacgaagg	ataacttctt	actccttct	tcactthtct	1380
gtatcttgca	tatagthaaat	atthaatgact	tgthttgcatt	ttgttatct	aacttggtca	1440
taaagaaaat	cagatgtctt	caccagtcgt	tcaaaactca	ggtctgccta	cagattcata	1500
gatggctgtg	gattthttata	atthttgtcac	aaagthtagtg	gthaaactacag	gthtatctcag	1560
aatatcttht	ttggcgtata	aatthtttht	ttthcttht	ttagacagtc	tctctctctg	1620
tcgcccaggc	tagagtgcag	tggcgtgatc	cggctcact	acaacctctg	cctcctgggt	1680

tcaagagatt cttaggcctc agcctcccga gtagctaggg ttacagggcg gcaccacctc 1740
 catgcccagc tcttttgtat ttttaagtag agacaggggt tcaccatgtt ggtcaggctg 1800
 gtctcgaact tctgacttca ggcaatccgg ccgcctcggc ctcccaaagt gctgggatta 1860
 caggcacaag ccaactgcacc cagccttatt accataaatc atcttgatgc tggtagctga 1920
 taagattcta tttgcttttc tttattcata gagaccacaa acagatcgca gatccagggt 1980
 tctcaaactg gagcatctgc ttaattttcc cataaaatca gtcttattct ttctgacagc 2040
 tctgagactc ctccggccac gactaggtgc tgtcctggag gaaacgggtg aggacggccg 2100
 cacaaaaacc aatctacctg atgaaaactc cgttcccttc tcgccagaaa cataaaatgc 2160
 gatggagacg ctctgtgc 2177

<210> 70
 <211> 226
 <212> DNA
 <213> Homo sapien

<400> 70
 tctcatgccc attcaatatg gaatgttctt cgcttgctga atttaagcct gtattttaag 60
 gttttgtggg tcctcggcca caatgggtga tgtcactgat agaacgaagc tgagtttcta 120
 agggtttggg gctgtgcaag agtaaacact agagcttgag ttgttatcca gctggcaagc 180
 acggaagtct ttgaagaatg taatgtaaaa agggaacaag aatgta 226

<210> 71
 <211> 2554
 <212> DNA
 <213> Homo sapien

<400> 71
 gcgggagagc cctgtcctta aacacattag gacaagtagt taaaacaggg ccaagaagta 60
 tggctgtgta gtgatcactg tacaagcaca cctggctgaa taaaccagtg ggggataaaa 120
 tccagctcac ctgccgctgg ctatgctttg tgcctcagga caaggggtg cttccttgc 180
 aattgacagg aaccatcttc ctgcccaact gcattccac tgcgtaggca ccttatctgc 240
 ccaatggggc tgtgaaccct aattggaagc ttgcaatc ttaacactat atcttcttga 300
 gctgggtttg agtccctatc caatcaagat gaaggcctga gaggactact caagttctaa 360
 catgatgtgg gggcaaggca tagtagtcca gatccgggac atgaggcagc ttttgctta 420
 gtatgacaat ctaatagttc ctaaaataga attatcccag gatggagctc cgtatgacag 480
 aagggtctt cataggtagt tggtaggggg aattgtgtat catgtaagaa gtaggaccag 540

53

atgtctttaa	aaagacctc	caactcta	gctacatgag	tctgtctagt	tgttatgttc	600
caacagggac	agctcttaa	atagtgtggc	aaagcaagag	atgagatttc	cagtgtctgac	660
tcggtggtgg	aatgacttta	gggcaggtat	ttaacctoca	cttccccaag	tacacaagtt	720
atttcacaac	tcttggcaa	aacagtgtctg	taaaaatcgt	aagtttattt	gttaaaaaaa	780
atactgtatt	tgaaaagtac	cttccttctg	ggattttcaa	ataatttgta	cactacattt	840
tattcatcta	cacattggaa	atgagtaa	atgagtaa	atgagtaa	atgagtaa	900
cacaaccagt	gcaaattctc	ctgcctctga	gaaggcagag	aagcccttta	ctcagaaggt	960
cttcaattct	agcattactc	caactcctag	ggaaatttcg	ggtgggtgcc	tatggctgta	1020
tgaccatctg	attcctcagg	gacaggacag	gaattcagca	agggagctta	aaatatttta	1080
agtaattgtc	aacattccat	ggtgactctc	cccaaaaatc	tagtggtagg	aaaataatct	1140
gtacttattc	ctctttctgc	acacaaagcc	ctcatttaaa	tttgtgagcc	tgcttgggat	1200
ccattaccta	gccattcaga	gatcctgtca	aatgcacagc	agattggata	ctcaccatcc	1260
caaaggggtt	cctcccacct	ggatggggcc	aatctctagt	tgacagtgcc	cctcagagtg	1320
caccatggag	atggaatgtc	ccttcagag	agacttttac	acagggaaaa	gcatttggtg	1380
gctgggctcc	aactctcatt	tggtacaaaa	agctttacat	tcttttcctt	ttttacatta	1440
cattcttcaa	agacttccgt	gcttgccagc	tggtatacaa	ctcaagctct	agtgtttact	1500
cttgcacagc	cccaaaccct	tggaactca	gcttcgttct	atcagtgaca	tcaccattg	1560
tgcccgagga	accacaaaac	cttaaaatac	aggcttaaat	tcagcaagcg	aagaacattc	1620
catattgaat	gggcatgaga	tatgcctatc	agattgtgtg	tgtgtgcgcg	ttttttaaag	1680
acagccaatt	acatcgtatc	tagtcaaagt	agcggattct	aaagcagcct	gctgggatgt	1740
tccacttagt	ctaattgctg	tgccactgta	cgccacagca	ccggacagtg	ttctttggga	1800
catctctggg	aaatgctctg	gaacatgctc	cttgatggaa	aacactaatt	tttgaagaa	1860
gtagatgtct	ggaggcaggt	ctggtgaata	aactgaatag	tactgccttg	gaccccagct	1920
gaggggtggc	agtaagcaat	gaggatgggc	tataaagctg	ttaactggct	aagggccatc	1980
cttgggcagg	catttcagac	acatctgtag	agagggcagt	agcatctccg	ataggccagc	2040
tctgaaggaa	gcttaatgct	taatacagtc	acactgcata	aattagctta	gaatgctctc	2100
ttgggtaaaa	aatattaata	gtgtatatgc	acttgaaaag	caaaattcct	caagaaaaaa	2160
agtttaatag	caaggagttt	ccatcagtcc	cggtctttgt	gaggattacc	acaacaaaca	2220
cttaaaagga	tacaacaggt	acttatta	tgctgccttg	ccttttacct	cttccttttt	2280
tttttttttt	tgagatggag	tctcgtctg	ctgcccagcc	tgaagtgcag	tggtgtgatc	2340

tcggctcaact gcaacctccg ccttccaggt ttaggtgatt ctcttgctc ggcctcccga 2400
 gtagctggga tggactacag gcacatgtca ccatgcccag ctaatttttt gtatttttag 2460
 tagagacggg gtttctgtgt tagccaggac ggctcggatc tcctgatttc atgatccgcc 2520
 cgctcggcc tcctaccct cgtgccgaat tctt 2554

<210> 72
 <211> 583
 <212> DNA
 <213> Homo sapien

<400> 72
 cagatcatga agcaattatc ttcctggaag ggtttttagc tatgctctcc agttgcctca 60
 gcagctttgg ctctgatgcc acagtgagcc caaggtggaa ggtgatggaa cagcatcaca 120
 tctgcaggct cagtgtgtcg taaggtaggg gtaaggggag ggcaagtgta gacggatgaa 180
 gaagatttct ccctattgct tccattttga tatttcttta acttcacatt tcatccatca 240
 ttctagaca gttgcctagt tatagaggat ttcttttatc tttttatca gagcatgcc 300
 aggtggaagt gaggtgctg ctggcctaca actccagtgc tcgcattcca aaatgccct 360
 ggatggaggg tggtagatg tcaacacagg tggaaaacag atccgagggc accataccca 420
 tacagacaac ctgtaaaagt cataataaag cccacactg cacggagcta aggcacaaac 480
 aacgcttccc aaccgatggc taagggcca ctaggcggca gatgagcaag ccgaagcatc 540
 accgaaatga agcagctcag aagaggacct aagccccggg aca 583

<210> 73
 <211> 981
 <212> DNA
 <213> Homo sapien

<400> 73
 gaaagaatga gatgttttca gacatttttag gtccctgaga catgttctctg ttcattggcc 60
 agaaactttt tggcgaacca cttcctattc aaaggcttcc tctccactaa taaagtagtc 120
 tgtggtacat gcagccctga ggcttgagat ggaactgcgc aggaagagcc caactgggta 180
 tcagaataag ccacatgcac cttctgaaac tgcccaaatc cacacctgca taagaatttg 240
 agcccagttc ataaagcaga tcatgaagca attatcttcc tggaaggggtt ttagcttgc 300
 tctccagttg cctcagcagc tttggctctg tgccacagtg agccaaggg gaaggtgatg 360
 gaacagcatc acatctgcag gctcagtggt ttggttggtg agggtaaggg gagggaatgt 420
 agacggatga agaaatttct ccctactgct tccattttga tatttcttta acttcacatt 480

55

tcacccatc tccatagcagt tgcctagtta tagaggattt cttttatctt tttttcagag 540
 gcatgccagg tggaagtgag gctgctgctg gcctacaact ccagtgctcg cattccaaaa 600
 tgccccctgga tggagggtgg tgagatgtca ccacaggtgg aaaccagcat cgagggcacc 660
 attccccttca gcaagcctgt aaaagtttat ataatgccca aacctgcacg gcgctaaggc 720
 aaaaacagtc ttoccaaccg tggcctagag ggccttctt aggtgtcaga atgagccaag 780
 cctgaagcac ttcacctgga attgatgtgt aggcttaagg agtatgtgac ccttacagtc 840
 tcactctgta tcaaacacag gataaattgt ttcttcatta aaaaataaaa aaccttcaag 900
 tctacttacc cttctcctgt ccacaataaa gttgagaaaa caaaaaaaaa aaaaaaaaaa 960
 aagatcttta attaagcggc c 981

<210> 74
 <211> 401
 <212> DNA
 <213> Homo sapien

<400> 74
 gccgcccggg caggtaccag gcagagggag gagcaccaag gtgggggata tttaggggac 60
 ctctttcctt caggaccaca cccttctagg tgaaagcacg aaccttgat tactttgcat 120
 tccatctgca aaaacaaatt taggttttga atatggtgaa aaacgaagaa aggaaaatat 180
 aaaactctgt attttatata cagtaaggaa taatggaggc tgataatgat cttgtgatca 240
 gctaagacaa tgtcagtaag caggtgagggt aggggtgctt ctatgggcaa aagggtgaat 300
 atcttgaatg accagaaatg actcgaagag ctgcattact atcatggtag catgcatgaa 360
 gtgatacatc taaacctttg ctaacctaac attattactc t 401

<210> 75
 <211> 1847
 <212> DNA
 <213> Homo sapien

<400> 75
 gccgatcttt tttttttttt ttttttattt ataaatttat tgcctgtttt attataacaa 60
 cattatactg tttatggttt aatacatatg gttcaaaatg tataatacat caagtagtac 120
 agttttaaaa ttttatgctt aaaacaagtt ttgtgtaaaa aatgcagata cattttacat 180
 ggcaaatcaa tttttaagtc atcctaaaga ttgatttttt tttgaaattt aaaaacacat 240
 ttaatttcaa tttctctctt atataacctt tattactata gcatggtttc cactacagtt 300
 taacaatgca gcaaaattcc catttcacgg taaattgggt ttttaagcggc aaggttaaaa 360
 tgctttgagg atcctgaata cacctttgaa cttcaaatga aggttatgggt tgtaattta 420

accctcatgc ataagcagag gcacaagtta gctgcatgtg ctctagactg tagagcgagc 480
 caccgttgag aagcaaagga cagcagcagg aagagcaatg gaacctcctc aggacttacc 540
 aggctgctgc acaggatcta gcttctccca cctaagatgg gcacattgaa agccttgttg 600
 cagcagcacc cccatctgtg gaagcacagg ctgcctgcac ttctccagct gctctagcac 660
 ctgacttctt ggtagtcagg gtaccagggg gagggaggag caccaggggtg ggggatattt 720
 agggggacctc ttctcctcag gaccacaccc ttctaggtga aagcacaac acttgattac 780
 tttgcattcc atctgcaaaa acaaatttag gttttgaata tggtgaaaaa cgaagaaagg 840
 aaaatataaa actctgtatt ttatatacag taaggaataa tggaggetga taatgatctt 900
 gtgatcagct aagacaatgt cagtaagcag gtgaggtagg gtgctttcta tgggcaaaaag 960
 ggcgaatatac ttgaatgacc agaaatgact cgaagagctg cactactatc atggtagcat 1020
 gcatgaagtg atacatctaa acctttgcta acctaacatt attactctca agctttatta 1080
 tctcaaggc ttaaattggct gtagctgttt aatttaaaag caaggcttaa aaaatagagg 1140
 ttactcataa ttccctttcc atatcccttt ttgacttgaa aattatttca ccaactactt 1200
 ttctggaatg ctgcttataa tacatattca cagattgccc tatgtgttat tctagtcatt 1260
 ggcccgtttt gcttataaaa aaggccatgt tttgtattcc taaaaaatct gcagacattg 1320
 ttaacataat acacgtcatt atacatcata tgtatgctac atctactcac tgacatttaa 1380
 aaaatgagct attttcaaag actaacacag gatctgttac tgagacgtgt aggaaggagc 1440
 tcagtgtaaa atattttctt tggatagatc ccttcaaagg gattaaaaca cacaaaatat 1500
 tatttatact aaactttctt aatgttcta tgatatttct atttcaaat tctcttattg 1560
 tgagaatatg tgaatatag atgtagcaaa ttcaacacat aagcttatac cccttagctt 1620
 gagtaaaaga cacatátatg gcttcccagc accaagaaga tggagaagaaac tctactgcaa 1680
 ctacttccct ttttccaagc agctcaaaat gctttagcaa ataccttggtg attctttttt 1740
 tttttttttt ttttgagacg gagtctcgct ctgtcgccca ggccggactg cggactgcag 1800
 tggcgcaatc tcggctcact gcaagccgcc ctogtgccga attctat 1847

<210> 76
 <211> 522
 <212> DNA
 <213> Homo sapien

<400> 76
 attttactct agtattaatg tggttttata aatgattata tgccttatat tctgggggga 60
 aagaaatgtg aaaatgtgct aacgtagaca gaaacagaat atataagtcg ttttgaatgt 120

tatttcctttt ttaaaaaatt tgcttggtgt catatagcca aaactattca tggtgacagt 180
 ttcattgcta tactttttat atgatttcag cgaattgaaa acatgtatat aatagcaaaa 240
 aactggactt catgctgagt atagatgata catataaaag aagtcaaaat ttggagaaaa 300
 aatttaaaaa gataagtaga aaaatgaagt aactgtagaa accatactta ctctttgatc 360
 tcaaatgctc aaaaactgaa tgaaaatgtg aatttaggcc gaccaggtag tcttgtcaat 420
 aaactaaaag caaaaacagg aaaatgaga aatatgttac aactataaca acacaaaaa 480
 gcatagtttt gaaacacttg cagttcttaa atataaaagc tt 522

<210> 77
 <211> 1643
 <212> DNA
 <213> Homo sapien

<400> 77
 actgtcaatc atcaattgac attaacatgg tcaattaagt aatgtttctc acccaacttt 60
 aaatttccat agtcataacc atggaaacat acaaaaaaca aacatgcaaa taaaatgtca 120
 aaataattga gctgagtact ttgcatgctt taggaaataa gatgtagggt ggttctttgt 180
 gccaatatat tcaagtaatt ggtttatctt cccatgtttt gctgctctaa acatgatcta 240
 atataactct cattcatggt gacatagcag agagctgcta ggagtaaacc tgttttctac 300
 acattaatca agctgttctt tcaaagtatt gtttgacaca ttgaatgttt tttattctgg 360
 aatattatca cagcaaaacc tcattaattg gatgctatca aaattatgaa aggaaactgt 420
 agtgagcaca cttgttttga aaagaaattg gtaaatactt ctatgatgca gttttaagtt 480
 atacaattaa ctgctatttg gaatttaata agtccactat aagcaatgtg cctgcacacc 540
 aattaaaggt tggatctgtc tcttcttgac aattttttag aagccattat ttcggtacca 600
 aataaacctg aagttaagaa atatttatat ttacatctat ttatatctgt tggagaatat 660
 ttcataactc agacttggtt gttttacaca gacttctccc cattatocaa catagtgaga 720
 tttttctata gttctatatt ttactctagt attaatgtgg ttttataaat gattatatgc 780
 cttatattct ggggggaaag aaatgtgaaa atgtgctaag tagacagaaa cagaatatat 840
 aagttgtttt gaatgttatt tcttttttaa aaaatttgc tgggtgcata tagccaaaac 900
 tattcatggt gacagtttca ttgcttactt tttatatgat ttcagcgaat tgaaaacatg 960
 tatataatag aaaaaactgg acttcatgct gagtatagat gatacatata aaagaagtca 1020
 aaatttggag aaaaaattta aaaagataag tagaaaaatg aagtaactgt agaaaccata 1080
 cttactcttt gatctcaaat gcccaaaaac tgaatgaaaa tgtgaattta ggccgaccag 1140

gtagtcttgt caataaacta aaagaaaaac aggaaaattg agaaatatgt tacaactata 1200
 acaacacaaa acagcatagt tttgaaacac ttgcagttct taaatataaa agcttttatt 1260
 agttaatttt ttaaaaggat ctcataggat tgacactgaa tcaggttggg aggtggaata 1320
 aggggatggg catattcttt ctgaattact tattataaca tttctagaat cattaggtca 1380
 gtgctacttt gttgtcgtca atgtacaata aaggaatcac aaattgatct tagtgataat 1440
 tttacagagg cagacattgc acataggat gactgcaaaa atgggtggct aactctggga 1500
 agatacttgt gttaaacttt atatgacatt taataaccct tcatcataag gcaatgtttt 1560
 ttacaaaaag atgcacaaa atcatgttag tcatttactc tgcaaaaatg gcacattagt 1620
 gggggttcca aaatccataa tga 1643

<210> 78
 <211> 755
 <212> DNA
 <213> Homo sapien

<400> 78
 cgaggtataa aaactacgtc actctaaaat gttacaaata ggtcatctac ttagtatgca 60
 tagccttgat aaaaacattg gtcaagtcgg gatgtagtgc gccaccaact agaaatgtgt 120
 taagattttt ttaagcagac ttgcttaata aggcaaggag tggggtcagg ttgttctagg 180
 ggccagcaga agggctctaaa atacagggta gtgaaaagag attacgagac tagtgagttt 240
 cctttaaatg cttactagt cattattaag acagccacat ttcagtgggg ctgagccaaa 300
 ctgctgagct tggaaatagca tatgcttggga atctgaatat gaataaggcc caggtgccac 360
 actttacacc acagatcctt tgctaaagag gcaacttttt gtctaacagg caaggaccag 420
 gctggcagtc aggaaggctg ggtttcggtg ctgatcttgt caccaactat gcactcttga 480
 acaagtcact tcacttcact atcctaagcc tgttatctca tctgaacaaa taacaggggt 540
 tagacttagc cttttacaat gacattttgt atatatctac tgagctctaa caattattac 600
 aacatatcta tgtctgacag ataggatagt cctacatatt caggaaactc cacgtatagc 660
 tctcctaaaa ctgattgttg cgtgttacca cacaacacaa caacatacaa acctgggcac 720
 tggcaacacg accggtcaat tctccaaca caacc 755

<210> 79
 <211> 1002
 <212> DNA
 <213> Homo sapien

<400> 79

59

tatttcatct ttatagggaa tttgctccca aggtatattc ggcacgagaa aaaacctcat 60
 atttaaaaac tacgtcactc taaaatgta caaataggtc atcttcttag tatgcatagc 120
 cttgataaaa acattgggtca agtcgggatg tagtcggcca ccaactagaa atgtgttaag 180
 atttttttaa gcgacttgc ttaataaggc aaggagtggg gtcaggttgt tctaggggcc 240
 agcagaaggg tctaaaatac agggtagtga aaagagatta cgagactagt gagtttctct 300
 taaatgctta actagtcatt attaagacag ccacatttca gtggggctga gccaaactgc 360
 tgagcttga atagcatatg cttggaatct gaatatgaat aaggcccagg tgccacactt 420
 tacaccacag atcctttgct aaagaggcac tatttgtcta acaggcaagg accaggctgg 480
 cagtcaggaa ggctgggttt tgggtctgat cttgtcacca actatgcact cttgaacaag 540
 tcacttcact tcactatcct aagcctgttt tctcatctga aaaataaagg ggttagactt 600
 agccttttaa atgacatttt tgtatatttc tactggctat aaaattatta caaatatcta 660
 tgtctgacgg taagatagtc taaatattca ggaaaactcc aagtatagct ctccataaaa 720
 tgatatgttg cgtgttaaaa aaagaaaaaa aagaaaagaa gaagggggag gaaaaataaa 780
 aatgaaaaaa acttcaaaaa tgcacggctg agttggtagc aaagaaggaa attctttgga 840
 ggccaaaaag atctagaaag tttaaatcca atgtgcagga gctggcattg cctagctaata 900
 ccctcatgat tgagaacctc agattataga cactcatggg gaccaagaga taaggcctgg 960
 ggctcaaaa aggccagagc cgaggtcgga tcaagaatc cc 1002

<210> 80
 <211> 374
 <212> DNA
 <213> Homo sapien

<400> 80
 tcttttctaa aactttaatt tccactatgg ctcttttgaa accattttaa tcaagtcaca 60
 tttcttagaa aaaattcact cagggttctg aaggaattag ttattttcta caagcaactc 120
 tgtcatgagt gatagagttg tagctctctt agaagttttt ttcccctttc aaagagaatg 180
 agaaatatgc agagatttcc ttactgactc actaaatgta aagattaaga ggacataata 240
 aaatttggga ctacagtagc atataggttt tcagtttatt tactactaac tagctataac 300
 ttagacaagt catttaacat gctgtgcttt agtttcatct ttgaaaccaa agagattcga 360
 accagaaatc tctt 374

<210> 81
 <211> 399
 <212> DNA

60

<213> Homo sapien

<400> 81

atggggaatt ccattgacac agtcagatat ggcaaagaat cagatttagg ggatgtagt 60
 gaagaacatg gtgaatggaa taaggaaagc tcaataaacg agcaggacaa tagtctgctt 120
 gaacagtatt taacttcagt tcaacagctg gaagatgctg atgagaggac caattttgat 180
 acagagacaa gagatagcaa acttcacatt gcttgtttcc cagtacagtt agatacattg 240
 tctgacggtg cttctgtaga tgagagtcac ggcatatctc ctcctttgca aggtgaaatt 300
 agccagacac aagagaattc taaattaaat gcagaagttc aagggcagca gccagaatgt 360
 gattctacat ttcagctatt gcatgttggt gttactgtg 399

<210> 82

<211> 517

<212> DNA

<213> Homo sapien

<400> 82

gaaagtatat tgacgtaggt agtggagacg ccatgagttc ataactctgc cagagtcgca 60
 gtatgatgta tccggcaccg gacaggtcaa gaaagaacta cttgtttcta ggaagaacat 120
 atgaagtgct taatttataa gcgggctgct gaatattatc caatatagtt tcttctgaaa 180
 agtgaagggg gatcatctat tgtagatta gggggctctg gaaactttt gaaaattcga 240
 atcagtggac caatgtacat gtgaaaacta aagagggcag gggttaaaat agggcttgaa 300
 tttctcattc tgtatagacc agcaaacctc cctgtgcaag gcaagtttac atcacaatc 360
 caagaatgtt tgcacctaata atgctagttt gcttcagccc ctagttaacc tcaggacttg 420
 gtttgcataa aaaaggtaga cagctgatat gttttcatga ataaatattg tcagccagaa 480
 aaggttggtg tcaggtaatg catatTTTTT taagctt 517

<210> 83

<211> 619

<212> DNA

<213> Homo sapien

<400> 83

acacaatgat acccattttt gcatgttaat gtattattaa atatcagtgg gaatagtctg 60
 catgctatTTT cacatctcag gcacacttaa ggaagacctt gtgatgtgca tgttgctcat 120
 ttaatctaga aaggatacca agattcattt agaacttctt tatgcacagt ttttttttga 180
 gtatgttatg tcttgaggca ttaagggtat tactaaagca agcagcggga cttctcagag 240
 aaattaaagg tttcatatca accacacggt gtcaaaatct tcactttgaa taggattaa 300

61

tgatgtttca tcagtattct tggcacacat gacattgttt ttaaataaac agttttatta 360
 ctctgggctg tgacagtttc tcagactttc cttaatatca tacaattctc caatttaaac 420
 tggatatagtc agttttacaa tattttaatt accctgtatt cattagcact ttcctcattt 480
 tctactacct cctccccage tggccctacc ctaggcaatg ccaaactctac tttctgtcta 540
 tatatttgcc tattcttgaa attcaaata aatggaatcg tataatacaa acaaaaaaca 600
 ggaaaaaaaa aaaaaaaag 619

<210> 84
 <211> 646
 <212> DNA
 <213> Homo sapien

<400> 84
 aatgatccat ataggcgaat ggtcatctaa atcatgctcg agcggcgag tgtgatggat 60
 cggcgccggg caggtaactc accccccagg atagagaagt gtttgtagg gagagaagag 120
 ggagaggcag gagccggccc aagcccaggg tccctgcttg ggccccagaa agcacttaac 180
 caggccocaa gccttcaagg gaaaccaagg cctcaaccag acaatcttga ggaaggaaa 240
 agccagactt tgggtttgtt ttttggggga attattggtt tttttttttt tatgtttctt 300
 ttggaatfff gtttggtggc aaattctgtg tgatcttttt tcataaaaaa aaagacaaag 360
 aatttacatt ggacaaaatt aaaaaaaaaa aaaaaaaca acaaaaaaa acaggcgtgg 420
 gcggtctacc tcaggtggcc atatgccggt gtgtcccggg ggtggtgaaa catgtggtgt 480
 tatctccggc ctcaacaaat tctccccac acaattccg tccaccgac caagccgat 540
 ctaacaacag gacatcatat agcaacctat atacgagcac ctcaacagca ccaacgacag 600
 ccaagcgaga cgaacgacca acagacacac cactcacaac caaagc 646

<210> 85
 <211> 419
 <212> DNA
 <213> Homo sapien

<400> 85
 cggcgccggg gcaggtactt tcgttgatac aggcgtggaa gaccttgagt tcccctgtgg 60
 ctaccccatc atagttcctc ctaaggctat accagataag ccatacggag cagatgacca 120
 gcaagaacct ttccagaatt attattctaa ctagaatctt agccaagaga atggaatcac 180
 cacaaatggt atcatgaaaa tcattctcaag taaatttctt attcattca tacggtaag 240
 ttgaggctcg atgatatacg aaaactttaa ctgaattgac ttcataaagg cttaatggtc 300
 ttcaaaatta tgctggttat atgaattcct aaattcaagc tcttttcaa ataataaatg 360

ataaaacaac attttaatta gtatthttacg taaaaatata tattaanaag taaatcaag 419

 <210> 86
 <211> 2133
 <212> DNA
 <213> Homo sapien

 <400> 86
 ggaagtacag gataatatta aagtcaaata gagtacagtt cttcagcadc ataatcaaa 60
 attcaattgc taaaaaatc aaaacttgte agactthttg ctttaataca aatagttgga 120
 atthctgagc aatcaggttt atctthaaat atgtthtttt ctgagcttht ttacttcaaa 180
 aacgatgaga attatcaatt thtcagtact actgacttgt tccttggtga agggaggaaac 240
 attaagtatt taaatcaatt thttaagtct tcgagtatca aatttatttt gthtaatctt 300
 tgatttaatg thtaacatgg gcactthtta tattctctta cctgagttag thttgaattc 360
 ctagaacatg thcattthta cagtggttgt gatattattt agthtaatact actgtctgga 420
 thattthtaa atcttggtac aatttgata aaacaacata acacttgthta acttgccagt 480
 cctctaggaa cttgtthctt thccttactc tgaatagact agtggttagct thccattatc 540
 thttacctta attacgattg thtgaaccac attthaaatc caaatctat attattggtt 600
 taaaagcttc aacttgacaa gatattatta acagtctaca tgaaatctc aaattatata 660
 tgaattthca aacattgata tcagctcctt gatttactth thaatatata thtttactga 720
 aaatactaat taaaatgthg thttatcatt tattattthg aaaagagctt gaatttaaga 780
 attcatataa ccagcataat thtgaagacc attaagcctt tatgaagtca attcagthta 840
 agthttctga thcatctgag cctcaactta acggtatgaa tggaaatagga aatttactthg 900
 agatgattth catgataaca thtgtggtga thccattctc thggctaaga thctagttag 960
 aataataatt ctggaaaggt thctgctggt catctgctcc gtatggctta thctggtatag 1020
 ccttagaggg aactatgatg gggtagccaa aggggaactc aaggtctthc accctgtat 1080
 caacgaaagt actctaattg ctgthttaca tactgggatt atttgtaaga thtcattthga 1140
 aaggaggtht cthtagacca agaaagaaaa ggaaaaaggt tgaaccaat gacctgctcc 1200
 aatctcttag aaactgaatc tcagagaagt taactthcaag gthaaagcat thgttagtgc 1260
 tagaggtaag atgaaaatc aagthtttht atthcttgte thcataataa tataattatt 1320
 gtgatgtctt thgtacaatt tgcataatac tatgtataca thcacatgta gtattthagt 1380
 tacataagtg atgggtacta tgaattact attgatcaag aatgactatt agattthaat 1440
 taagattaca cthtatttht thgthaaaggt gattthaaat gcacattctt thaccaatcta 1500

.atttgaatca tgattagcct cagtttaatt atccttacia aaatattttt gagtgggtgg 1560
gatcagtttt aagttgagct cctagatttg ttgaatagga aaggatacta ataactgttc 1620
taggggaaat gattttgtaa tatttcacct tgaatttttg aactgaacct tataaactag 1680
tcttcagaat gactaagcag gttaaagtgt ttagcattta aatgtcaaat agagaaatca 1740
atctgacttt tggaaaaaag aaagatgttt aatttaaaat atgtaaagca aacttccaaa 1800
tttcttccat cagtaagagt aactaactgt ctgaatgtag ttattattat tgtgtcaagt 1860
taaagatttg tacatacttt cctttacaga tttggataag tgaagacagt aataacattg 1920
aagcagtгаа ccagtggaaa gagacagtaa taaatccaga aaaggttggt atcaggtggc 1980
acaattaaa tccatcttga agacttcaca cattaatttg gtgaagaact tgacattctt 2040
ttagaagact tatgatttca atttgctacc aatgagaaga ggcaaatcaa caaatttgtc 2100
aatttatggg ggctataatt atggtatata atg 2133

<210> 87
<211> 493
<212> DNA
<213> Homo sapien

<400> 87
gcggccgccc ggcaggtctt cgatctcccg gggtgctggg attacaggtg tgagccacag 60
cacctagcct taccttcaaa ttctaaacca agctatttaa atagccactg tttgattatt 120
tgaattaaca tggagcatct tctgggatat tgttcagga aatatgagta gatcaaggta 180
ttttggggat gtaaaccctc atgtttgata aaataaatga tattttgagc tactgtttgc 240
tgggaacaga aagtaagaag ggaaaaggag cgaccataca ggaaagtaa aataataaaa 300
gaaaatttag aaaactagag gaaaaggat gaaaggataa atcctccatc ccatctgat 360
aatggccttt gagcatcact aagccccttt gcttctcca ttaagcaaag gatgatgact 420
gaggaggaac aaacaaaaat agacatcatt ataaaaata cccaagactt ttagatgttt 480
ctctaacatt tgg 493

<210> 88
<211> 1412
<212> DNA
<213> Homo sapien

<400> 88
tgaattagcc atacaaaaaa aataaaaaat tactgttagt caccctacag tgcaaggtaa 60
cactagaatt tatctttcca tctagtaacc actgtttttt aaagagacag agtatctccc 120

64

tgttgcccca gctggagtgc agtggcacia tcatagttca ccacaccctg gaactcctgg 180
 gctaagggat cctccttagc ctcagcctcc caagtagcta ggtatacagg catgtgctac 240
 catgcctggc taattaataaa agattttttt agagatgagg tcttgctgtg ttgccaggc 300
 tggctcaaaa ctctgggct caaacaatcc tcccaccttg gcctcccaaa gtgctgggat 360
 tacagggtg agccacagca cctagcctta ccttcaaatt ctaaaccaag ctatttaaat 420
 agccactggt tgattatttg aattaacatg gagcatcttc tgggatattg ttcagggaaa 480
 tatgagtaga tcaaggtatt ttggggatgt aaaccctcat gtttgataaa ataatgata 540
 ttttgagcta gtgtttgctg ggaacagaaa gtaagaaggg aaaaggagcg accatacagg 600
 aaagtaaaaa taataaaaga aaatttagaa aactagagga aaaggtatga aaggataaat 660
 cctccatccc atactgataa tggccttga gcatcactaa gccctttgc ttctcccatt 720
 aagcaaagga tgatgactga ggaggaacaa acaaaaatag acatcattag aaaaaatacc 780
 caagactttt agatgtttct ctaacatttt ggggtcattt tcagattacc agtgttcatt 840
 tgctgaggta tattaacgga tatttgact taatttga aaatagcagga tccaaaccag 900
 aggtctgtat aagagcaggc ggcatgctg tctggagagc tgctgcctcc acaagtattc 960
 tgacagcact gggctgctag tgagacctg atggccaccc tccccatgtc atggccatgg 1020
 gttttcggga accgtttcct ccttttactg catcacagtt gcaaactcgt ctatttattt 1080
 ttctcttgat taacaactgc actctgacat tgcagcagtg ttgatgaaga caatttaact 1140
 catgtttttg ttaacataat aattgtctgt cgtaactaaa atataagttt cttgaaagct 1200
 ataacaggt atagagaaaa tctttgttat gcacaatacc agggcaggta atatctgtaa 1260
 tatgtattaa cagcaattca ctaaaccatt aatgtctctg tatgctggca cctgtgctaa 1320
 agatttgctg tataaagata aataggaaat tgcctcttct cccacgaaac tcaaaacatt 1380
 tattgaatga ataaataata ggtgaattaa ta 1412

<210> 89
 <211> 624
 <212> DNA
 <213> Homo sapien

<400> 89
 ggtacttgag gtgtttctca ggttccagaa catccgtgtc atcttaccag atccttcaag 60
 gattcagctt aaagatcagc tccaccagga agccttcctg gatttcccct cttagtttcc 120
 aacaagaatc eggctcttcc gttctctgcc caccttgagg tagcagtagc gttcagctgt 180
 gagactctcc gtgttttcc cgttacagtc gtttgtagc gtgcatctc tttcagctga 240

65

attagttaga tgtgagacc taggactctc ttgttttctt cgttacagtc tttgttgctg 300
 catcctctct cactgaattg ttgaattgtg agaccctgtg agggctcggca ccctgtgata 360
 ctggccagga aagggttgtt gcaaggggat catgggattg ttgaatgggt tttgatctgg 420
 attttgatgt tggaaatcaa gttcccaaat gttttcaacc ttgggtaaag gaacatgtaa 480
 tgggtgtttt taacaaaaca aaaaattaaa aaaaaaaaaa aacaaacata aaaaacaacc 540
 aacggctggg ggcacccggg ggcaaagggg gcccgggggg acattgtttt tcccgtataa 600
 atcccaaat tgggaaaaa aagt 624

<210> 90
 <211> 659
 <212> DNA
 <213> Homo sapien

<400> 90
 accacgcctg tagcctctgt ctagagtagt tcacacatgg atgctgtctc tctggactt 60
 ggggtgttct caggttccag aacatccgtt catcttacca gtccttcaag gttcagctta 120
 aagatcagct ccaccaggaa gccttcctgg atttcccctc ttagtttcca acagaatccg 180
 tctcttccgt tctctgcca ccttgagtag cagtagcttc agctgtagac tctcctgttt 240
 ttcccttaca gtctttgttg ctgcatcctc tttcactgaa ttgttgatgt gagaccctag 300
 actctcctgt tttttcgta cagtctttgt tgctgcatcc tctctcactg aattgttgaa 360
 ttgtgagacc ctgtgagggt cggcacctg tgatactggc caggaaaggg ttgttgcaag 420
 gggatcatgg gattgttgaa tgggttttga tctggathtt gatgttgaa atcaagtcc 480
 caaatgtttt caacctggg taaaggaaca tgtaatggtg ttttttaaca aaacaaaaa 540
 ttaaaaaaa aaaaaaaca acataaaaa caaccaacgg ctgggggcac cggggggcaa 600
 agggggcccg gggggacatt gtttttccc gtaaaatccc caaattggga aaaaaaagt 659

<210> 91
 <211> 556
 <212> DNA
 <213> Homo sapien

<400> 91
 aatthttcaac tggcccatatc tttatagtga tggaaagcgc ataacactac ttgtaaatca 60
 ttaaaatagg gtgataactg tgataatagt gtttcttgca ttctagaaaa ttattttatt 120
 aactacattc aaaaccagc atttcacagg ttccatcatt agaaacagta tagttctagt 180
 taacatgatt ggagagtthc aggggaaagg tttacattht ctgaaactgt atttggatg 240
 tgactcaatg tggatthtca gtcttgtag tcacttacat gactgacgth tgcaaggatt 300

66

tattgccaag taaaatttga tcagagtgca ctgagaatag ctacataagg ggaaatctct 360
 caaaattcct tctgttctact ttaattcgga gcatatgttt caactcattt tcacacatct 420
 gtcccacagt tgaagcatta acacacatct tcacgacaca atgaacacat acacattagc 480
 aaacataagt ctcttaatgc aaaattacta gttgactaca atatagctac cttaaaagca 540
 gagcttgcta taattc 556

<210> 92
 <211> 635
 <212> DNA
 <213> Homo sapien

<400> 92
 acaaaatata atgttataaa tgtattttaa aaaaagaaat acaaattcta tggctctttg 60
 cattttactg cctcaaagca gaattagcaa agctgatgaa gaatgaacat tttcccttgg 120
 gcgggtggcc cttggctcact cccacaggca cgttaccggg ctccggcgtg tgctcccacc 180
 aaccacggca aacaaaggcg tcctcctcac ttgaagtccct ggctgtggt tgtttcatct 240
 gtttttttgc tcagtgaaca aaacgttctg aaattagaac tcaccaaagt taaaagcagt 300
 aaaacaacat atgctactta agacattttg aagcggaaag taaagctatg tgaatgccgt 360
 ccttccttcc ttcctttttc tacagcttgg aaacctctga gaatttgctg gcgggtggca 420
 gaggagggct tcgtctagct cttgaacgga caggaactgt ctggctagac agctctccag 480
 accacgaaag cccaggaggt gccctcttcc acacaacaga ctaagcactg caccacttt 540
 ctttgatcca gaaagcatcc ctactgacct tgtaacctac accctctctg tccaaagaac 600
 agaggccgac cagagtagcc agcctggaga ggcac 635

<210> 93
 <211> 8156
 <212> DNA
 <213> Homo sapien

<400> 93
 cggggcgtgc gcgtcctcct cccacaggccc gccgcctccc tgccaagaat ctgagagagg 60
 ccgagtggag ttcggctcctt ctctgaacag ttttagctga gagtaccagc atccaactgg 120
 gagcgttgtc attgcatttc cacattccca ggaaagccca ggtgctggct gccagctgct 180
 gcgcccccca tgtagaaggc gcacctcctg ggagcaggca cgtcttttgg ctcttctgac 240
 catggagaga taggacggtc cctgcagccc gcgacacaga aagctgtgcc gccaccaccg 300
 gccgcgtccg tccttcggat ggatcgcaac agagaggccg agatggagct gaggcgaggc 360

67

cccagcccca	ccagggccgg	ccggggccac	gaggtggatg	gggacaaggc	tacctgccac	420
acctgctgca	tctgcgcaa	gagcttcccc	ttccagagct	cgctttcgca	gcacatgcg	480
aagcacacgg	gcgagaagcc	ctacaagtgt	ccctactgcg	accaccgggc	ttcccagaag	540
ggcaacctga	agattcacat	ccggagccac	cgcacgggga	ctctgattca	gggacacgag	600
ccggaggcgg	gcgaggcgcc	gctgggtgag	atgcgcgcct	ccgagggcct	ggacgcctgc	660
gccagcccca	ccaagagcgc	ctctgcctgc	aaccggctgc	tgaacggggc	ctcgcaggcc	720
gacggcgcca	gggtcctgaa	cggggcctcg	caggccgaca	gcggcagagt	cctgctcggg	780
agcagcaaga	agggggcaga	ggggtccgca	tgcgccccgg	gggaggccaa	ggcagcggtc	840
cagtgtcct	tctgcaagag	ccagttcgag	cgtaagaagg	acctggagct	gcacgtgcac	900
caggcgcaca	agccgttcaa	gtgcaggctg	tgcagctacg	cgacgctgcg	ggaggagtgc	960
ctgctgagcc	acatcgagag	ggaccacatc	accgcgcagg	ggcccggcag	cggcgaggcc	1020
tgcgtggaga	acggcaagcc	cgagctgagc	cccggggagt	tcccgtgcca	gggtgtgggc	1080
caggccttca	gccagacctg	gttcctgaag	gcgcacatga	agaagcaccg	gggctccttc	1140
gaccacggct	gccacatctg	cggccgtagg	ttcaaggagc	cctggttcct	caagaaccac	1200
atgaaggcgc	acggcccca	gacgggcagc	aagaacaggc	ccaagagtga	gctggacccc	1260
atcgccacca	tcaacaacgt	ggtccaggag	gaggtgatcg	tgcgcggcct	gagcctctac	1320
gaggtctgcg	ccaagtgcgg	gaacctgttt	acaaacctgg	acagcttgaa	cgcccacaat	1380
gccatccacc	gcagagtcca	ggccagccgc	acgcgcgccc	cggccgagga	ggggcgggag	1440
gggcccctcg	acaccaagca	gttctttctc	cagtgcctga	acctgaggcc	gtcggcggcc	1500
ggcgactcgt	gccctggcac	gcaggccgga	cggcggggtg	ctgagctgga	cccggccaac	1560
agctaccagg	cctggcagct	ggccaacgcg	ggtaagggtg	ccgagccggc	cgagtacctc	1620
aagtacgggg	cctgggacga	ggcgtggcc	ggggacgtgg	ccttcgacaa	ggacaggcgc	1680
gagtacgtcc	tggtagcca	ggagaagcgc	aagcgtgagc	aggatgcacc	agccgcgcag	1740
gggccccgc	ggaagcgcgc	gagcgggcct	ggggaccccg	cgcccgcgg	ccacctcgat	1800
ccccgctcgg	ccgcgcgcc	caaccgcagg	gccgcagcca	ccaccggcca	gggcaagtcc	1860
tccgagtgct	tcgagtgcgg	caagatcttc	cgcacctatc	atcagatggt	gctgcactca	1920
cgcgtgcatc	gccgcgcgcg	ccgcgagagg	gacagtgacg	gggacagggc	ggcgcgggcc	1980
cgctgcggat	cactcagtga	gggtgactcg	gcctcccagc	ccagcagccc	tggctccgcc	2040
tgtgccgctg	ctgactcccc	gggctctggc	ctggccgacg	aggctgccga	agacagtgg	2100
gaggagggcg	cccctgaacc	tgcaccaggg	ggacagccgc	gccgctgctg	cttttccgaa	2160

gaggtgactt cgaccgagct ctccagtgga gaccagagtc acaagatggg agataacgcc 2220
tcggaagag acaccggcga gtccaaggca gggatcgag cttctgtgtc cataactgaa 2280
aacagtagca gagagacttc tagaaggcaa gaggcagaca gattttctat ggacttaaag 2340
atgccagcat ttcaccccaa gcaggaggtg cccgtccctg gtgatgggtg ggagttccct 2400
tccagtacgg gagcggaggg ccagacgggt caccctcgag aaaagctgtc cgatttgcac 2460
aacaaggaac actctggggg agggaagcgg gcgctggccc cagacctcat gccgctagat 2520
ttaagtgcga ggtcgacgcg ggatgacccc agcaataagg agacggcctc ctccctcgag 2580
gcgcttttag tcggtcaccg gtgtccttac tgcagccaca agacctacta ccccgaggtc 2640
ctgtggatgc aaaaacgcat ctggcaccgt gtcagctgca actccgtggc tccccgtgg 2700
attcagccca atggttacaa aagcatcaga agcaatttgg ttttctcttc cgggagcggg 2760
cgcacgggcc ccccgctgc cctcgggtggc aaagaatgcc agccttgct ccttgcctcg 2820
ttcaccgcga ctcaggtgcc aggggggatg cgggggtcca aaagtggctc ttctccctg 2880
ggagtggca caaaagccgc tagcatgcct aagaataagg agagccattc cggaggtccc 2940
tgcgctctgt gggcgcccgg ccctgacggg tatcgacaga ccaaaccttg tcacggccag 3000
gagccacatg gcgcccgcac acaggggccc ctggccaagc ccaggcagga ggctagctcc 3060
aaaccggtgc ctgccccggg tggcgggggc ttcagcagga gcgccacccc tacgcccacc 3120
gtcatcgccc gggctggcgc gcagccctcg gccaatagca agcctgtgga gaagtttggg 3180
gtccccccag cgggggctgg ctttgcccc acaataaagc acagtgcccc ggactccctg 3240
aaagccaaat tcagtgtca gcctcagggt ccacctctg caaagggcga agggggcgct 3300
cctcctctac ctccccgcga gccccctcg aaggcagccc aggagctgag gactctggcc 3360
acctgtctg cggggtccag gggcgacgcg gccttgacag cccagcccgg cgtggctggg 3420
gcgccccccc gtctactact ccatcaaaca ggagccagtg gccgaggggc atgagaagcg 3480
cctggacatc ctcaacatct ttaagacgta cattccaag gactttgcga ccctctacca 3540
gggatggggg gtcagcggcc ctgggttggg gcacagaggg aactccgga cgcaggcccc 3600
gccaggagag ttcgtctgca tcgagtgcgg aaagagcttc caccagcccg gccacctcag 3660
ggcccacatg cgggcacact cagtgggtgt tgagtccgat gggcctcggg gttctgaagt 3720
tcataccacc tccgcagacg cccccaaaca agggagagac cattctaaca caggtaccgt 3780
ccagacagtg cctctgagaa agggaacctc aaggcgtgtt tccgacgcac cccaggtccc 3840
cgtaacggcc attagcagta ccctcacgat gtcccagcag cctcccacct gtgacctggc 3900

cgctccatgg aagaacagcc ggggaactcc tgagcagaca cctcacatcc cgagccgctg 3960
cgctggagtg gaaactgaag gcagatgcct ctccttgta aacgttcaga aataaatgaa 4020
gatgctatat tctagaaata catgtagata ctatatacgc atttacgtgc tcatcgtcca 4080
tagtcccata ttttcttata ataaacagta gtactggcag gcacagtagg ggcaacaaggc 4140
atctgtctta ttcaagacaa gtttgagaca ctggaaaaaa agataactgt tgtgtgtgtt 4200
ggacagagtg gcgaggctga gcactgtcac aggggcctcc catgttaaga gggactgtgg 4260
ggatgatgtc agaacaagac gtggtggatt tgaggttgat cgagtattaa tactactgcc 4320
tctccttgtc ttagtgggta tttaaaatag taaataagag agaggaagga ggtgacgttc 4380
agggtcgtg ggaagcaggc ttggcggagg ggtatgatga tgagaccctc attgttctact 4440
ggctccatcg cactcctccc tggggccgtg tgectgttcc attcttcca ccattcgaac 4500
tgagcgaatc tggcaaagga gacacgtctg tgggaatgcg tagattccgc ctcggaagag 4560
agctagcgca aactaagaa aagcaggcct cttgtttatt ctcaggacct ttttgtaaca 4620
gggctacatt ctgcaaactg cttacaaagg aagactatac gtcttaacaa attatctagc 4680
cactgagtcc tcccgattcg gacctgttt agtaatggca gaagaatccc tgagcaggtt 4740
cagggtccct agatgactag ggtgctgagc tctggcgcct tctgtcccca ctctttgctt 4800
ccccgccct tccctgagcc accccagcaa gtgggtgtct tttctccctg ggcctgggta 4860
cctccacagg atgagtgact ttgttcataa aggggtggga tcaccagccc cttgggtggg 4920
ggacgggttc atatacctct tcctcagtaa tgcaaatgcg agtttttggtg gtgggggtta 4980
aggcccataa caaaggatct taaaccatgc agtgtacgca attgaaatgg tattccacag 5040
atataaatat tttcttttcc cattgccgtg acactatgtg tgatggtaat atttctgaga 5100
gtttcagatt tttgcacata tgattttatg cattatcaaa agttactgct gccttgaatg 5160
aaaatgttct gtgaaatttt ttgcaaaagc tttactaggt ttttttttaa ttgtgaaatt 5220
ttgtaaaggc aggaaatgga ttaaaacgag catgctaat atatttttca aaaaagcaat 5280
aattttcat gtacagaaat taccetaacc ttaatactg gcgagagcaa cagtttactt 5340
aatcggtaa tggactagtg cagttttgt agacagtggg cttctgatac aaagtctgt 5400
ttaaacacag acacacacac acacaacac acacacacac ctaaagtgt gggtttctctg 5460
ttctaagat ttgttgaata ttattatatt attattatta ttattattat tattgttatt 5520
gttattagta atgtttggtt ctggattcta cttgttactg agtttaaatt acttgacgggt 5580
tcaggttact ttgcaacact ttcaaacgat gcaatgtaac tggctagctt atatatatat 5640
atatatatat atatatatat atttttttt ttttttactt attttttct gatattctta 5700

caccagatat gtacgaaaat gatctgtcct gttggtgtaa ttaggaatgt ccatgcagat 5760
acagttaaac aactgtaatt gactgttctg taaagttatt ttgggcaaag ttgcgagac 5820
acattcctct gtccacctaa gaaatcagaa gactcttctg ttgatttatg tttaatcatt 5880
tcagtagttt cccacagtg atcatttctg cattttctgg cttttgtttt cttggctgaa 5940
agtgaatggt gactgttagg aatgtcaggg actagtgacc cagtctctgtt tctctgtggt 6000
ttagttatta aaaagaaatt ctgtacccaa agtgacacga aagtgtagtc tacattttta 6060
ctgtttcaga agcggcatgg aaaagtgcag ttggcctttg gagctggaag tgtcttgctg 6120
gtgaggctcc atcctggagg ctctgggtgg gagtgggctg gcgctggggc cctgcccggc 6180
gcgtgctgga tccttcctgg cttgcaggag agcaggcgtg gaggacagtc agcttgccgg 6240
gccgcgcagg gtgcacagag tgcaggagga aggttttcac ccagttaaac agactgggga 6300
gccccccaa gaacgccatc ccttgaggcc agctgtgggc aggcctggat gtgtggtccc 6360
ttccttcac ccacgctcag tattgtgttg gtttgttaat ttgttgattt ggtcatagta 6420
tttaatatga tttgtgttc cttatttatt tagccaccgt tttgattgcc ttttttttc 6480
cgaatggtaa tttctgcatg atacacttct gtacgttctc ttctgactgt tacagacttt 6540
ctactacctc tccgatctg ctgtttcctt gtttctaac aggatttttt acagtgttc 6600
gtctaatagt acttagacaa taaagggttt ggttgbctac actgcagctt ctcggtgtct 6660
ctcccctgct tccgctcgc tgcctcccgc tctgcccctg ctggggcctg gctgcacct 6720
ggcctgcctt cctatactct cctgtttccc gctcatatct ctctctcatt tttgcttca 6780
aataacacac agctaatagt cttctaaaaa tcttttcagg ttgttcactt gtattcctta 6840
atltgaagaa tgaatattta aattctctca aaagtcagat attgaggatc ttctctggga 6900
aattggccac tgtacctgcc caccttctg cctggttccc tgggaaggctt tattgtcatc 6960
ttagacggac agatttcatt ctacgacca tacagatttg gcttcaaagc caggtgaatt 7020
ttgccttga ggctctgaaa agtattaagt gttttaagag gtcccccaat atttacttat 7080
ttatttttta aaccaagaaa gacactggtt ccctgaaaag caggtgcttc aggaagtagc 7140
aattgggagt tgcatacagt tacttcgtca gagaaaggag cgccagtat gacaggcccc 7200
cacccctgat cggccactg tgcacaggtc gctgaggggtg tgagaacacc tctgcagggg 7260
ctccggcaca tgtgggtttc atcgtctcac actcctcag gctgcagggg ttgagtgag 7320
aaagggcaag cttcatctcc atgggtcctc tccaggtgga caactctggt cggcctctgt 7380
tctttggaca gagagggtgt aggttacagg gtcagtaggg atgcttctg gatcaagaa 7440

71

agtgggtgca gtgcttagtc tgttgtgtgg aagagggcac ctcctgggct ttcgtggctg 7500
 gagactgtct accaacattc cttccttcaa actagacaaa ccctcctctg ccacccccag 7560
 caaattctca gaggtttcca agctgtagaa aaaggaagga aggaaggacg gcattcacat 7620
 agctttactt tccgcttcaa aatgtcttaa gtagcatatg ttgttttact gcttttaact 7680
 ttggtgagtt ctaatttcag aacgttttgt tcaactgagca aaaaaacaga tgaacaacc 7740
 acagggcagg acttcaagtg aggaggacgc ctttgtttgc cgtgggttggg gggagcacac 7800
 gccggagccc ggtaacgtgc ctgtgggagt gaccaagggc cacccgcca agggaaaatg 7860
 ttcattcttc atcagcttgg ctaattctgc tttgaggcag taaaatgcaa aagaccatag 7920
 aatttgatt tcttttttta aaatacaatt tataacatta tattttgtac tctttatatt 7980
 agaatttgta actagattga tgtatttaac tattttctgaa aaagtaattc aatgttttag 8040
 ttgtgtgata aaaatattta gataaaacat attcattcta ttggaatttg aaataaataa 8100
 aaacatcttg gagttctgaa aaaaaaaaa aaaaaaagat ctttaattaa gcggca 8156

<210> 94
 <211> 668
 <212> DNA
 <213> Homo sapien

<400> 94
 tggtcgcggc cgaggtatcc cttagaaatt gacagcttct atgaaattta caatcaagaa 60
 ggcataagaa caactgctgc agcttcagaa ctatgcagaa aataaaatgt caacagctgg 120
 gagaaaaaat tctcagtgag caacagagcc agtgaaaaat atcaccagta gagcaggcac 180
 ctgagacagg gaaatggcac ccactaagtg caggtctaca ggggctgacc ttgcagacca 240
 tttatggatc ccaagttacc aaaaccctta tttgattatg aactgcatag tagataccaa 300
 tacatcagag cattgtctca atgttaatat ctatctgtgc tgattgtatt gtggtcatgt 360
 aacagaatga atgcccttgt tcttaagtga tacgtgggaa aatatttgag gggatgaagtg 420
 tcatgatgtt tgctacttac tctcacatgc tttggcaaaa ataaaacatc tctctctctc 480
 caggcaaaat atagtaatca gtaaaatatt aacaatctgt aaaaccacac cacaacccaa 540
 caaaaaaggt tgggggacaa ccaagggcaa aagggtgttc cgggggtgaa atttgttttc 600
 gggccaaaat tccccacat ctccgcaca aagcgggagc aaaaaaaccc acaaaaaaca 660
 cacatata 668

<210> 95
 <211> 746
 <212> DNA

<213> Homo sapien

<400> 95

gactaagaca cctttctaga cagagaggag gccgatggca gacattctca gataggtttg 60
tagctattga cctggctgca tcaaaggaga tgaaatccct tagaaattga cagcttctat 120
gaaatttaca atcaagaagg cataagaaca actgctgcag cttcagaact atgcagaaaa 180
taaaatgtca acagctggga gaaaaaattc tcagtgcagca acagagccag tgaaaaatat 240
caccagtaga gcaggcacct gagacagga aatggcaccc actaagtgcg ggtctacagg 300
ggctgacctt gcagaccatt tatggatccc aagttacca aacccttatt tgattatgaa 360
ctgcatagta gataccaata catcagagca ttgtctcaat gttaatatct atctgtgctg 420
attgtattgt ggtcatgtaa cagaatgaat gcccttgttc ttaagtgata cgtgggaaaa 480
tatttgaggg gtgaagtgtc atgatgtttg ctacttactc tcacatgctt tggcaaaaat 540
aaaacatctc tctctctcca ggcaaaatat agtaatcagt aaaatattaa caatctgtaa 600
aaccacacca caaccaaaca aaaaagggtg ggggacaacc aagggcaaaa ggggtgtccc 660
ggggtgaaat ttgttttcgg gccaaaattc ccccatctct cccgcacaaa gcgggagcaa 720
aaaaaacccac aaaaaacaca cataca 746

<210> 96

<211> 978

<212> DNA

<213> Homo sapien

<400> 96

cgcccgaggt accctgtgcc aactagggga ggagaactgt tgcctgctaa gttggtggca 60
ggaaccagct tcctgaagga tataaattcg ctggaagaaa gaccttggtt tatgttccag 120
tgctgttttc ccatctctag agcagtggct ggcagacatg tggctactca agaatggttc 180
ccagatgaat gaatgcagga aacagttccc tcctccaatg agattaacag ctgatccatg 240
cttataatga ctgaactctg taaagagggg aaccctctgc caatggggga tcaaaatggt 300
taatgagagc cctgctgtgg agagaggtag gacagcaagc acagaagcac ctgaccccat 360
gcagaggacg ggaggcagaa gcaggggcca gactggaggg agtgatctcc atatgcccat 420
atagcatcca tctgtcttgc caagcaccct tctggggtcc tctaccttta ccagagcata 480
gtctcttctg cagatcaatg aaacgaagcg aagctggata gactatggcg agagcacttc 540
cctctctgcc tcccccaaga tgaatggcta cttgtggaga ggagctgtgg catataaaca 600
ctttcatcct tacccaaggc ccccatctc cctcaaatgg cataacaggc agacgagcgg 660
ccaaccaact ccctctgatt ctcatacaca ggggtgtggc ttccattggc atctttgagt 720

```

ggccccaagt cgtacgcaga ttgtggactc agtacacagc ttgcgcaaac ctggacaatg      780
tggccatggc ccatcataca cctctacacc acctcatagc gacgttgaat atgagatcca      840
cccgtagtgc ccagctcata acatcctctc cattaagatt gaccacaggc aacttaccat      900
tgactaggac caagtcccc aaacacaaa attgagaaca gagcaacatg gtgccaalaa      960
tatcacagag aatcaac                                                    978

```

```

<210> 97
<211> 787
<212> DNA
<213> Homo sapien

```

```

<400> 97
acctggcaca aagcaaacia taaatattat tgttattggt gttataattg taaaatgaat      60
gacttcaaaa acatagtccc agtttgaggg gattttgtga tgcagaatat ctaagtcata      120
gaaatagaag acaggtggaa taagtatatg ttcagagttt ttagatgtgt tgagtagaga      180
cggtataaat ggaagcatta aatacaaatg aaaatcacac cagatatccc tgaaattcaa      240
gcaaagaaag ttcacatcatg attcttgggc agcaagagaa aggactaggg ttatggcaat      300
gtgtggaaaa gttgaggctt gctaagggtt gagatctggt ggtagccctg gatcacatgg      360
ggtcagcacc aggcagtgcc tctgaaagcg gagagaggtc ctggacttcc cttgtgtata      420
acagttccta gtgtccaaca atgaggaaac ggtgaagcat ggttacaaaa ctgtgacaaa      480
acatatttac atctagcact gttaccactc acctagccaa acattggctg cacacgtgca      540
gcccttattt gtaattaaca tcaaaagact agatctgaag ccttcataa atgagagacc      600
attcatatgg cattcctgga acaaaacact gcacaggtac caaggctctc cactccctga      660
cgggttggtg ctgaacagtc agggattgtc ttgactagac ttctgatgct tctgcatctt      720
cttctctctt cccggaatc caaataacca attcatacca ttgtatttat gcttcgggta      780
acctagt                                                    787

```

```

<210> 98
<211> 3670
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (3416)..(3416)
<223> a, c, g or t

```

```

<400> 98

```

74

agcggacagc	cgctccctcg	ctctgctggg	gcctccggac	gcgcttccca	cgcggtctc	60
tggaacactc	ggtccgaacg	cacgcctgct	tgcactcaca	ctgcggttca	cacccggagg	120
cgctctcgca	ctcacactgc	cgctcacgcg	cgctcacact	ccccacgcg	cgctccgctc	180
eggctccagc	cccgcgcca	gcgaaggcgc	aggcactgct	gccgagagcg	ccgaggggcc	240
ccgcggcctt	cccatggcgg	acctgagctt	catcgaagat	accgtcgcct	tccccgagaa	300
ggaagaggat	gaggaggaag	aagaggaggg	tgtggagtgg	ggctacgagg	aaggtgttga	360
gtggggtctg	gtgtttcctg	atgctaattg	ggaataccag	tctcctatta	acctaaactc	420
aagagaggct	aggatgacc	cctcgctggt	ggatgtccgc	ctctcccaa	attatgtggt	480
gtgccgagac	tgtgaagtca	ccaatgatgg	acataccatt	caggttatcc	tgaagtcaaa	540
atcagttctt	tcgggaggac	cattgcctca	agggcatgaa	tttgaactgt	acgaagtgag	600
atctcactgg	ggaagagaaa	accagcgtgg	ttctgagcac	acggtaatt	tcaaagcttt	660
tcccatggag	ctccatctga	tccactggaa	ctccactctg	tttggcagca	ttgatgaggc	720
tgtggggaag	ccgcacggaa	tcgccatcat	tgctctgttt	gttcagatag	gaaaggaaca	780
tgttggtctg	aaggctgtga	ctgaaatcct	ccaagatatt	cagtataagg	ggaagtccaa	840
aacaatacct	tgctttaatc	ctaacacttt	attaccagac	cctctgctgc	gggattactg	900
gggtgatgaa	ggctctctca	ccatcccacc	ttgcagtcaa	ggtgtcacct	ggatattatt	960
ccgataccct	ttaactatat	cccagctaca	gatagaagaa	tttccaaggc	tgaggacaca	1020
tgtaaggggg	gcagaacttg	tggaggctg	tgatgggatt	ttgggagaca	actttcggcc	1080
cactcagcct	cttagtgaca	gagtcattag	agctgcattt	cagtagccaa	agaggacagg	1140
aacaagtctg	tcttcatgag	ggaggaagac	aatggtccta	taatgccctt	ggataagaaa	1200
aggaactttt	tgagctgcac	cttcagttta	tctcaaacg	ctgcgttgtt	tgtcttcac	1260
taatccagct	ttgatggaca	tctgtgatgg	ttgcctgtac	acttgctgaa	atgaaatatt	1320
agaaatggct	gtatattcca	aagaaaccct	atattatata	tccacattac	tgctgctagg	1380
attcatagtt	gcacatactg	tttattgctt	atgtgtagaa	ggaatgaaac	tagtttccag	1440
agttgttatt	aatatgaata	tatatcatgt	gttaatattg	agaaaggaaa	aatacattcc	1500
cgggtgttagt	agttcttcat	ttcctgtctc	caacagaaaa	ttcaactcatt	ttagaactag	1560
tgtaattctt	gataataaaa	taagagtttt	gattaagaac	agcatagagc	ttcaaaatgc	1620
aaagtgaatg	attagtaaaa	ttatgtctca	ttttattttt	tcagcaccca	taccacaatt	1680
aatattaggc	tggattgcca	tgggaaacat	tttttggcat	taatgcagca	acataatact	1740
cactttaggt	attactacat	agttgaagga	tttaactgaa	tgtatggatc	aaatttattt	1800

atttgacata ttcgaagctg tggtttaata ggaatttgag aaaggtgtaa gaaataggat 1860
aaaaagaagg tcagcaccat gtaccaggaa tagctttact ttccatacat agaaatataa 1920
athtagtggt atcctatatt actttagtggt cgtacgcttt gtaagactta aatattttat 1980
tctattgatt ccactacttt ggtatgtaa gacatttctt taaagatgac caacaatatc 2040
cttatttttag gtgccactag cagatgtaag cgtatactta gttgccgta gatgtgacag 2100
aatgagataa tttatgtaa gcagtagagt acctggcaca aagcaaaca taaatattat 2160
tgttattggt gttataattg taaaatgaat gacttcaaaa acatagtccc agtttgaggg 2220
gatttgatgc gcagaatatac taagtcatag aaatagaaga caggtggaat aagtatatgt 2280
tcagagtttt tagatgtggt gagtagagac ggtaataatg gaagcattaa atacaaatga 2340
aaatcacacc agatatccct gaaattcaag caaagaaagt tcatcatgta ttcttgggca 2400
gcaagagaaa ggactagggg tatggcaatg tgtgaaaag ttgaggcttg ctaaggggtg 2460
agatctgttg gtagccctgg atcacatggg gtcagacca ggtagtcct ctgaaagcgg 2520
agagaggtcc tggacttccc ttgtgtataa cagttcctag tgtccaaca tgaggaaacg 2580
gtgaagcatg gttacaaaac tgtgacaaaa atatttacct ctagcactgt taccactcac 2640
atgcaaaaca ttggctgcac acgtgcagcc ttatttgtaa ttaacatcaa aagactagat 2700
ctgaagcctt ccataaatga gaggccattc atatggcatt cctggaaca aacactgcac 2760
aggtaccagc ctctccactc ctgaccgggt tgggtgctgaa cagtcagggg ttgttcttga 2820
actagacttc tgatgcttct tgcaatcttc tttcatcttt cctgaaata cacaaaataa 2880
acaaatacaa taacaaatag taattaaatg actttcagga taacatctag ttgttcagac 2940
ttcacccttc acaggtgtgt gtgtatgtgt gtttatgtct gtatattgaa gcaatttgaa 3000
tttatttact gtatatttct tgagtaaaag actgaaatga actacttggg tcagatcatg 3060
gtgtccattg gtgacattgt ttggaggcat aatattcttt atatggaaaa tcctttaatt 3120
ccacagttag ttacctcaga ttcagaatat gaatactggt tataatacgc ttttgtagga 3180
atgaattcga aaggtagttg tcagtaaaca aaagcacaac aaactaatct cagagtctgc 3240
cctgatggct gtgatagggg cagaaagcta aaccctactg ctgacgcgcc ccgcacattg 3300
ggcgcagaat ttccaagaa aacggggcaa atcaccgcca cggtcctaac tctgaactct 3360
atacgggcca tctcgcctaa accactacaa ggcacgcacg ggaaaggact ctccgntcgc 3420
gactcgcaag cctacggccc ccgaacgaca ggcgcaaccac gacaccaccg gcgcgtctac 3480
gagacatgat cagcgtcaag ggcacctgaa aaaacgatgc cccaactagt gcggccccga 3540

76

accaggcaga cactaagctt gatagcacag cgactgcacc aagagctaata caccgacaca 3600
 accaaagaca gaaactaccc actctatcac tacacggacg aactagaaa caacctgcaa 3660
 ttgttactgc 3670

<210> 99
 <211> 938
 <212> DNA
 <213> Homo sapien

<400> 99
 ccacccccga cgacgacata ttaggggaac gggccactag atggctggtc gagcggcgca 60
 gtgtgatgga tgcccgggca ggtacataat gttcagacct cctccatcct tttaaagcc 120
 tgctgcagta aataactagt ttgagtagaa ctatgctctg tctatctatt tggcacatgt 180
 tctgctgcct ggggagtaag caagctaaag ggatgagaaa gaccacctcc cctaccctg 240
 gaaattgcac tgcaaggcag ggcgagaatg gggtagctgg cagacctggc ctccctgttc 300
 ccagctcttag ttatttcttg cagagattca gtattcagta aagaatagca ttcaattagt 360
 caaaaaatat atatctaact tcttcttttc ccttcccctg aatcattgca cgtcattccc 420
 taagctttct tctctttcca cctcatggcc tgctcagtct tcccatcctt accaatcaca 480
 gactctcagc ctatagacgc agtcacagta tctcaactca tccgcctctg cttcacacta 540
 cattaacaat acctcctcac tcacatacta cataactcca gctctagtct tccaaaattc 600
 acctttcatg atgccactca gcatctcaaa tacctttcat gggctctctg ctgcccagg 660
 ataacaggtc aaagtcatta gcctcaacag tgggcttcaa ccagccttgg gacctcagcc 720
 catttatcca tcacagaggc tggttaactag tctcactgct caggctgtga gtgttcctga 780
 tccttgtagc attctgtgct gtgctttaca tggacaggt ctttctctc tctggcccat 840
 tcgaatctc taatcaagcc catctgattc tgtacagaac acattttcaa gttcaattcc 900
 ctggatgctg ttgcgcgaaa agttgcttaa tgactggg 938

<210> 100
 <211> 376
 <212> DNA
 <213> Homo sapien

<400> 100
 tactcttggg tttcttctc caagactact ccttactcat atcagcaaat agcagctctt 60
 ttcaagtgtc cagtgtaaaa acctacaatt aatccttgat ttctctttca gtcagcctat 120
 actaaatcaa tttcatttaa aatctctcgg ctactactct gcatctccac tgctaccatc 180
 ggctctcca gtcacattct ccaagagcac tctatctcat taaaagaca aaatctctgc 240

agtggcctgt gatgctcctt aatggcctac ataatccagc cctcaagcac ctccgtgac 300
 tctgtaaaac tttcccttgg tcaactgtgct tcagccacat taaccagctt gcatatttct 360
 cacattcacc aagctt 376

<210> 101
 <211> 3661
 <212> DNA
 <213> Homo sapien

<400> 101
 ggacacaact caaccagta acagttagtc aatggctgtg gcaggctaaa tgtggctccc 60
 aaatatgtcc atatcctaata cctacagcc tgtgaatatt accttatata gccaaagagga 120
 ttttgcagat gtgattctga gattgagaga ttatgccaga ttatccaggt aggcccaaaa 180
 tgtaatcacc acagtcctta taggagagggc aagaaagtca agtgtagaag gaggcgatag 240
 aaggagagag ggatttgaag attaataaggc tgcttgcttt gaagacagag ggaagggacc 300
 atcaaccaga aataaacctc tagaagctgg aaaaggcatg gaaatagacc ctcccttaag 360
 gtctctggag ggagtgcagc tttgatttct accgagtaaa attgattttg tacttcagac 420
 ctccaaaact gtaagagaat gactgttgtt ttaaaacat tgagtttgta gtaatttgtt 480
 gcagcagcca caagaaactg gtacaacatc tatatagaat tttttcagat aattgggagg 540
 aaatttgaat atggatggca tattaatatt actgaatcag cattaattt gttagggtgta 600
 ataatgtgat tgtagctatt taggagaata tcctattttt aagagacatg ccaccatatt 660
 tagggagaag tgccaacata tttgcagttt attttcaaat ggttcagagg ctgtctgtgt 720
 acatgagaag acaaagataa ggcaaatgca gcaaaattgt aataattggt gaatccagggt 780
 gaagggacta tggctggtct ttgtactttt ttttccaact tttctgtagg tttaaaattt 840
 tcaaaataaa aatgggaaa tactttaaaa attgtaatca aagacattag tacagaaact 900
 ttcataatgt attttatttt tacagtaaaa ttaatttatg taaattgata gaattttact 960
 aatttcactc ccaagttaca ttaaaaggct tacatatggt tgataatagc atatgtaaac 1020
 tagaactctg aatgatatcc attggtcata atacgtacta tgtagcggtg atggtgactt 1080
 ttgtgattgc acaagtctag agatgcccc aatgacattg acttagacat ctggttattc 1140
 taaggctgaa actgaagttg aatagaagggt tttagtcaaa tactgagatg aaaactgagg 1200
 cagtctctggc ggggggggagt gagtgtgtgt gtatatatac acacatagac atcatgcttc 1260
 taaacattta cagaaagaaa gggtagatta tctacaaaaa aataagaatc agactgatat 1320
 gagatcttac aaacctaacc cccttctctt tcctaaactc cagatttctca tatttctgac 1380

ttcctatattg	atatttacac	ttcgatattt	accaggagtc	ttcaacattt	tgttcaaac	1440
agtactcttg	gttttcttcc	tccaagacta	ctccttactc	atatcagcaa	atagcagctc	1500
ttttcaagtg	ctcagtgtaa	aaacctacaa	ttaatccttg	atctctcttt	cagtcagcct	1560
atactaaatc	aatttcattt	aaaatatctc	ggctactact	ctgcatctcc	actgctacca	1620
tcggcctctc	cagtcacatt	ctccaagagc	actctatctc	atttaaaaga	caaaatctct	1680
gcagtggcct	gtgatgctcc	ttaatggcct	acataatcca	gccctcaagc	acctccgtga	1740
tctctgtaaa	actttccctt	ggtcactgtg	cttcagccac	attaaccagc	ttgcatattt	1800
ctcacattca	ccaagcttgt	tcctgccttg	gggcctttgt	acttaccatg	ttctgttctg	1860
agaatactct	gcctcaagat	atcctacaac	tatcttactg	tattcagctc	tctgctcaag	1920
tattaactga	tgaaacctgt	catccctact	ccactccatg	ttctgcttta	cttaacagca	1980
attgcacata	tggccccctg	aataatatac	athtagtcac	ttatttttac	ttatctgcta	2040
atataaatgt	agactttttc	tattctgttt	actgctgtat	tcccagcatg	ttttatccga	2100
atgtgcagtg	gtttcttttc	ttctccctta	tcgtgggaag	tgatgtgca	aaatacacat	2160
aatggagcct	gaatgtcata	ttgctttcat	acctgtgtga	attttggtaa	gaaaggaaaa	2220
gtagcgattg	acaggtaata	taattacatt	aagtcactct	catagttagc	tgtttattgc	2280
tttctgctc	ttattctcag	tccccaggac	caaatgttga	ccactacctt	ccccacata	2340
taattagggt	atattaccgaa	cgccatgcag	gtggctgtta	aaaggaagat	atatacttac	2400
cttataaact	caacttttcc	ctgttgctct	tctgtctcac	ccctacctcc	atgctttaa	2460
ttaacttttc	aggcttaggc	cttatctctc	agtagagcca	tataaggat	gtgtaaaagc	2520
aggaaaatgt	ttcctgggga	tgaagctttg	aaaagctttt	tttttttttc	ttttggcaat	2580
aaaataaggt	agattcagca	caatacctaa	taactaaaaa	atctgttttt	aattgggtgg	2640
ggcagacagc	aagtgtgtca	tcctggaaga	tactatttgg	gattttatgt	aggtacataa	2700
gagaaaaaag	tgaacaaaag	caaggggcta	ccaggacgcc	gcagtatgct	taacatgtat	2760
tttctaagtt	tgtattatgc	ctttatcttg	gtacttttat	cttctgttct	cacttgatct	2820
ttttgaaatg	tattttaaat	cctaataaaa	atatataaag	tctggaatta	ataaaggatt	2880
aaatgaaact	tttgtatatac	tcactgaaat	tctcagaaaa	aaggggggtg	tggggagggg	2940
gaattgcctg	gggtagtgag	tgaaaattgt	gaccaggttc	ttactaagga	atattggcaac	3000
tgcataatca	aatgtcagtg	gttaccaaac	ttatgaatca	cctgggtgtg	tgtcatagat	3060
tgtctatcct	tgctctctgc	ccccagtgat	ttagatcagt	ggaactatgt	ggggtttaag	3120

79

aaatatacaa tatatatattg tatatatattg tgtgtctcga aagcttcagg gttaaataag 3180
 ttttaactgt ttaggaaaca ctattgtttt aggtatccag tctcaaagac gaaggccttt 3240
 aaaacttact taatttttca ttacatttct tgcccagaaa attgtaaaat acccaacgat 3300
 aacaatgggg aattgtctat cagcacttga ctaaaaagct ttactatcca tgacagcagc 3360
 ctttgcatta ctcaattctg atggcattta acgtcttgaa acccagaaat aaatacctat 3420
 agactcacag tacctgaaag gaatacctaa ttgagacaag agagctatat aaacccaaaa 3480
 ttgcttcaac cacagaatgg aggtctacag gtgcggaagg aaagtttata tggtagaggct 3540
 tggtcgcaaa actattagga atattttcag gttactacac aattttgcca gctcaatatg 3600
 cagttaacac tttttccctc gaatctcctg agcagattta cattgaccgg acccgtagca 3660
 t 3661

<210> 102
 <211> 698
 <212> DNA
 <213> Homo sapien

<400> 102
 acatttccat ttcaccggc ttggagcaga gctgtcgagg agtgctattc taggatcctg 60
 atgatgacca caagggcagt ttgtattcag ctgtccctgg gaacacttcc ctgaaagcgc 120
 tcagggacat tttcatcagg cacagtgctc caggctacgg cactctgtat tgttccctgg 180
 tggcttttagg gggctgggca tcgtagctga aataggacaa cagggagatg gctgagtggt 240
 tttcccaact gccagatgac aacaggtcta tcagataaaa gtcatcatat aacttagaag 300
 aaaccttacc ctcggtgaaa tctcccagca gatcagcaac gaaatggact aagcaacttc 360
 ggtagaaca catggggcta ggatataaac agttcatagg aaaggacacc tgatatcatt 420
 aatgattagg gagagaaatt gggtagctaa cagcaggggt gagagagaaa ctttatagta 480
 ttttctctg tagcttttga attttaagac atatgaatgg atttttttt taattgtaat 540
 taaagtataa tttttttaa agagaaatgt ggagtcattt aacttgtaag acaaaggcta 600
 tcttgaata agaatactgt tcttctatt tgctctagat ttaagtttg gatgggctac 660
 atggtttctt agggcagaac cactcttata gactattt 698

<210> 103
 <211> 1217
 <212> DNA
 <213> Homo sapien

<400> 103
 acatttccat ttcaccggc ttggagcaga gctgtcgagg agtgctattc taggatcctg 60

atgatgacca caagggcagt ttgtattcag ctgtccctgg gaacacttcc ctgaaagcgc 120
 tcagggacat tttcatcagg cacagtgctc caggctacgg cactctgtat tgttccctgg 180
 tggcttttagg gggctgggca tcgtagctga aataggacaa cagggagatg gtgagtgtgt 240
 ttccaactg cagatgacaa caggtctata agcataaagt catcatataa cttaaagaaa 300
 ccttaccctc ggtgaaatct cccagcagat cagcaagaaa tagactaaca attcggtaga 360
 aaaatggggc taggatataa acagttcata ggaaaggaca cctgatatca ttaatgatta 420
 gggagagaaa ttgggtagct aacagcaggg gtgagagaga aactttatag tattttcctc 480
 tgtagctttt gaattttaag acatatgaat ggattttttt ttttaattgta attaaagtat 540
 aattttttta aaagagaaat ttgggagtca ttttaacttgt aagacaaagg ctatcttgta 600
 ataagaatac tgttcttctc atttgctcta gattttaagt ttggatgggc atacatgggt 660
 tttcttaggg cagaaccac tctactagac ctatttaacc ccatgacaga gcctagaagg 720
 aacaggtgta atagaagatg gcatttatgg caagaagggt gatcaagttc tccattagaa 780
 tttgaaccag atctaagtc ttttcttccc ttgtttaaga acggcccggg atgttgact 840
 tcacgggcaa ggccaagtgg gatgcctgga atgagctgaa agggacttcc aaggaagatg 900
 ccatgaaagc ttacatcaac aaagtagaag agctaaagaa aaaatacggg atatgagaga 960
 ctggatttg ttactgtgcc atgtgtttat cctaaaactga gacaatgcct tgtttttttc 1020
 taataccgtg gatggtggga attcgggaaa ataaccagtt aaaccagcta ctcaaggctg 1080
 ctcaccatac ggctctaaca gattaggggc taaaacgatt actgactttc cttgagtagt 1140
 ttttatctga aatcaattaa aagtgtattt gttactttaa aaaaaaaaaa aaaaaaaaaag 1200
 atctttaatt aagcgggt 1217

<210> 104
 <211> 193
 <212> DNA
 <213> Homo sapien

<400> 104
 ccgggcagg tacaatagga tttcaaaata acgttccactg gtaatccttc ctgatgcca 60
 ttttaaaatg aagaccgtct aaatttttct gaccagttat tagttgccct gcctctcgga 120
 aatgtgttta aacttttctt tcaattattt gatacctttt gcccaagaga ttactatctc 180
 tctctttttt ttt 193

<210> 105
 <211> 542

81

<212> DNA
<213> Homo sapien

<400> 105
 ggccgcactt tttttttttt ttttttagtt atatatttaa tgaatcattt ttattgcaaa 60
 gggtaaatta catgaaattg acaaaattta gtccatgtaa tatctatcaa aatacacata 120
 tgtaagtgtg tgtatattta tatatgtata cagtacagtt ttcacaaaaa gcttcaacat 180
 tcctaagaaa cacagacata gtcattctgg tacaatatgg atttaaaata agttcatggg 240
 aatccttcct gatgccaatt ttaaaatgaa gaccgtctaa atttttctga ccagttatta 300
 gttgccttgc ctctcggaaa tgtgtttaa cttttcttcc aattatttga taccttttgc 360
 ccaagagatt actatctctc tctttttttt ttttctttta agacagagtg ttgctctgtc 420
 actcaggttg gagtgcagtg gcacaattcc tgatcactgc aacctctgcc tcccaggctc 480
 aaacgatcct cccacctcag cctccccagt agctgggacc acaggcacat accaccaagc 540
 tt 542

<210> 106
<211> 715
<212> DNA
<213> Homo sapien

<400> 106
 ccgcccgggc aggtcctaaa tagaattcaa gattagacta aatgattttc agcagagcac 60
 attcaagggt ttacattcta tgattgaaaa aaattttttg aaaacttttt atttcattct 120
 ttctgttagg attttgctac aaataacttt gggaatgaat aaagtggaat ggtaactttc 180
 cagtggttca gaattgaatt agacttcttg tgactgtgat gtttggtttc cattgaaata 240
 tatgaagtga gatgtcatat cctgaatata gtttgtcttc cccaattact tgatagcatg 300
 tctgtcagcc agtaaagatt aagaacagag tttctctaaa ttctccgat tattccacta 360
 aggcacatta aaatacttaa ttttgggaaa ccagacatca cagatttctc catgaagtcc 420
 taaatcttct ttaaagtcag aataggtatc ttagttactg acagtattca ggtttttttc 480
 tcccttggtg atatgtcatt ccatcagtga aaaaatattt tctccaagg gatatagaaa 540
 ggtattctgg taatacatta tcatcaatcc tttaacagta acagtctggc acttatcaca 600
 aaaacgacca tttctataa ccagaaagat atcttagatg tcttcacata tatttactat 660
 gctgtagata aagatgcccg ggttatgggc tccatttcat ggctggggtt acgtg 715

<210> 107
<211> 1716
<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1594)..(1594)

<223> a, c, g or t

<400> 107

```

agactgcaat ttctgactaa agcttttaat gccaggttaa acaggagaaa ctttttccac      60
tagaagaaaa tccttgctat ctattttttc caatagaaga aaatcctgct atttatttta      120
tttgatgaat aaacaaattt attgcagtag cttaaaaaaa tttttttttt aaacagtctc      180
actctgtcgc ccaggctgga gtgaagcaat gtgatctcag ctcaactgcaa cctccacctc      240
ccgagtagct gggattacag acatgcacca ccacctcag ctaatttttg tatttttagt      300
ggagacgggg tttcgccatg ttggccaggc tggctcttaa ctctggcct tacgtgatcc      360
gccccccctt ggcttccaa agtgctggga ttacaggtgt gagccactgc acctggcctg      420
tagtagctta aaattttcct tgagaaaatt cctgacttta aaaataacct ttatataagt      480
acaagtgatt gtgacaaatg acgtaaaaat ggcattcatg atgtctgaaa caagcctaaa      540
tagaattcaa gattagacta aatgattttc acaaagcaca ttcaaggttt tacattctat      600
gattgaaaaa aattttttga aaacttttta tttcattctt tcctgtagga ttttgctaca      660
aataactttg ggaatgaata aagtggaatg gtaactttcc agtggttcag aattgaatta      720
gacttcttgt gactgtgatg tttggtttcc attgaaatat atgaagtgag atgtcatatc      780
ctgaatatag tttgtcttcc ccaattactt gatagcatgt ctgtcagcca gtaaagatta      840
agaacagagt ttctctaaat tcctccgatt attccactaa ggcacattaa aatacttaat      900
tttgggaaac cagacatcac agattttctcc atgaagtccct aaatcttctt taaagtcaga      960
ataggtatct tagttactga cagtattcag gtttttttct cccttgggtga tatgtcattc     1020
catcagtgaa aaaatatttt ctcccagga taagaaagggt attctggtaa tacattatca     1080
tcaatcctta aacagtaaca gtcttggcac ttatcacaaa accgacctat ttcttataac     1140
cagaaagatt atcttagact gtccttcaca ttatacttta cctactgcct tgtaagaata     1200
agagttgctc actgtgttta cttgctgtcc tccatattct ccattgcacc attggtgtat     1260
aacgttaaga gtttcattga atattatttt aagtattaca aaaggcagct tgcttcttaa     1320
tctatgcac tttgggggtt ttgaagaaat ttaattcttt gatgtaaaaa ggaactgtta     1380
aaaaagtgg aagctctgca cctgtgtata tatatatttt agcaataaag cagcatgggc     1440
tgagaatgca ctgaaaaaaaa aaaatgctag tgacttcagc aagtcataat cttcctgctg     1500

```

83

gtggagggtc tcactgcat gtggatggcc gctggggctg accaggggtg tgggtggcaga 1560
 aggctggggc ggctgtggca gtttcttaaa atangacaac aatgacattt gccacattga 1620
 tagacttttc ttttcacaaa agaagtctct gtagcacgtg gttgctgttg gtgcacttta 1680
 cccacagtgg aacttctttc aaatagtctc aatcct 1716

<210> 108
 <211> 666
 <212> DNA
 <213> Homo sapien

<400> 108
 tcgcgccga ggtacttaat aatgactgaa tttcatgttc ctacagtcac acatattcat 60
 tagaagtttt atgttgttg tctgatctga ttcttctttg tttgtgggtg gaacggcact 120
 gagagaagta tagtttttta aacttgaaca tgttcagtag ttacattgcc ttagaaaacc 180
 cagacacata gcagtggaat tgaagaaat ggcatcagaa gtgacttaac ttagcaattg 240
 tgattcctct tgtaaaacaa acaaaaaaaaa caatgccata ttttttggag aaaagttggc 300
 aatatagggg tttcgtgtgc tgtttcacaa gaagactcat ttgttctttt gggggaacca 360
 gtgccttaca gattttgtat atactgtaat tattcaggac tagggaacaa acaattgtat 420
 tgtatttgtt acagattgta tatggctttg ttttaacatt cccctaaata aaatggcttc 480
 attctcccct tggaaaaaaaa catgactgtt atgttataaa acaaaaaaaaa aaaaaaaaaa 540
 aaaaaagggtg ggggtaccgg ggcaaacgt gtcccggggg gaatggttc ccggcccaca 600
 aatccccac attgagagaa aaccgtgcga acaaaaaaaaa aaaaaaacg aaaaaaaaaa 660
 acaggg 666

<210> 109
 <211> 1983
 <212> DNA
 <213> Homo sapien

<400> 109
 gaatttcgta atccttgaaa ttgaaaaaaaa aaaaattgtg tttttaaaga gtgaaaacag 60
 ttaggaaaca agtagaactg taatcagaac gctgcttcaa ttgatattaa aaataacctc 120
 aataataatg taaaggttcc tttctcttgt gtcagttata ttcttaggga tagcctagaa 180
 ggaatatatg gttagaacta agtgtgacta atcatctgag ccttgaagag aaacttcagt 240
 gcctctaac agatcatcta caaaacaaca ggtaaacatt tatgccagtt aagtgggtca 300
 tgtttttgtt tcttgggttt ttctaaatt taagtgaggt tgggcttacc ttgtagataa 360
 aattatgtt tcttttgggt aaatactga atgtggataa cgtcaaatca gaatattttg 420

tgaggagggtg atgatttgaa attaagctag atttctaggg aggtggtggt tccaatgaag	480
gatgggaaga aattaaata gtcttcaaac ttcttcctta ttatatttgg ttgctttgga	540
aaagattggt cctatcctca atctaattta ttcactatta atattttaaa aacattcctg	600
agatacttaa aaagaccac ttagcgatta tagttgctca atgaaacaag aatttattta	660
tgcatagatt tttctctgta tcttaccaaa atccacttta cttagataac actaaattgt	720
tcttaaagac tactcatttc ccaataatcc tttatgattt caaaatttct agtggctcag	780
aagtgaattt tattttattt gtctttcact tgaataaatg agaaccaga aattaataat	840
gttgtttatt gcttactgtc aggactattt caaagactaa gaagagttc ttctaacccc	900
tccctctcaa aggaatccta aattattagt tgtagataa gttttgtatg ctaagatatt	960
caggtttata gtttatgtat gtgtgtatat atataaatat atatgtatat ataaatatta	1020
tgttcagttt ggagctctggc acaactccat tatgtggatt agagagtaag atattatgga	1080
tgataaagta ctaaagaaa cataatattt atttataaaa gtgtgtagat tgttaaatca	1140
caaaaagagt gctatgacca ttatgtatga ggaaacaggc ctttgacctc ctggaaagca	1200
ctgctcaaaa gtcattagtg cccatttttg aattccccaa acagaaagct tcttagaaaa	1260
cacgctgaga ttttatttac aggaattctt ttgacacatt tcaattggtg tgtagtcaag	1320
tatagcaagt acttaataat gactgaattt catgttctca cagtcataca tattcattag	1380
aagttttatg ttgttggtct gatctgatc ttctttggtt gtgggtggaa cggcactgag	1440
agaagtatag ttttttaaac ttgaacatgt tcagtagtta cattgcctta gaaaaccag	1500
acacatagca gtggaaatga aagaaatggc atcagaagtg acttaattta gcaattgtga	1560
ttctcttctgt aaaacaaaac aaaaaaaca tgccatattt tttggagaaa agttggcaat	1620
atagggggtt cgttgtctgt ttcacaagaa gactcatttg ttcttttggg ggaaccagtg	1680
ccttacagat tttgtatata ctgtaattat tcaggactag ggaacaaaca attgtattgt	1740
atgtgttaca gattgtatat ggctttggtt taacattccc ctaaataaaa tggcttcatt	1800
ctccccttgg aaaaaaacat gactgttatg ttataaaaca aaaaaaaaaa aaaaaaaaaa	1860
aaaggtgggg gtaccggggc aaaacgtgtc ccggggggaa tggtttcccg gcccaacaat	1920
ccccacatt gcgagaaaac cgtgcgaaca aaaaaaaaaa aaaaacgaaa aaaaaaaca	1980
ggg	1983

<210> 110

<211> 758

<212> DNA

85

<213> Homo sapien

<400> 110

aaaaaaaacc acaaacaaga gaggattgat tgataatatg gggcatgctt aatctaataca 60
 tgctcgagcg gcgcagtagt gatggatcga gcggccgccc ggcaggtacc taacatatag 120
 tagacagtgg agagtggttc tctttcgttg tctcaggggc agacagatgg ggtgctggag 180
 tcctctatca aagagtcaga gctctatccc agatgtgtaa tgaacgtggt cacagacata 240
 ttgtccatt accatttacc ttccctataa ccaactgtgcc tccagccttg tagaatagac 300
 acataggagc gcagcaatac gtctaaaaat aggagtgaga gagggcaggg catgcccgtt 360
 cttgtggtag aagaaaagaa tgtcaaagaa agcagctggg actaatgaac tttacattag 420
 ccatattcca ttatttcagc ttaagtcaaa tgtcggctct catgaggcaa ctggctttga 480
 caggagctac gctaattgac cacttaccaa cctttaattt ctgggtaaaa gcagaaagag 540
 aaaaactaat ggatttttca tttccagaa gagacaagaa tcaactacac tagtagtctg 600
 tcagaacaaa agaaaacctg catccaatta caagaattat tactgtctct ttaataaata 660
 accacattat taaaaaaaaa aaaaacaaaa aagggttggg ggtaccgggg ccaaggggtc 720
 ccggggggaa ttgtttcggc ccatatccat acaaaaaa 758

<210> 111

<211> 3575

<212> DNA

<213> Homo sapien

<400> 111

atgaaattac aactcaggat taagagtctc actcaaaacc gcacaactac atggaaactg 60
 aacaacctgc tctgaaatga ctactgggta aataagaaaa ttaaggcaga aataaataag 120
 ttctttgaaa ccattgagaa caaagacaca atgtaccaga acacagctaa agcagtgttc 180
 agagggaaat tcatagcact aaataccac atcagaaatt gggaaatacc taaaatcaac 240
 gtgctaacat cacaattaaa agaactagag aagcgagagc aaacacattc aaaacaagaa 300
 ataactaaga tcatagcaga actgaaggag atagagacac aaaaagccct tcaaaaaatc 360
 agtgattcca ggagctggtt tttgaaaag attaacaaaa cagatagact gctagccaga 420
 ataataaaga agaaaagaga gaagaatcag atagacacaa taataaatga taaaggggat 480
 atcaccacta accccacaga aatacaaaact gccatcagag aatgctatca acacctctac 540
 ataataaac tagaaaatct agaagaaata ggccggggcg agtggctcac acctgtaatc 600
 ccagcatttt gggaggccaa ggtgggcgga tcacctgagg tcaggagttc gagaccagcc 660
 tagccaacat ggtgaaacct cgtctctact aaaattataa aaaattagcc ggggtgtagt 720

gtacacgcct gtagtcccag ttacttggga ggctgaggca tgagaattgc ttgaaccag 780
 gaagtggagg tggaggtgag ccgaaattgt gccactgtac tccagcctgc aacagagtga 840
 gacactgtca cacaaaaaag aaagaaatat cacaatatgt cacaataggc cgggcgcagt 900
 ggctcacacc tgcagtcca gcactttggg aggccaaggc agatggatca cctgaggtca 960
 ggagtttgag accagcctgg ccaacgtgac aaaaccagc ctactaaaaa tacaaaaatt 1020
 agccaggcgt gatggtgggc acctgtaatc ccagctactc aggaggctga gacatgagaa 1080
 tcgcttgaac ccaggaggtg gagattgcac tgagctgaga tcctgccact gggctccagc 1140
 ctgggtgaca gaatgagact ctgtctttaa aaaaaaaaaa aaaaaaaaaa aatcacaata 1200
 agtcctaggg taaagatggg ggtacagaaa acaattaaat agaacaaaaa caactgtttc 1260
 cttttcctgt gattcaagaa gggcttagat cttctactca gcatcctttt actaatgccc 1320
 tccattggct ctcacgcca acatttcctt tttatagct tattttgtaa tgcctcctta 1380
 attatccttt aatagaagcc accgctgata agctacctac actcatacag aagcattaat 1440
 ataatgcccc agatgtactg tttcagggca aaaaggaaaa taatttcaa caaagtgggtg 1500
 tgtgtctcac tgtcagatgc ttgcacttac acacggaatc gctgtgcatc cgacagaggc 1560
 tgattggcac atggggcacg gggattgtca gctcaaacac cgtcagcagc gttgccttg 1620
 gaaatgggat ttcccagaac agtaaacgtg tctgtccttg atttacagag tagctacatt 1680
 cctaggaaat ccagggtaca ttaaaactca ccatgttacc caggctggtc tcgaactcca 1740
 ggctcaagc aatcctccca catcagcttc ccagaatfff gggattacag gcatgagcca 1800
 ccacaccag ccagaatatt ttatttctgt tagacacaga gcgttcgttg actcgtctgg 1860
 gcgttagtgt taatattctg tacttgaagc aagcccacca agcggctgaa ctgggtggat 1920
 aatggaaaat gtcctgtgga tttgggagtg agacaaaccg gcttgagtct aacctctcag 1980
 ttagtctaag gctccaagct tgaaagggtt aaatgaagta ctatatttgt tttgtttcgt 2040
 tttcgttttg tttgaggctt tgctctgttg cccaggctgg agttagtgg cacaatctct 2100
 gctcactgca acctccatct cccaggttca ggcgattctc ttgcctcagc ctccagagta 2160
 gctgggatta cagggtgccc ccaccacacc cggctaagtt ttttttgta tttttagtag 2220
 acacagggtt tcacatggtt ggctagactg gtctcgaact cctgacctca agtgatccac 2280
 ctgccttggc ctcccaaagt gctgggagta tgggtgggtga gccaccacgc ctggcctaaa 2340
 tgaagtacca catgaccgac cgaccgacct ggggaacata gcaagacccc atctctacaa 2400
 aaatgtaaaa aataaaaatt agccgggtgt agtggtagat gcctgtaatc ctagatactc 2460

87

gggaggctaa ggcagaagga tcacttgagc ccaggagttc gaggctgcag tgagctgtga 2520
 tcgtgccact gcactccatc ctgggtggca gaggtaggcc ctgtctcaaa ataaataatc 2580
 cagtcccccc caagaaagga atgaagtgtc ataatgagaa aaatcctagt acctaacata 2640
 tagtagacag tggagagtgg ttctctttcg tttctcaggg gcagacagat ggggtgctgg 2700
 agtcctctat caaagagtca gagctctatc ccagatgtgt aatgaacgtg gtcacagaca 2760
 tattgtccca ttaccattta ccttcctat aaccactgtg cctccagcct tgtagaatag 2820
 acacatagga ggcagcaat acgtctaaaa ataggagtga gagagggcag ggcagccccg 2880
 ttcttgtggt agaagaaaag aatgtcaaag aaagcagctg ggactaatga actttacatt 2940
 agccatattc cattatttca gcttaagtca aatgtcggtc ctcatgaggc aactggcttt 3000
 gacaggagct acgctaatta ccacttacca accttaatt tctgggtaaa agcaaaagag 3060
 aaaaactaat ggatttttca tttccagag agacaagaat aaaataatag tagtctgtag 3120
 aaaaagaaa acctgcatca attacaagaa ttattaatgt atctttaata aataaccaca 3180
 ttatttagct gtttaatttc ctaaaaaaaaa aaaaaaaaa aaaaaaaaa aaaaaaaaa 3240
 aaaaaaaaa aaaaaaaaa aaaaaaaaa aaaaaaaaa aaaaaaaaa acaaaaaaaaa 3300
 aggagggggg gggggcgaga aaaagagccg aggggggagc acagagcggg ccgccgcgca 3360
 catatgaaaa aagcgacca gaagaagaaa cacaaaacca gcaagcgcaa acagaagaaa 3420
 taagaaagag aaaaagtac gagacgaata gaaaggaaat aactacagga ccaacacggg 3480
 acaaaccaaa agcaataaa caaagaaat aagacagaca caagatgcca acgagctaac 3540
 gcccgacaa tggaaacagg taaacaacat aaagc 3575

<210> 112
 <211> 442
 <212> DNA
 <213> Homo sapien

<400> 112
 actgagcagc tacggaagtg caaggcactg taggagtagg gtgagtatac tccccacaag 60
 ggctcagggt caggcagggg acggtagaga taaaaacca cagaccatac acatagctgg 120
 cactgtctct gagggttttg tgaggcacac aaatgcttag gagactagac gaagtaagac 180
 aatgtctttg acatgaggca gaaatcaacg gaaagcatgc gcttttagaa catgtgtggg 240
 actgtttttt ggtatcagca gactgaagag gctttttaa cgtggagga aggcaactg 300
 aggcatagag atgccaatac caggtcttgt caggaagaac agagtccaat ttggctgcag 360
 gatagggcat atgtagggga ggggataaga ctggcatggg ggcagagggg gacttgaatg 420

tcaggtgaca gagtcaaagc tt 442

<210> 113
 <211> 412
 <212> DNA
 <213> Homo sapien

<400> 113
 tgtcatacta taaggcgaac tgggcctcta gatgcaattg ctcgagcggc gcaggtgatg 60
 gatgttcgcg gcgaggtatc agaagctgtg atgtctgcct tgtagtcttg tgcttggtac 120
 tgtaatTTTT tttttttttt tacgaagcac gtgactggac taatgtaagg cagatgacgt 180
 gatctttaag actgctatat atatcagtct cttactctat aaggttttaa attagaaaag 240
 gcttatatgg ttaactacct tagactatat ctacagcagg gtctggtttg ccagaacaag 300
 tttaaagtgg ctgtttatta agttggctat ttcagaatt gaaactataa gaccgccatt 360
 tgacactgaa acttgcgtga atcctaaatt gcatcaatta tctatttgat aa 412

<210> 114
 <211> 625
 <212> DNA
 <213> Homo sapien

<400> 114
 gcaacaacaa ctgaatggct gtaatacatt aatgtataca gatgaattga gaagtcttct 60
 agtgaaatgg ctcagatctt tgttcttggc ccagtcctgt tcagtttttg atcagtgcct 120
 tgcaatatca cttgatcgac tcaactaaca ttatacaag agtgcagagg cctcctcaga 180
 gaatggatgg tagaaatgca ttgatgagag aacgtttatc tatctatctg tctatctatc 240
 tatctatcta tctgtctcta tctaagaagt cataaaggct gagtctaata aggcaaaaaa 300
 aaaaagaaaa aaaaaaaaag ctgtggcgat acccagggcc aaagcgtgat cccggcgcca 360
 actgcgaaat cgcctcaca tccaacaac acccccacaa ccccccccc agccccaaca 420
 ccaacacctc acaaaaacct cacaacacct ccaccacctc acacagccac cccctacca 480
 cacaaccaca tcacaccacc accgccccac ccacaccaca caccacacac aactccgaa 540
 caccaccgcc cactccacac acccaccac caccagcacc aacacacca cacaccaca 600
 ccgccacccc cacaccacac gcagc 625

<210> 115
 <211> 378
 <212> DNA
 <213> Homo sapien

<400> 115

89

gcggcgcccc ggcaggtaca tagtgcagat gcagtatata atttcaggct aggaaaatta 60
gctactagta tgtatctgac agtccctaag agctaagagg cctaagaatg cagacgggga 120
gaaaaaaaaac caaaaccaa aaaaaagaca cctctccaat tgctgggagg gcctgggaat 180
aggatgaagat caaacacag tgggagagga gggtaaagat gtgagcttca agcgggtaat 240
gggcaagcca cacctcccag ttcctaggag ggaatcgcca cggccgactt cagcattctc 300
gtctttacta agacttacc atagagaact acagcaggaa accgatttct tcattcattc 360
tctttaaaaa gtatgaat 378

<210> 116
<211> 8905
<212> DNA
<213> Homo sapien

<400> 116
atggcggcgg cgctggggcc cccagaagtg atcgctcagc tggagaacgc ggctaaagtt 60
ctgatggtga ggacgcgcg cccctcagac cccgggattc gcgggcccc ggtcgccct 120
gccactccag gcottgctgc tcgctgggct ggcgactggc aagggcctgc agggagcctg 180
gaagtggagg aggaggtggc ggtggcgtag cgcaggattc ttcagcctac tttcctcctg 240
ccgtcgtccc ctccttcag gagctgtccc cttcccctgg ctgccagca cccagtcgg 300
gcgtgggaat atagtgggtg agcaaagaga atttctcac cttacacct gccccacaga 360
ctgggtcgca gagcaaggcg cgggaagga gttgggggta tccccgcagg gcttcgggccc 420
tctcatatac tagtccttct gtctggaatg ctttcttcc ctgtcacttc atccttcagt 480
tctctcagta gtcagtttct caggaagcc ttccttagcc tgctgaaag tataccctgg 540
gtgatagatt ggattggatt ggattggatc ggatcggatc ggatcggatt ggattatatt 600
gtatttattt ttaagagaca gggcagctgt caaaatggaa gttcagggtc actagagggt 660
ggcacatgtc tccagggtaa acacatgagt gcttgcatte atctttggat ccctgcgttc 720
gcttctggtt tagcttttga tgattcctta atttctctg ccacagccat aatggaagca 780
gttgctcgag agtggattct cttggaaaa ggtagcatcg agtctctgcg aacattcctt 840
ttaacctatg tcttacaag gcccaacctt caaaagtatg ttcgggaaca gattctacta 900
gcagtagcag taattgtaaa aagaggatca ttagataaat caattgactg caaaagcatt 960
tttcatgaag tcagccagtt gattagtagt ggcaatcca ctgtgcaaac tctggcctgt 1020
tctattctga ctgcgctatt gagtgaattt tcaagtcaa gtaaaactag caacattgga 1080
ttgagcatgg aattccatgg taactgcaaa aagagttttt caggaagaag accttcgtca 1140

gatcttcatg ttaactgttg aagttctgca ggagttcagc aggcgggaaa acctcaatgc 1200
tcagatgtct tcagtatctc agcgttacct tgcactcgcc aatcaagtct tgagctggaa 1260
ctttcttctc ccaaatttgg gcagacatta tatagctatg tttgaatcct cgcaaaatgt 1320
gctgttgaag ccaacagagt cctgctggag actcttctgg acagcagagt tatggagctt 1380
ttcttcacag tacatcgaaa aatccgagaa gcattcagat atggcaccaa gattctctgc 1440
agtgccctgc ccagttagct tctcttcatg gacctatctt cccagatgaa ggatcacaag 1500
ttgattatct agcacacttc attgagggat tactgaatac tatcaatgga attgaaatag 1560
aagattctga agctgtgggg atctccagca ttatcagcaa cctgataacc gtgttccac 1620
gaaatgtttt aactgccatt ccaagtgaac ttttctctc ctttgtaac tgcctcacac 1680
acctcacttg ttcttttggg cgaagtgctg cattggaaga agtgcttgat aaagatgaca 1740
tggatacat ggaagcatat gataaattgt tggagtctct gttaactttg gttcaagatg 1800
acaaacattt ccataaaggc ttttttacc aacatgcagt tcaagttttc aattcctata 1860
ttcagtgcca cctagctgct ccagatggca caagaaattt gactgccaat ggtgtggcct 1920
ctcgtgagga ggaagaaata agtgaacttc aagaggatga tcgagaccag ttttctgatc 1980
aactggccag tgtaggaatg ctaggaagaa ttgctgcaga aactgtata cctcttctga 2040
caagtttatt agaagaaaga gtaacaagac tccatgggtca gttacaacga catcagcaac 2100
agttacttgc ttcaccgggt tcaagcactg ttgacaacaa aatgcttgat gatctctatg 2160
aagatattca ctggcttatt ttagttagc gctacctctt agctgatgat actcagggag 2220
agactccgct aatacctcca gaaataatgg aatattccat taagcattca tctgaagttg 2280
acattaatac aacacttcaa attttggat ctccaggaga aaaggcttct tccatcccag 2340
ggtacaacag aacagattct gtgattagc tggttgtctgc cattctcaga gtttcagaag 2400
ttgaatctcg agcaataaga gcagatctca ctcatctact aagtccccag atgggcaaag 2460
atattgtttg gtttttaaaa cgctgggcaa agacttatct cctgggtggat gaaaaactgt 2520
atgatcagat aagtctgcca ttcagtacag cgttcggagc agatacagag ggttctcagt 2580
ggataattgg ctaccttta caaaaagtca tcagtaacct ctcagtctgg agtagtgagc 2640
aggaccttgc aatgacact gtgcagctcc ttgtcacttt ggtggaaaga agagaaaggg 2700
caaacttagt aattcaatgt gagaactggt ggaatttagc taagcagttt gcaagccgaa 2760
gcccacctct taatttcttg tcaagtcttg tgcagaggac attgatgaag gctctagtct 2820
taggaggttt tgcacatag gacacagaaa ccaaacagca gtattggaca gaggttcttc 2880
agccacttca gcagcgattc ttaagagtga taaaccaaga aaacttccag cagatgtgtc 2940

agcaagagga agtcaagcag gaaatcactg ccacactaga ggcctgtgt ggcattgctg 3000
aggctaccca gattgacaac gtagcaatcc tgtttaattt tttaatggac ttccttacca 3060
attgcattgg attgatggaa gtttacaaga ataccccaga gactgtcaat ctcatatag 3120
aagtttttgt tgaagttgca cataaacaga tatgctatct tggagagtcc aaagctatga 3180
acttatatga agcctgcctt actttgttgc aagtgtattc taagaataat ttagggcggc 3240
aaagaataga tgttacagca gaagaagagc aataccaaga cctgcttctc attatggaac 3300
ttcttactaa cctgctgtca aaagaattca tagatttcag tgatacagat gaagtgttta 3360
gaggacatga gccagggtcaa gcagcaaaca gatctgtgtc agcagcggat gttgtgttgt 3420
atggagtaaa cctaattctg cccttgatgt cacaggatct cttgaagttt ccaacccttt 3480
gtaatcagta ctacaaatta atcacattta tctgtgagat ttttcctgaa aaaataccac 3540
agcttcctga ggatctgttt aaaagtctga tgtactccct agaattagga atgacatcaa 3600
tgagttcggg ggtttgccag ctttgccctgg aggccttgac accgtagct gaacagtgtg 3660
caaaagcaca agaaacagac tcaccacttt ttctagcaac acggcacttt cttaaactgg 3720
tttttgatat gctggttttg caaaagcaca acacagagat gaccactgcg gctggcgaag 3780
ctttctacac gttggtgtgt ttgcaccagg ctgaatattc tgaactggtc gaaacattac 3840
tatcaagtca gcaagacca gttatttacc agagattagc agatgccttc aacaagctca 3900
ctgcaagcag cactcctcct acgctggatc ggaagcagaa gatggccttc ttaaagagtt 3960
tagaagaatt tatggcaaat gttggtgtgt tcctttgtgt aaaataaaca acagaacttt 4020
atgcttaatt tagatccttt ctgcaaagtg cactgaattg ctgaaagtgt acttgagtct 4080
tgtcctattc ctgagttcat ttggccattt tggattttgg agagcctgaa actttgatat 4140
gtatgtaata cagtgaaca ggagaggtca acttggcatc agcttctgct gttaagtgtt 4200
agccacaatc tgtcatatat atgtctttta gattctgaat ggtgatttaa aattttcaaa 4260
atgaaattcc atatatgtgc aaacagatat ggcaccacg aaatacatat gcagtgcctt 4320
ttttcctttt aacataggtg gctagccaaa gtttagaatt tttgtcatta aatatgaaat 4380
ggatataatg taggcagtgt ttctcaaaat ctccacagat cgcctgcac cactgaggag 4440
ctggtgaaaa ggcagattct taggccaac thtagacctt cagagtcaga atgtctggtt 4500
gttgggcca ggagtcttca tgtaataag cttctccctt tcgtcacc ccaagttttg 4560
aatcaatgaa agagacattg aaaactctta agaggttttg tgctttctag cttttcctcc 4620
ctttgatgat tgggttttat aattcagcag gaaggggaaa catcatcagg ggtttgttgg 4680

ctttttctta gcttgcttcc ttgcttgctt gctttcttgc ttttcttgct ttctgtctct 4740
ctctttcttt tctctctctc tctcacatca acccagtgtc gcaggttttg tgtaatacaa 4800
gtcactaatc atactctgat gcctgaactt gaggaggaaa atacatgtat atttttgttc 4860
cgtaaaaaata accttaggaa ctgtagccat ttcattgcct taattttaag aggaaaaatac 4920
aaaaacagct gatttgtttt agtaagaaac cacgtcttga tgcttcagag ttgggttagg 4980
gtgttagctg ctatgaacct gttgccctt tcgatcgtgt atttatgtag gtttatcagt 5040
gaaatgaaag gcttgtttcc gtctagtcta actttttgag tgtgtttcta tccagccaca 5100
tagcccatat ctactctaaa tggcttgctt aagcaataat tattttaaag gatgtgaatc 5160
actgattcac acagactatt gcacgttggg gcattagggg caataattct tatccagaca 5220
tgaggagccag tgaatttaac ttcagagatt aaaaattcac tttagatcct ctagttagat 5280
ctcttaataca ggatttttat acagctgccg ggctccccta attcagtgtg ccagcttaca 5340
atgtggaaat gaaagctaat ttatacacag caggcatatg aaactccact cattgcagta 5400
ctttcacagc acagtgcagc gtagaggact ctggcacagg tgcaactcatg aaactctgct 5460
tccaccatgt tcotgacacc tatctattaa accattctgc aaatacgggt tttctacctg 5520
attgcataata gcatatgtgt cattacatgt gatgctgtgc aaaactttgt ataattctgt 5580
gttattaaca gttaacaaaa ctggagcacc tgaattacat ccaacctgtg catgtgatgt 5640
taggtagatg tgaatgcagg gccttgggcc ataacttaca tttctctcaa tttgattagc 5700
tttgagtcac aattaagggg aagcaaaaac atcttgaaaa gactgctagg aaggaaatta 5760
atatcagtca tccagaagta cacgtttctg tattttaaaa aatactttga tgcatttatt 5820
tttaggtgtt ttttttttcc ccttaaaaaa cttgaagtga tatgcagcag taatctattt 5880
gttttgcatg gttcttggtg ttttgtgtt occagatccc tcaagctttc tcagctgttg 5940
cgaattatgt gtatctgtgt gtgtgctaag tacagtctct ttaccaaagg gcactgaaac 6000
acacaattga ctggacaggt ccacgcgcca tgacaaaact ataatacagt tattaact 6060
aaagaggagt gggaaaggaa tgccttggtg agtaaaaagg catctatatt taataacttt 6120
tatccagatg gcaacatatt tgcaaaatth gccagatcc tattacaata ctaaaaatag 6180
aaaatttcac ctccatattc ctgaggtgta atttcattag actagtttta gtttaaaaag 6240
accttcttca gattggacca aataaactt ataagatcag cagaatgttg aatattagct 6300
cactgggggtg gggagaagcc actaccatth tttaggtgat ggggatgcca ctgagttgca 6360
acggctagac cttttcaggg tgggtgtgtc catgtttgcc tgattggatg cttattcact 6420
ttgtgttttc ttttgtttta ttttgtccaa ttttgtctth agctgtgttt attaacttct 6480

ccggctcttgt tttgttttaa tgctcttggc ccagtgggtg tcaagaacac tggcttaatt 6540
 caagtcagtt gatttttttt ctattaaaac tgttgtaaa atatttttta aaacaaaaac 6600
 attatttgtg ccctcttcta tatatgtcaa agggacactg tcaagtattt catttttaga 6660
 tttttgtttt ataaaatttc tgttgttcat atagtatcct ttaacctcta gttttccata 6720
 catccttctg ttgtttctca ttttattttc cttgacctat ttatttccca aggcacaatc 6780
 actaaagact ttgtactttc acagtctggt aatgtggtag cacctgtaac tgtgttcttg 6840
 ttctgttaaa aggattgatt tgcttttata gtccttctgc tggatgagtg gctgcctcag 6900
 tagcaaaact acctgacagt atttgacagt gtcctttcca gcaccattat ttgggtcttt 6960
 cagggtggcc atctctgtta gaagacagta gcatgttaac atcactgcat tgagtttttg 7020
 tctggtgtaa agtatgactt ttaatgtaaa caaactgcag gtttttttca aactaatttt 7080
 aagaatttag tcttatttcg ttgtaaactg tgtatctaata tatattacat tactctgttc 7140
 agatgggatg gttactacca cttgtccatg attttcattt gaaaagcaag tatctatata 7200
 atttcccccc agtcagcatt atttaacact ccccttaact gtctttgaac tttctctttt 7260
 aacaaaaatg tcaagtcttt acagttgtaa tatcaccatg tttccattt ctgttaatac 7320
 ttctatgaac ccctaaagta ttgaaggaa ctagctgtca gtttcaagga ttacaagttt 7380
 gagtctccta gtattcaaca tcattctgaa ccctgaaata atatttttct ctgttaaaaca 7440
 atttttatct gtttgccacc tctgttgta gaggtggtg tcaattgacc ttactaagtt 7500
 agctgtcttt gatgaggaat tattgttatt ggctcctgaa taaaacatta accttttaag 7560
 tcagaaggaa cctcggact tcttaaggtt tgtttgtgtt ttctaaaacc agagaataag 7620
 gaactgattt ggctatgagg ttaaacatta taattttctg taagctttcc cacaaaaaaa 7680
 cattgtgat ttgaggatat aataatgttt taatcttttt aaaatataag tggttattct 7740
 ctgacttggg aactatgttc tgaaaacact gcatttaaga atttttaaaa attggttttc 7800
 taaaattaaa atgtccaaat taggcatatt gctgagctca aattgatgtg aaatgccatg 7860
 gttccagttg aattttaagc atattttcat ttagatataa aatatatgaa gtatgctttg 7920
 ttgattatag tgagaacca tgacatagtt aaccaaagaa tatgtttggt tcaaataaaa 7980
 atagaagctt aatactgggc attcatactt ttaaaagaga atgaatgaag aaatcggttt 8040
 cctgctgtag ttctctatgg gtaagtctta gtaaagacga gaatgctgaa gtcggccgtg 8100
 gcgattccct cctaggaact gggaggtgtg gcttgcccat taccgcttg aagctcacat 8160
 ctttaccctc ctotcccact gtggtttgat ettcacctat tcccaggccc tcccagcaat 8220

94

tggagaggtg tctttttttt ttggttttgg ttttttttct ccccgctcgc attcttaggc 8280
 ctcttagcta ttaggaactg tcagatacat actagtagct aattttccta gcctgaaatt 8340
 atatactgca tctgcactat gtacctacta gggatctgac ctcaagtgtt ttctgagccc 8400
 aggcttctctg gtgtggtgtc ttttaccaca taaaattatt acaaattgca aatgttggtg 8460
 ttgtgatttg attatctgta caaagaaaga agctctatgc agtgagtgtg tggtttaatg 8520
 gtcacaaaaa tgtagcact gctaccactc agcacgtgta aaatttttta aatttataaa 8580
 tattaaatt ttaaacttac actaagactt ttcagtttta tttaaagacc cagggatgag 8640
 tgtactgttt aaatatttac ctctattaac ataactaatg aaggataaaa attgcattta 8700
 gtttttcaga agatgctgca atatgatttt aggaaataag gctatgtatt gagccagtta 8760
 taggctgaat atcaggttga taaaatttta tttgtatttt taaaattcat aaatgggagt 8820
 taaaatgtgt cttttcacta aatattttta ttacaaaaaa aaaaaaaaaa aaaaaaaaaa 8880
 aaaaaaaaaa aaaaaaactg cggcc 8905

<210> 117
 <211> 827
 <212> DNA
 <213> Homo sapien

<400> 117
 tcgcgccga ggtaccctgc atcaactgcc tggttgtgct attctcatct caacatagaa 60
 ttggtgggtt ctctaagggt tgcaggaac ctctaaaag atgtgattct ttgggagggg 120
 atatttgaat ttccaacttc cattccccct agcaaaagga agcagctgct gtttaagggg 180
 tttatctgag ccactttaa gatgaatcca tggattact ctggatacta gccattcctt 240
 aggattttaa ggtcacattt tattcctgga tgctttatgt cccacactcc acctgagccc 300
 tcatcctctg ttccctacta tactcccac ttctactctt tgttttatcc acctatocct 360
 attacctgac cttttgtctt cctgtctcc catccttggg gggacatgca gcctgtggt 420
 catggttctg atgacatcat cagggcagcc ctctgccca ggtattatgg cctgtcagca 480
 ttccctgtgc cctccaaacc ttaggcctag aatgcggagc tgccaacata acattcacc 540
 ttttgaacag atggagtcag gcacactaac acagccttct gtcctcaata acacagccat 600
 tattgccact tggctcagtc gtcaatgtaa accctcagag tcagctgaac tattttaggc 660
 caaacatact gtttttgtaa agtatttttc attaataaat ctataagaca gttctattta 720
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aggctggggc gaaccggggc caacggctcc 780
 cgggggaaat tgtttccgc caaattcccc caaaaaatgg caaaacg 827

<210> 118
 <211> 6470
 <212> DNA
 <213> Homo sapien

<400> 118
 ggctccctgg tagctatagc agccgcggcg gttaagtatg cggcgccagg agctgctaaa 60
 tgtgaacaat aatgtcttgg aagagaaatt atttttcagg gggtcgtggg agtgtacaag 120
 ggatgtttgc acctcgaagc tcaacctcca tagccccag caaaggcctc agcaatgagc 180
 cagggcaaaa cagctgcttc ctcaacagtg ccctgcaggg tttgtggcac ttggatatct 240
 tccgacgtag ctttaggcag cttacaactc acaagtgcac gggagattcc tgcactcttt 300
 gcgctctcaa ggggaatcttt aaccagtttc agtgtagtatg tgaaaaagtg cttccatctg 360
 aactctccg cagtgctctg gcaaagactt tccaggatga acaacgtttc cagctgggaa 420
 ttatggatga tgctgcagag tgctttgaaa acctcctgat gagaattcac ttccacattg 480
 ctgatgaaac caaagaggat atatgtactg cccaacactg catttcccat cagaaatttg 540
 caatgacatt gtttgagcag tgtgtatgta ctactgtggg tgccacttct gatccgctgc 600
 ctttcatcca gatggtacat tatactcca ccacttcctc ttgcaatcag gctatttgta 660
 tgctggaaag acgagagaaa ccttcaccaa gcatggttgg tgagctgctg cagaatgcc 720
 gcaccatggg ggatctgagg aactgtccaa gcaactgtgg agagaggatc aggattcgcc 780
 gtgtgttgat gaatgtcca cagattatca cgattgggct ggtatgggac tcagaccact 840
 cagacttagc agaagatggt atccacagcc tgggaacctg ccttaagctg ggtgatctgt 900
 ttttcagagt gacggatgac cgggccaagc aatctgaact gtacttagtt ggaatgatct 960
 gttactatgg caaacattat tctacattct tttttcaaac aaagattcgc aaatggatgt 1020
 attttgatga tgctcatgct aaggagattg ggcccaaatg gaaggatgtg gtgaccaaat 1080
 gcatcaaggg gcattatcag cccctgctgc tgctttatgc agatccccag ggtaccccag 1140
 tttccacca ggacctgcct cccaagctg agttccagtc atacagcagg acatgctacg 1200
 acagtgaaga ttcaggacac ctgactgata gtgaatgtaa tcagaaacac acatccaaga 1260
 aagggtcact gatagagcgc aagaggagct ctggtcgggt taggaggaaa ggcgatgagc 1320
 cccaggcctc gggataccac agtgaaggag aaacactgaa agagaagcag gctcctagaa 1380
 atgcctcaa accatccagc agcaccaaca ggctgagaga ttttaaagag acagtacgca 1440
 atatgatcca taacagacca tccttggtct ctcagaccaa tgtaggctct cactgcaggg 1500
 gcagaggagg agaccagcct gacaaaaaac ctcctaggac cctgccttta cactctcgtg 1560

actgggaaat agagagtacc agcagtgagt caaaatccag ttcttccagc aagtatcgtc 1620
 ccacatggag acccaaacga gaatctctga atattgacag tatctttagt aaggacaaaa 1680
 ggaagcactg tggctatacc cagcttagcc ccttttctga ggattcagct aaagaattta 1740
 taccagatga accaagcaag ccaccttctt acgacattaa atttgggtgga ccaagcccc 1800
 agtacaagcg ctggggccca gcacggccag gctctcacct tttagagcag ccccccgac 1860
 taatccagcg aatggaatct ggctatgaaa gcagtgagag gaacagcagc agccctgtca 1920
 gcctggatgc agccctgctt gagagctcaa atgtctacag ggatccaagt gctaagagat 1980
 cagctgggtt ggttccttcc tggcgtcata tcccaaagtc gcacagcagt agcatcctgg 2040
 aggtagactc cacagcatcc atgggtggct ggacaaagag tcagccttcc tctggtgagg 2100
 agatatcttc taaaagtgaa ctggatgaat tgcaggaaga ggtggccagg agggcgagg 2160
 aacaggaact tcgaagaaaa cgggagaagg agttagaggc agcgaaaggg tttaacctc 2220
 atcctagccg cttcatggac ttggatgaac tgcagaatca ggggaggagt gacggctttg 2280
 agaggtcctt gcaagaggca gagtcaagtgt ttgaagagtc actacatctg gaacagaaag 2340
 gagactgtgc tgcagctttg gctctctgta atgaagctat ctctaaacta agacttgccc 2400
 tgcattggtc cagctgtagc acgcacagca gagccttagt cgataagaag ttgcaaatca 2460
 gtattcgaaa agcacggagc ctgcaggatc gcatgcagca gcagcaatca ccacagcagc 2520
 cgtcgagacc ctgagcctgc ctcccaacac aggcggggac tctctctcag ccaacaagtg 2580
 aacagcctat cccgctcaa gtattgtaa gccaaaggc ccaactggaa tccggcatgg 2640
 atacagagtt tggggccagt tctttcttcc attcacctgc ttctgcccatt gagtcaact 2700
 catcactatc tccagagtca tctgccccac agcacagctc cccagtaga tctgccttga 2760
 agcttctgac ttcgggtgaa gtagacaaca ttgaacctc tgcattccac aggcaagggt 2820
 tacctaaagc accagggtgg actgagaaga attctcatca tagttgggag ccattggatg 2880
 cccagagggt taagctgcaa ggctctaggt gtgacaacag cagttgcagc aagctccctc 2940
 cacaagaagg aagaggcatt gctcaagaac agctgttcca agaaaagaag gatcctgcta 3000
 acccctcccc ggtgatgcct ggaatagcca cctctgagag gggatgatgaa cacagcctag 3060
 gctgtagtcc ttcaaattca tcagctcagc ccagccttcc cctgtataga acctgccacc 3120
 ccataatgcc tgttgccttct tcatttgtgc ttcactgtcc tgatcctgtg cagaaaacta 3180
 accaatgcct ccaaggccaa agcctcaaaa cttcattgac tttaaaagtg gacagaggca 3240
 gtgaggagac ctataggcca gagtttccca gcacaaaggg gcttgtccgt tctctggctg 3300
 agcagttcca gaggatgcag ggtgtctcca tgagggatag tacaggttcc aaggatagaa 3360

gtttgtcagg tagtctaagg aagaactcct ccccttctga ttctaagcct cctttctcac 3420
agggtcaaga gaaaggccac tggccatggg caaagcaaca atcctctctg gaggggtggg 3480
atagaccact ttcctgggaa gagtccactg aacattcttc tcttgcccta aactctgggc 3540
tgcctaattg tgaaacttct agcggaggac agcccaggtt ggcagagcca gacatatacc 3600
aagagaagct gtcccaagtg agagatgtta ggtctaagga tctgggcagc agtactgact 3660
tggggacttc cttgcctttg gattcctggg tgaatatcac aaggttctgt gattctcagc 3720
ttaagcatgg ggcacctagg ccaggaatga agtcctcccc tcatgattcc catacgtgtg 3780
taacctatcc agagagaaat cacatccttt tgcattccaca ttggaaccaa gacacagagc 3840
aggagacctc agaattggag tctctgtatc aggccagtct tcaggcttct caagctggct 3900
gttctggatg ggggcagcag gataccgcct ggcaccact tagccaaaca ggctctgcag 3960
atggcatggg gaggaggttg cactcagccc atgatcctgg tctctcaaag acttcaacag 4020
cagaaatgga gcatggtctc catgaagcca gaacagtgcg tacttctcag gctacacctt 4080
gccgaggcct cagcaggag tgtggggagg atgagcagta cagtgcagag aatttacgtc 4140
gcatctcacg cagtctcagt ggcaccgttg tctcagagag ggaggaagct ccggtttctt 4200
cccacagttt tgattcatca aacgtgagga agcctttgga aaccgggcac cgttgttcca 4260
gctcctcttc cctccctgtc atccatgacc cttctgtgtt tctcctcggc ccccaactct 4320
accttcccca accacagttc ctgtccccag atgtcctgat gccaccatg gcaggggagc 4380
ccaatagact ccaggaact tcaaggagtg tccagcagtt tctggctatg tgtgacaggg 4440
gtgaaacttc ccaaggggcc aagtacacag gaaggacttt gaactaccag agcctcccc 4500
atcgtccag aacagacaac tcctgggcac cctggtcaga gaccaaccag catattggga 4560
ccagattcct gactactcca gggtgcaatc ctcaactaac ctacactgcc aactaccag 4620
aaagaagcaa gggcctcag gttcctcaca ctcagtcctg gagtgatctt ttccattcac 4680
cctcccaccc tccattgtt catcctgtgt acccaccatc tagcagtctt catgtacccc 4740
tgaggtcagc ttggaattca gatcctgttc cagggtcccg aaccctggc cctcgaagag 4800
tagatatgcc ccagatgat gactggaggc aaagcagtta tgctccac tctggacaca 4860
ggagaacagt gggagagggg tttctgtttg ttctatcaga tgctcccaga agagagcaga 4920
tcagggctag agtcctgcag cacagtcaat ggtaaagggtt attcctttcc tttcctggag 4980
ctacaccttt ctttgtaaaa ctgtactgtg ggccgggcgc ggtggctcac acctgtaatc 5040
ccagcacttt gggaggctga ggcgggtgga tcacgaggtc aggagattga gaccatcctg 5100

gccaacatgg tgaaaccccg tctctaccaa aatacaaaaa attagccagg cgtgacggtg 5160
 cgtgcctgta gtcccaacta ctcggaaggc tgaggcagga gaattgcttg aacccgggag 5220
 gcagagggtg cagtgagccg agatcgcacc actgcactcc agcttggcaa tagagtgaga 5280
 ctccatctca aaaaacaaaa caaaacaaca acaaaaataaa ctactgtggc agcgttggta 5340
 ccctgcatca ctgccatggt tgtgctattc tcatctcaac atagaattgg tgggttctcc 5400
 taaggggtgc aggaacctct aaaaagatgt gattctttgg gaggggatat ttgaaattcc 5460
 aacttcatt ccccctagca aaaggaagca gctgctgttt aagggtttta tctgagccac 5520
 tttaaagatg aatccatggt attactctgg aactagcca ttccttagga ttttaaggtc 5580
 acattttatt cctggatgct ttatgtcccc acctccacct gagccctcat cctctgttcc 5640
 ctactatact cccaacttct actctttggt ttatccacct atccctatta cctgaccctt 5700
 tgtcttcctt gtctcccata cttgggggga catgtagccc tgtggtcatg gttctgatga 5760
 catcatcagg gcagccccc tgcccaggta ttatggcctg tcagcattcc ctgtgccctc 5820
 caaaccttag gcctagaatg cggagctgcc aacataacat tcaccctttt gaacagatgg 5880
 agtcaggcac actaacacag ccttctgtcc tcaataacac agccattatt gccacttgct 5940
 cagtcgtcaa tgtaaacctt cagagtcagc tgaactatbt taggccaaac atactgtttt 6000
 tgtaaagtat ttttcattaa taaatctata agacagttct atttaaaaaa aaaaaaaaaa 6060
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaacaaaaaa aaaaaaaaaa aaaaaaaaaa 6120
 aaacaggtgg gggccgcgcg cgggcccgcc cccgagagaa aatttccaca aacaccctg 6180
 ggggggcccgc gggcgcccc aagtgagtgc agagaaagaa agacgaggag cacaacagga 6240
 ggtccgtcct ccagaagaaa acaaccgcgt gcggcacaga acaaaggagg tggcgggggg 6300
 tgcgctccac cacgacaaat aagaaaacct cgcggggggg ggaaaaacag cagacgagtg 6360
 tcgtgaagaa caacaacca caggagagag gcctcgtgga caaggcacac agggggtgct 6420
 cacaaaacaa ggggggtaca agaaggagac gcaagaaaac ataattgccc 6470

 <210> 119
 <211> 435
 <212> DNA
 <213> Homo sapien

 <400> 119
 gtataatcat ataggcgcac ggttctctaa tgctgctcga gggcgcgctg tgatggatgc 60
 gtggcgcggc gaggtacctc tcaacactga gaactgtagt agttgaaacc actgttctag 120
 tgggcagtta gaacagttgt tttcccctgc ttgttcccga cagagctgcc caagttatta 180

tctgctcctg ggggttgacc atctgtttta tgacagttat gatattggtg tttaaaaaaa 240
 atccaaattg ttactttgat ttatatgatc taactctgaa tcacggaagt attactatga 300
 tgttcaaaac tctgattgac tctacttgct ttaaaaactc tcagatccct tctgcattta 360
 tcatcagaga tcggtaaaga tgacaacaag caggctctaaa gttctgagat gtttagcacat 420
 acccttttca caatt 435

<210> 120
 <211> 1262
 <212> DNA
 <213> Homo sapien

<400> 120
 ggccgagttt tttttttttt tttttttttt tgtttttttt tttttttttt tttttttttt 60
 tgttttagat atgttgcttt tattcaaaag aataaaatgc ttgacaaact ctttaatcac 120
 aaggtttgaa ccaaaccacc agtcttctac aacaactctg tgaggtaggt atctgcatag 180
 ccacaaggga tccacatagt cctttcttcc cttgtacctc tcaaactctg agaattgtag 240
 tagttgaaac cactgttcta gtgggcagtt agaacagttg ttttccccgt cttgttcccc 300
 acagagctgc ccaagttatt atctgctcct ggggttgacc catctgtttt atgacagtta 360
 tgatattggt gtttaaaaaa aatccaaatt gttactttga tttatatgat ctaactctga 420
 atcacggaag tattactatg atgttcaaaa ctctgattga ctctacttgc tttaaaaact 480
 ctcagatccc ttctgcattt atcatcagag atcggtaaag atgacaacaa gcaggctctaa 540
 agttctgaga tgttagcaca tacccttttc acaatttagg aagctttaag atcatttagt 600
 atttttttat gttacaaaat ttggtacaat acacctctt caggaaagtc ttagtagtaa 660
 ctccaaatat tataattatt gtaaccagaa ttgtgacact tggagcagaa tgcattgcaca 720
 caaaataaaa tcctgtcaaa aatgacatc accattcccc cacaccaaat gtgtaattgg 780
 taggaaatgc atttccagtc tggtagatgg cagtgtgaca aactcctact cactcgtttt 840
 tcaagttggt gactgcagct gaaatgtttt tctgtgatgt atgccaccct tttacctatt 900
 tgatttgtaa gtgtagaatt cggattcatg tcatctccac agacctttcc tcttaggagt 960
 gcctaagctg tcttactctg atggaggat aatgtagcac gaaagacttc caaagaacca 1020
 gtttctctct tgctgttctt cttacaactc ttcacgtcta tctaaacatt ctatgcagga 1080
 gtccactaa gaaattttgg tgtaatgcca ctttgatcag ttatttgttg tatgacttca 1140
 ttcaaaaaca ctttcatcaa tagcatgggg attgtatcta tgaaagggaa gttggtgtcc 1200
 tgcgttcctc acaaaattat ccaaaggata aatgaaaag tatgtgagaa acctgcttta 1260

at 1262

<210> 121
 <211> 562
 <212> DNA
 <213> Homo sapien

<400> 121
 ggtaccaact tgagtgcctc ctaaaagtgt aaccttgggg gcggggatac agaaggatga 60
 tgctgacaat ggaatttaaa aacaaacagc aacatthttgt ggtgtctaca ggcgtggggg 120
 tggaggagct gcagcgtcac catgggaaca aaagtctccc acgcatctca ggccccagga 180
 atctttaaag agggagagtg ggcatgggag gaggacttaa gctattagtc atattttatt 240
 tcgaaaacta gatcttaagt aactgtagca aaatgttaac aattcttacc ttggaatacc 300
 ggttacatgg gattcatggt actctatthtt ttcacatggt gcaaatattht tcatatthttg 360
 acaattaaaa ctaaatagta gctthttata aaagtggcat atgcactgaa gtataatgtg 420
 ctaatthggg attcgtthta ataaaacagc thttcttaca aaaaaaaaaa aaaaaaaaaa 480
 aaaaggthtg gggaaacaag ggcaaaaggg gthccccggg ggaaatggtt accgggtcga 540
 aatthcacia ttggagaaaa ac 562

<210> 122
 <211> 695
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (13)..(13)
 <223> a, c, g or t

<400> 122
 ctggagcatg gthntgcagga gtgcaagact gcaagcctcc tccacggcca cactccagg 60
 cctggataaa gaatthcgtg catatthtcag ggaacagaat gtccccctggg gcgaaagggg 120
 atgaagthcat thtactthgta ccaactthgag thcctctctaa aagthgtaacc thgggggctg 180
 ggatacagaa ggatgathgct gacaathgaa thtaaaaaa aacagcaaca thttgthggtg 240
 thtacaggcg thggggthgga ggagctgcag cgtcaccatg ggaacaaaa thtccccagc 300
 atctcaggcc cgaggaathc thaaagaggg agagthggca thggaggagg actthagct 360
 thtagthcat thttatthcga aaactagath thaaagthact gtagcaaaath gthtaacaath 420
 cthacctthg aathaccgth acathggath atgthactct atthththcat catgthgaaath 480
 atththath thgacaatha aaacthaata gtagctthth ataaaagthg catathgact 540

gaagtataat gtgctaattt gggattcgtt taaataaaac agctttctta gaataaaaaa 600
 aaaaaaaaaa aaaaaaaggt tgggggaaac aagggcaaaa ggggttcccc gggggaaatg 660
 gttaccgggt cgaaatttca caattggaga aaaac 695

<210> 123
 <211> 386
 <212> DNA
 <213> Homo sapien

<400> 123
 aaccctggc caggcccagc tgccacaccc tttctgggag aagcatggcc tacagaatga 60
 agaggggggac caggaacccc tgtgggagag gcttagacct gaagcagtgc ccaactctggc 120
 tctcctgcc ttggctgact gggttcctgg accatgtgca tttcactggg ccatgggatc 180
 tacatctcct tgcaccccca gctggtctga tccctgccag ggccccttcc ttctctgtca 240
 tggctctcag gtggcctgat catggaaagt aaggagttag gcattacctt ctgggagtga 300
 accctgactc catcccccta ttgccaccct aaccaatcat gcaaacttct cctccctgg 360
 ggtaattcaa cagttaaag aagctt 386

<210> 124
 <211> 654
 <212> DNA
 <213> Homo sapien

<400> 124
 atgataaacc acctcagccc ccaccaagcc gccgcacccg tagaccagac cccaaggacc 60
 ctggccacca tgggcccagag agcattacct tcatctctgg ctctgctgag cggccccttg 120
 agtccccac ctgctgacct ctctggcgac cctgggtgtg ggagtgggtg cgggctgcct 180
 tctgcttccg ccgctgccgg gattgcctcc agcgtgtgg aggcctgtg cggggatgca 240
 gccctgcct gtctactgag gactcccctg aggggactgc tgaagccaac tggccaagg 300
 agcacaatgg agtgcccccc agccctgatc gtgcagcccc ccgccggcgg gatggccagg 360
 cgggctgcaa gtcaaccatg ggcagcagct tcagctacce cgatgtaag ctcaaaggca 420
 tccctgtgta tccctaccga gaggccacct cccagcccc tgatgcggac tctgctgca 480
 aggagccact ggccgatccc ccaccagcg agcacagcct gccagcacc tttgccagta 540
 gtctctgtgg ctccgaggag tactattctt tccatgagtc ggacctggac ctgccggaga 600
 tgggcagtgg ctccatgtcg agccgagaaa ttgatgtgct catcttcaag aagc 654

<210> 125

102

<211> 684

<212> DNA

<213> Homo sapien

<400> 125

acatgcagat gtgcatgtta cagagataaa gtgatcgaga caaggactga ctgggtatag 60
 aaggaagaca gactcctgtc ttcactccta aatgcagttc tttggaatca ccctactgtg 120
 atgggcgtag tagggagcca tcagctagga agaaacgtgg gagatgtgaa ttccaagagt 180
 tgcctggaca gggcaagtca tgttagcgtg ggtcacactt ccaagatatt taaagcaaat 240
 acaaaacaga acagaggatt caaacgcaa gtatgggaga tttaggccct gcagaggcag 300
 accattcctt agtatctcac aaagcagagt aatactggag gcagagtagg gggtggttgg 360
 agagcagtta gtaccaataa caatgaagtc tgtgtttgat ctgatcgata ctttccagtc 420
 ccgaatcaaa gatatggaga agcagaagaa ggagggcatt gtttgcaaag aggacaaaaa 480
 gcagtccctg tgagaacttc ctatccaggt tccggtggag gaggaggttgc ctggtgatct 540
 ctgtcctaac gatgaagact gggctattca caggcagctc tctgccctca gtggtcaggc 600
 gtgcacattt ggtctgcgcc acataacatt ctgaagcttg ggtatcatgg tcatagtgtt 660
 ccgtgtgaat gtatcgtcac atcc 684

<210> 126

<211> 2671

<212> DNA

<213> Homo sapien

<400> 126

ctgccgaaga gttcaaaaaca gaagagcaag atgcctcagg gagtatagaa tttggtgat 60
 cttttcctga tagggaatca tcatctatgg aaacatccat cgaacaaaaa gcaactgaaa 120
 cttctcacac agaggaatt actgccattg aggagagctg ggagtctatg tttaacgatg 180
 atggtgactg cctggatcca cgtcttctac aagagttatc aggaataacc aagagcagag 240
 agagcatcca ggaacctaga tctgattact acaatcatga agttcctgat attgacctca 300
 gtgattgtga attcccacat gtcattgaaa tttatgactt tccccaagaa tttcgtactg 360
 aagaccttct acgggttttc tgagttatc aaaagaaagg atttgatatt aaatgggtgg 420
 atgatacaca tgccttagga gtattctcca gtccaattac agctcgtgat gcgttgggta 480
 ttaaacacac catggtgaag attcgtccct tgtcacaggc cacaagagca gccaaaggcca 540
 aagctagagc ttatgctgag ttcctccagc cagcaaagga gcgtcctgag acttcagcag 600
 ccctagccag aaggtagtc atcagtgccc ttggggttcg aagtaagcag agcaaaaccg 660
 aacgagaagc agagctcaag aaactgcaag aagccagaga gagaaagcgg ttggaagcca 720

agcaacggga agacatctgg gaaggcagag accagtctac agtttgaaca tcactcaatg	780
aaagggataa ttccatgaat cagaaaatgt ttccatagcc ttcagataag atgatccttc	840
cagagctcta tgtacatgca gatgtgcatg ttaaagagat aaagtgatcg agacaaggac	900
tgactgggta tagaaggaag acagactcct gtcttcactc ctaaatgcag ttctttggaa	960
tcaccctact gtggtgggcg tagtagggag ccatcagcta ggaagaaacg tgggagatgt	1020
gaattccaag agttgcctgg acagggcaag tcatgttagc gtgggtcaca cttccaagat	1080
atttaaagca aatacaaaac agaacagagg attcaaaccg caagtatggg agatttaggc	1140
cctgcagagg cagaccattc cttagtatct cacaaagcag agtaatactg gaggcagagt	1200
agggggtggt tggagagcag ttagtaccaa taacaatgaa gtctgtgttt gatctgatcg	1260
atactttcca gtcccgaatc aaagatatgg agaagcagaa gaaggagggc attgtttgca	1320
aagaggacia aaagcagtcc ctgtgagaac ttcctatcca ggttccggtg gaggaggagg	1380
ttgctggtga tctctgttcc taacgatgaa gactgggctt attcacagca gctctctgcc	1440
ctcagtggtc aggcgtgcaa ttttggctct cgccacataa ccattctgaa gcttttaggc	1500
gttgagagag aagttggggg agtggttagaa ctgttcccaa ttaatgggag ctctgttgtt	1560
gagcgagaag aaaaaaaga tgaagaatga gaacgcagac aagttactta agagtgaaaa	1620
gcaaatgaag aagtctgaga aaaagagcaa gcaagagaaa gagaagagca agaagaaaaa	1680
aggaggtaaa acagaacagg atggctatca gaaaccacc aacaaacact tcacgcagag	1740
tccaagaag tcagtggccg acctgctggg gtcccttgaa ggcaaacgaa gactccttct	1800
gatcactgct cccaaggctg agaacaatat gtatgtgcaa caacgtgatg aatatctgga	1860
aagtttctgc aagatggcta ccaggaaaat ctctgtgatc accatcttcg gccctgtcaa	1920
caacagcacc atgaaaaatcg accactttca gctagataat gagaagccca tgcgagtggg	1980
ggatgatgaa gacttggtag accagcgtct catcagcgag ctgaggaaag agtacggaat	2040
gacctacaat gacttcttca tggtgctaac agatgtggat ctgagagtca agcaatacta	2100
tgaggtacca ataacaatga agtctgtgtt tgatctgatc gatactttcc agtcccgaat	2160
caaagatatg gagaagcaga agaaggaggg cattgtttgc aaagaggaca aaaagcagtc	2220
cctggagaac ttcctatcca gggtccggtg gaggaggagg ttgctggtga tctctgctcc	2280
taacgatgaa gactgggctt attcacagca gctctctgcc ctcagtggtc aggcgtgcac	2340
attggtctgg ggcgcttacc ttctgaaget taagcgtgcg cacggactgg gggcccgttc	2400
aactggcccc attaagggac cccgagataa cgagaaacgt acaccccatg gtgaaaaaca	2460

104

ccgcacaaat ccacggaccc ggagacaacc caggccaggc gcaaaaagca agaccacacg 2520
 gatatcacc aagcgagcga gaagggacca cacacacacc cgcacaacag gacacccaag 2580
 cggcgccaca acagtcacga caccacaagg ccacgaagca acacacagaa acatacacag 2640
 cagcacacgg ccatacaacc gcccacacag c 2671

<210> 127
 <211> 420
 <212> DNA
 <213> Homo sapien

<400> 127
 acgggccgca gtgttgatgg atgcggcgag gtaactctct ctcccttaag agttatgagt 60
 tatcaagagg agacttctta aagacagcaa cgcaattctt gtaacttggtg taaatagccc 120
 catctttcag agtgatacca tttctacatt tgataatgcc tgtattcctg taggatgtat 180
 atagtttagg ggattttttt ttggttgggg ttttggtttt ttagaaggtc aatatgtctg 240
 gttttattta tgtgcttgaa aaagatcatt tgaaaaaaaa aaatacattt tcaaccacaa 300
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa ggcgcggggg ggaacccggg gccagagcg 360
 ggccccgggg ggcgaaattg gttctcccgg cccacattcc cccaaaatat tggcacacag 420

<210> 128
 <211> 2269
 <212> DNA
 <213> Homo sapien

<400> 128
 taccgaggag ggaacaagct acatgctatt ttgtttgtag tattgtggaa cagtcttggt 60
 atggagtgcc agcttagagg ttgttgcaaa cttgtctaga agtgagagca tggttttttt 120
 tagccctttg agagtctaca tctaataaac attcttgctc acccataaat aacgtcaagc 180
 ctcaatgtca ccgtcacgtt gggatactct ttctcatctg gcacccctaga caggacaagg 240
 ttggttacct ttccttccat gaaccatgaa cctgtgacgg catcattcat cctgacttca 300
 ccaagctccg cctgtgggtg aggccagagc tcccactggc aatttttaga agagccagag 360
 gctccctgct tcctctagaa ataacagttc aggggtgaagc atggaggggtt tcagttccca 420
 gacaatggaa ccatttagag acaacacagt tggacatttc cactttttcc ttgattcctg 480
 gaagtccagt gggttctgca gctgaaaaag ccctgggtcc cagcagcaga gagacaggac 540
 agaggggatg cttgggcggg gagggacggt aacctgcaga acagattcca tttttataga 600
 acgagtacac gtttgctaaa acagtcctgc tttcccagac tggattccca ccacagggac 660
 agtcggaact caggactagc tccagcgaca tctttcctcc gaattcaagc cttctatcac 720

aatgtcaaaa cagctattta taaagccatt ttcattgtac ttgataacag cacgagtccc 780
aaaactttta gaaataaaaat aggacattgg cttgattgaa aagagggact ttttaaaaat 840
tgttctttcg tcagaagcct tttggatgac ttacaatagc tctgatgaag ataccacccc 900
agcgtcagtc caataggta gtagagttca acaggcattc atccctccca tgaagggatt 960
ctggatgatg gaagtttctg taatgacagg aaagcattga ccctcattga ttgtcaactt 1020
tggatattagc catgaaagac aggatgctca ttgggtgttc ttagagtgta ggaatgctgc 1080
ctattccctc ccagaacgct tgaccagggt gtgtgtgttg aggagccctg ggggaaatgg 1140
accaagtttt cccacagagc agtattaggc tgaagagcag gtgactggta ggccccagct 1200
cccatcattc cctcccaaag ccattttgtt cagttgctca tccacgctgg attccagaga 1260
gttttccaat ttgggaagcc atgagaaaagg tttttaaatac ttgggaagat ggagagaggg 1320
acataggata gttgactcca acatgacagg aagaggctgg agattgggaa ttggccatca 1380
accaagcctg tagtagtaaa gccatggtcc cgcattggaa ttacttgggg aacttataca 1440
gttctgatac ccaggctctc ctagaccagt tcaaccaatt ctaggtgggg gactcaggca 1500
tcagtgtgtt tcgtagctcc ccgggtgttt tcctgtgca gccgagcttg ggaactgccc 1560
atgctttttg gatgtcaagg cgctgttga ggctgggtgt gacagcacag agccaggttg 1620
tcttgtggaa accacagcca cgggtttgcc actggctcag catggcctca ctgccagctc 1680
cagcctggct gagggacaag atggtttctc ttgggagttc ctgagtggag cacccttcca 1740
ggctttttga aagccagctg atctgtggag ccttgttaag ggactcaata cgggtgttgg 1800
atattgatgt ttttccttga gactgtcttg tccatcaata aagatggagg atgtctcctc 1860
tttgaacccc gcttccccac cagtactctc tctcccttag agtttatgag ttattcaagg 1920
aggagacttc ttaaagacag caacgcaatt cttgtaactt gtgtaaatag ccccatcttt 1980
cagagtgata ccatttctac atttgataat gcctgtattc ctgtaggatg tatatagttt 2040
aggggatttt tttttgtttt ggttttgttt tttagaagtc aatatgtctg gttttattta 2100
ttgcttgaaa aagatcattt gaaaaaaaaa aatacatttt caaccaaaaa aaaaaaaaaa 2160
aaaaaaaaaa aaaaaaaaaa ggcggtgggg gaacccgggg cccagagcgg gccccggggg 2220
gcgaattggg ttctccggc ccacattccc ccaaaatatt ggcacacag 2269

<210> 129
<211> 750
<212> DNA
<213> Homo sapien

106

<400> 129
 gccgcccgagg caggtaacca agtttcagtt acacaggagg catgagattg atctagtgc 60
 aaaaatgatg agtataataa ataataatgc actgtatatt ttgaaattgc taaaagtaga 120
 tttaaaattg atttacatac atattttaca tatttataaa gcacatgcaa tatgttgta 180
 catgtataga atgtgcaacg atcgagtcag ggtatctgtg gtatccacca ctttgagcat 240
 ttatcgattc tatatgtcag gaacatttca agttatctgt tctagcaagg aatataaaa 300
 tacatttata tgttgactat ggcctatcta catgttgcaa ctaaacta gattttactt 360
 cctttccaac tgtgggtttg tattcattta ccacctctt ttcattccct ttctcaccca 420
 cacactatgc cgggcctcag gcataacta ttctactgtc tgtctctgta agcgattatc 480
 agttttagct tccacatatg agagaatgca tgcaaagttc tgtctttcca tgctgggtct 540
 tatttcactt aagcaaaatg acctccgctg tccatccatg ttatttatat tacccaacta 600
 gtgttcataa aactagtata tacaccacat agtataccac agaaacggac cactgcggat 660
 aaacaggatt tctggtccac acttttgtcc catacgggac cgtggggcaa tctgattacg 720
 cgcacagcaa gagcaacca gtaagaaaca 750

<210> 130
 <211> 738
 <212> DNA
 <213> Homo sapien

<400> 130
 gcgtgggtgc gcccgaggta ctgtgaatta cggatgctct ttgaaggaaa gaaatatcga 60
 ttctaattgtt cttcagaagt tctggcaggg ataagcagga catcgactgg aacgtatgct 120
 aatgaaagc agacaaattt ctattttctt acctgagcaa atattttatt gaaactgctt 180
 atgtatgcca aaggagccca caacttcagc tacacaactt tttgtattga aagaactcat 240
 actttttgta gcttttattt cacatttaac ttaaagtgac ttttagcact aaaatgccta 300
 gaagatttta ctccagacct ataaggaat gtttagtttt tatgaaaaat gacaagtcga 360
 tggtaaact tctcatgtct ttgggtgctt gccctaata gcactggaca acaccacgac 420
 cacatggaaa catatttttg gaagcaaac ttaatttta tataacgtat gctatggaga 480
 gtaagacaa ttaaggact acttgtttc tattttttt ctaataaaa tggaatccac 540
 tgtgttgaag actcttgata ttcattgtct tgtctaacca tttttgttt tataattaga 600
 ataaaatata gttgtgataa tggcatcga atggatttt tttggaaagc tacatcttat 660
 ttgtgaaatg ttttttaaaa tcagagtaac tatcaactga ttcagctttt tgttgtttt 720
 ttcttggtat aatacttg 738

<210> 131
 <211> 1875
 <212> DNA
 <213> Homo sapien

<400> 131
 tggcaacgat ctggaccgct acaaccgct aagctccagc gccttgtgcg caacgcgctg 60
 gcgcacgtgg tgccaaggag cgcgagctga gctggcgcac tcggagagtt tcgccgcctg 120
 tgccgctacg gcaagcgcga gttcaagatc ggcggcgagc tgcgcatcgg caagcagccc 180
 taccggctgc agattcagct gtcggcgagc cgcagccaca cgctcgagtt ccagagtcta 240
 gaggacctga tcatgggaga agcgacgcaa cgaccagat cgggcgcgcg gccctgctg 300
 caggagctcg ccacgcacct gcaaccggcg gagccggagg agggcgacag caacgtggcg 360
 cggactacgc cgctcccgg gcgccccct gcgcccagct ccgaggagga ggacggagag 420
 gcagtggcac actgatgggc gagctgagcg cagagctgcg aaggggaact gtttgagta 480
 gcagccgctg ctccctttct ccctctcttc ctccctcttt tgccactgtc tgggccccat 540
 ctgggattcc tgggcccttt ggaaaagagt tggtgaaatg cgcagccggc tgtggacggg 600
 ggaggaggaa ggggacagag ggagcaggaa taagactgta gaactgtttt gtactgtgaa 660
 ttacggatgc tctttgaagg aaagaaatat cgattctaata gttcttcaga agttctggca 720
 gggataagca ggacatcgac tggaaactg gctaaatgaa agcagacaaa tttctatttt 780
 cttactgag caaatatttt gttgaaactg cttatgtatg tcaaaggagc ccacaacttc 840
 agctacacaa ctttttgtat tgaaagaact catacttttt gtagctttta tttcacattt 900
 aatttaaagt gacttttagc actaaaatgc ctagaagatt ttactccaga cctataagga 960
 aatgtttagt ttttatgaaa aatgacaagt cgatggtaa acttctcatg tctttggctg 1020
 tttggcccta atagcactgg acaacaccac gaccacatgg aaacatattt ttggaagcaa 1080
 aactttaatt ttatataacg tatgctatgg agagctaaga caatttaagg actacttggt 1140
 ttctattttt tttcttaata aaatggaatc cactgtgttg aagactcttg atatcatgtg 1200
 cttgtctaac cattttttgt tttataaatt agaataaaat atagttgtga taatggctcat 1260
 cgaatggatt tgtttgaaa gctacatctt atttgtgaaa tgttttttaa atcagagtaa 1320
 ctatcaactg attcagcttt ttgtgtttt gttcttggtc ataatacttg tgactcatga 1380
 agaattatgt tgacaaacag gataaattcc acatgcattt tatttcccag tgagttgtat 1440
 aaactttatt tttgttgaag gttgtatgtt aaatcaatgt tacattctta tatcacttct 1500
 tgagaaggaa gttccgattt gaaattgtat catttccttc aaaatgaagg gcagtgttta 1560

108

gttaaataaa agattgatga tatcttttaa gccaaaaaaa aaaaaaaaaa aaaaaaaaaa 1620
 aaaaaaacga accaaaccaa taaaaacaag aagcacacag accgaacacc acacacacaa 1680
 gccaccagag ctacataac gcgcgggcaa acatccacac ggccacacac agcaaccac 1740
 tatgagagcc accccgcgga acaaaagacc ccacacacaa ccagagacaa gaaacctgcg 1800
 agccacgccc tccacacca caaccacgaa tagtcacctc agtaacaaaa caaacacaga 1860
 cggaggcgcc gacaa 1875

<210> 132
 <211> 828
 <212> DNA
 <213> Homo sapien

<400> 132
 tggtcgcggc cgaggtacaa taggtctctt gaatttatcc ctctgtctta attgaaattt 60
 gtatcccttg accaacatct tcccagtcac acccccatcc ctctggtaac catcattcta 120
 ctctagttgt atgagttcaa tttttttaga ttccatttat aagtgattta attaatatct 180
 ttatcctctt tccagataat tcaaggacct tagcatttta actctagtca actgtaatat 240
 tacattccat cgtattgcag tattttagtc ttctctatt aagccttcca aattggatat 300
 tagcattatt gtggttgttt cacattagca ttattgtggt tgtttcagat agtcaatatt 360
 gatgcagatt tacctgaata ttacccatga ttaccatcat tccttcttc tacttagatt 420
 tccatcatcc ttcttcttga aatataattt taaaaggtc cattgaagaa gttctgttga 480
 tggtaaatac agttttactt tctttgaaaa tatctttatt ttgccacat cagttatttt 540
 attgttcagt attaagaaaa cctaattcct gtgttttctt cccatcattg ttgatattga 600
 gttgtgtgcc atcaggcaaa tgtcattact ttttagatat tctaaacctg ttgtttcttt 660
 aagtaagtac attgtctccc ccttaatctg ttctccttcg taatgtttta ttatttctct 720
 cactattatg gattctggac aggtttcttc tgggtccttc tttcaggttg ctattctcta 780
 ttcaggtgtg tttatctgct atttatcctc cctccagttt tttccttg 828

<210> 133
 <211> 1023
 <212> DNA
 <213> Homo sapien

<400> 133
 tggtcgcggc cgaggtacaa taggtctctt gaatttatcc ctctgtctta attgaaattt 60
 gtatcccttg accaacatct tcccagtcac acccccatcc ctctggtaac catcattcta 120

109

ctctagttgt atgagttcaa tttttttaga ttccatttat aagtgattta attaatatct 180
 ttatcctctt tccagataat tcaaggacct tagcatttta actctagtca actgtaatat 240
 tacattccat cgtattgcag tatttttagtc ttcttctatt aagccttcca aattggatat 300
 tagcattatt gtggttgttt cacattagca ttatttggtt tgtttcagat agtcaatatt 360
 gatgcagatt tacctgaata ttacccatgg attaccatgc attccttctt tctacttaga 420
 tttccatcat ccttcttctt gaaatataat ttttaaagg tccattgaag aagtgtctgt 480
 tgatggtaaa tacagtttta ctttctggtt aaaatatctt tattttgccc acatcagtta 540
 ttttattggt cagtgattaa gaaaacctaa ttctctggtt ttcttcccat cattgttgat 600
 attgagttgt gtgccatcag gcaaatgtca ttacttttta gatattctaa actgttggtt 660
 gctttaagta agtacattgt gctcccctta atctgttctc ttcgtaatgt tttatttatt 720
 tgtctcacta taatgaattc tggacaggtt tcttctggtc tttctttgca gtttgctaat 780
 tctctattca gctgtatcta atctgctatt taattcatcc atcaagtatt ttttccctag 840
 tattttggtt taataatttt atttactatt tctagatttt tttctaataca tcttggctct 900
 tgtcatagta tcttcttctt tatatacatt ttatttatgt atctgataac attaataact 960
 taaacctttg taagttataa gtagtgtttt agttttggtg ctgatttggg tcaaataaac 1020
 ata 1023

<210> 134
 <211> 757
 <212> DNA
 <213> Homo sapien

<400> 134
 gagcggcgcc egggcaggta ccttcgtgcc cctcagtagt tgttttagcc taatgtagag 60
 tcaatctagg acttataatt attcatcatg attttgagta gattgtaatc atcaagaatt 120
 tttcatagat cgtttacttc caattgaatt tagctcagaa gtgattgctt tctctttatt 180
 tgagatagga gctctcgcac tgtcgccagg ctaggagtgc aagcggcat gatcgtcgcc 240
 tcaactagcaa cctctgcctc cggggttgaa gcagatatac ccctgacctc aagcctcctg 300
 cagtagctag ggactacagg tagttcatcg cttgtcctta gcttggaac taggatgcac 360
 aaacacatgg gttattatac tcgtacacgg agctggtcac acaacggaac tagactctct 420
 ctccaatgt gataccacac agacaacct cagaactacc ttcgagcctt acttaagatc 480
 atcccttcac tgatctaaca aacttataaa cattaataca accagatact gcgtctcgac 540
 tattgcacgg caaatcaaaa tacaacagg tctccactaa agaccagggtg gtgacatgct 600

110

ctagagatca acagaacaat ctaatcctga ccctcacgcc aactatgatg acacgatggc 660
 cgctggccca cacaggaagg cggacacggg ccgcgctcaa agaccaccca tgtccggacc 720
 tagcctaaaa aaaactcacg ccccgccgcc cctaact 757

<210> 135
 <211> 1513
 <212> DNA
 <213> Homo sapien

<400> 135
 gggggagcct gggcgccgag ccgggtgtga gctgcctgaa aatgcactcg gatgccgccg 60
 ctgtcaattt tcagctgaac tctcatctct caaactggc aaatattcat aagatctacc 120
 acaccttaa taagctgaac ctaacagaag acattggcca agacgatcac caaacaggaa 180
 gtctgcggtc ttgcagttct tcagactgct ttaataaagt gatgccacca aggaaaaaga 240
 gaagacctgc ctctggagat gatttatctg ccaagaaaag tagacatgat agcatgtata 300
 gaaaatatga ttcgactaga ataaagactg aagaagaagc cttttcaagt aaaaggtgct 360
 tggaatggtt ctatgaatat gcaggaactg atgatgttgt aggcctgaa ggcatggaga 420
 aattttgtga agacattggt gttgaaccag aaaacgtgag tcaaacttac tgagtgggt 480
 gaatcagttg gttgtttttc atacttaaat ctttgttctt tagcaaataa atagaataat 540
 taaaaagtag tggtagttaa gtttttatga agcagtctaa gaaataagtt ctaattctag 600
 tttgacttat aagcagattc tccattcttg taagtatat ggtgtaacta cagttatfff 660
 ttctctcatt taatttcttg tatgtaaaag gtacagtaag ccagatgctt acaaaatggt 720
 gtggccacat gtgcctacaa tgacggatca actggaggcc acattgtacg ctgtgtacct 780
 tcgtgccct cagtagttgt tttagcctaa tgtagagtca atctaggact tataattatt 840
 catcatgatt ttgagtagat tgtaatcatc aagaatffff catagatcgt ttacttccaa 900
 ttgaatttag ctcagaagtg attgcttttt tttttttgag ataggagctc tcgactgtc 960
 gccaggctag gagtgaagc ggtcatgatc gtcggtcac tagcaacctc tgcctcccgg 1020
 gttgaagcag atataccct gacctcaagc ctctgcagt agctagggac tacaggtagt 1080
 tcatcgcttg tccttagctt ggaaactagg atgcacaaac acatgggtta ttatactcgt 1140
 acacggagct ggtcacacaa cggaaactaga ctctctctcc aaatgtgata ccacacagac 1200
 aacactcaga actaccttcg agccttactt aagatcatcc cttcactgat ctaacaaact 1260
 tacaacatt aatacaacca gatactgcgt ctcgactatt gcacggcaaa tcaaaatata 1320
 acaggttctc cactaaagac caggtggtga catgtcctag agatcaacag aacaactata 1380

111

tectgaccct cacgccaact atgatgacac gatggccgct ggcccacaca ggaaggccga 1440
 cacgggccgc gctcaaagac cacccatgtc cggacctagc ctaaaaaaaa ctcacgcccc 1500
 gccgccccta cct 1513

<210> 136
 <211> 738
 <212> DNA
 <213> Homo sapien

<400> 136
 gcgtggtcgc ggcgaggtac caaccccagc acacccaac agcctttcct cggccccctc 60
 ctcaggcctc ctaattactc tttctcagcc tggagtgtgg ggccgttacc gtcctcttcc 120
 cccttctcct tccatactgc acttaacctt gctggaagat ttaatgatgg agatttaggg 180
 caactgtggc tgcttgggac ccttccctgg gaccaaagga acttaaaacc caatacctga 240
 cactggaatg aaatccaagt ttttaaatat caccttcaa tcaactcacag atctcacatc 300
 tatcttaaaa tactcagcct cactccttaa ctgagtgtct gcctgagagg gagaaaagtt 360
 ccattttaaa aacgtattca ctttactgat tactgtgcaa tttgaattaa gtcacgattc 420
 tttagtcatg gaggtcgaga atctcagatt caaattgtca gagaccatga tttagaagtc 480
 taccaaacac ccagtttcct tccactgttt tagggtaaca ggaaaacatg agattggggg 540
 ggtgtccgct attaaatgga accacacatc atgaaattca attctcatgt taagacattc 600
 tgtattgtgg gatgtcaaaa gtatctccca aaactttcgt ttgacctgtc agagtgggga 660
 tggttactcc ctatacttca gtttgtttca caagcttggc gtaaccaggc atagtgttcc 720
 gtgtgaatgt tcgtccac 738

<210> 137
 <211> 1350
 <212> DNA
 <213> Homo sapien

<400> 137
 atggttatgg agaagcccag tccgctgctt gtagggcggg agtttgtgag gcaatattat 60
 actttgctga ataaagctcc ggaatattta cacaggtttt atggcaggaa ttcttcctat 120
 gttcatggtg gagtagatgc tagtggaag cccagggaag ctgtttatgg ccaaatgat 180
 atacaccaca aagtattatc tctgaacttc agtgaatgtc atactaaaat tcgtcatgtg 240
 gatgctcatg caaccttgag tgatggagta gttgtccagg tcatggggtt gctgtctaac 300
 agtggacaac cagaaagaaa gtttatgcaa acctttgttc tggctcctga aggatctggt 360
 ccaataaat tttatgttca caatgatatg tttcgttatg aagatgaagt gtttggatgat 420

112

tctgagcctg aacttgatga agaatcagaa gatgaagtag aagaggaaca agaagaaaga 480
 caaccatctc ctgaacctgt gcaagaaaat gctaacagtg gttactatga agctcaccoct 540
 gtgactaatg gcatagagga gcctttggaa gaatcctctc atgaacctga acctgagcca 600
 gaatctgaaa caaagactga agagctgaaa ccacaagtgg aggagaagaa cttagaagaa 660
 ctagaggaga aatctactac tcctcctccg gcagaacctg tttctctgcc acaagaacca 720
 ccaaagccaa gagtcgaagc taaaccagaa gttcaatctc agccacctcg tgtgctgtaa 780
 caacgacctc gagaacgacc tggttttcct cctagaggac caagaccagg cagaggagat 840
 atggaacaga atgactctga caaccgtaga ataattcgct atccagatag tcatcaactt 900
 tttgttggtg acttgccaca tgatattgat gaaaatgagc taaaggaatt cttcatgagt 960
 tttgaaacg ttgtggaact tcgcatcaat accaaggggtg ttgggggaaa gcttccaaat 1020
 tttggttttg tggtttttga tgactctgaa ccagttcaga gaatcttaat tgcaaaaccg 1080
 attatgtttc gaggggaagt acgttttaaat gtggaagaga aaaaaacaag agctgcaaga 1140
 gagcgagaaa ccagaggtgg tggatgatgat cgcagggata ttaggcgcaa tgatcgaggat 1200
 cccggtggtc cacgtggaat tgtgggtggt ggaatgatgc gtgatcgtga tggaagagga 1260
 cctcctccaa ggggtggcat ggcacagaaa cttggctctg gaagaggaac cgggcaaatg 1320
 gagggccgct tcacaggaca gcgtcgtga 1350

<210> 138
 <211> 569
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (509)..(509)
 <223> a, c, g or t

<400> 138
 cgcccgggca ggtcgcccat gtgctgtgat gtcagtgagc gggcggagt caggctggtc 60
 agtgccaggt gtccttctc ccaccgaga acagtggcca ggttgctcct caggcacctc 120
 gggcaactgc cccttcctt ccagtggggc ctgacctggc taccgagctt ggcagctaat 180
 aggcggggccc ctcagcattc acgctcctga gctgctttat caaactagga ttgttcccc 240
 aggtctaaga aaaccatcca ttcactgcaa agttagtat tactgcggat gggctaggag 300
 ttagaggaag agagtgactc aaatcacaac acctcctgga cgaagctgga agcggattaa 360
 aataccgggc ctaatttcag aacaacaaaa aaaaaagaaa aaaaaaaaaa agcgcggggc 420

113

ggaacccagg ggccaaaagg gtgggtcccg gggggggaaa tctggttacc gcggcccaaa 480
 attcccaaaa aaatttgggg gggccaaang caccgcgctc tctgcccccc ccacgcccgc 540
 ccccccccc acaaccatc gccgccccg 569

<210> 139
 <211> 739
 <212> DNA
 <213> Homo sapien

<400> 139
 tatatcacta taggggactg ggtcctctag atgctgctcg agcggccgca gtgtgatgga 60
 tccgggcagg tactgcctgg ttttacaaga attaatgcag tttcacagtg aagcatgtaa 120
 gatattgaat tttagagaca atagaccaga tacctttcta atctcatttt attcattaat 180
 gtcaaataat accattttta aaaatatggt gcttatttgt ctagcaagta acctatagaa 240
 aagtattatt ttatacaaaa agatgattag gtcacataaa ggaattggaa tcttaagttt 300
 aaaatacact tctgttttta gccagaaggg agaaacgatg gttggattta tgccattttt 360
 caattaaaaa ccatgtggta ctacttgaag cagtttctga gtaaattggag gtgtttaaag 420
 atttgtatta ttctctccca atgactagat agtagtattt tacaattggag acttaaaagt 480
 tttttgtggt ttattctttc gcttttctat gccctcaatc caaagaacac cagaaataca 540
 cttgtagtcg gaaaacttgg gtttatcact cgcacaaagg aatgacacac accatgggcc 600
 actctggagc ctctcaataa aaggatgttt caaaggaaca acaacaaaaa aaaaaaaaaa 660
 aaaacgttgg gggaaacaca gggcacaag tgtcccgggg gaaattgttt tccgccacaa 720
 tcmetaattc acaaaaacc 739

<210> 140
 <211> 1131
 <212> DNA
 <213> Homo sapien

<400> 140
 aagttgatag tatatccacc acctccagct aaggagggca tctctgttac caatgaggac 60
 ctgcactgtc taaatgaagg agaattttta aatgatgtta ttatagactt ttatttgaaa 120
 tacttgggtgc ttgaaaaact gaagaaggaa gacgctgacc gaattcatat attcagttct 180
 tttttctata aacgccttaa tcagagagag aggagaaatc atgaaacaac taatctgtca 240
 atacagcaaa aacggcatgg gagagtaaaa acatggaccc ggcacgtaga tatttttgag 300
 aaggatttta tttttgtacc ccttaatgaa gcgtgagtaa gaatttcctt taaaggaaaa 360

114

tctttaaatc atgtaaatga tgacaatddd taaataatga gtatgaggtg aagaattcat 420
 tttaaaacat ctttctgaaa tctcttgtgt atattcatat ttgtactgcc tgttttacia 480
 gaattaatgc agtttcacag tgaagcatgt aagatattga attttagaga caatagacca 540
 gatacctttc taatctcatt ttattcatta atgtcaaata ataccatddd taaaaaatg 600
 gtgcttattt gtctagcaag taacctatag aaaagtatta tttatacaa aaagatgatt 660
 aggtcacata aaggaattgg aatcttaagt ttaaaatata cttctgtttt tagccagaag 720
 ggagaaacga tgggtggatt tatgccattt ttcaattaaa aaccatgtgg tactacttga 780
 agcagtttct gagtaaattg aggtgtttaa agatttgtat tattctctcc caatgactag 840
 atagtagtat ttacaattgg agacttaaaa gttttttgtg tttattctt tcgcttttct 900
 atgcctcaa tccaaagaac accagaaata cacttgtagt cggaaaactt gggtttatca 960
 cttgcatcaa ggaatgacac acaccatggg ccactctgga gcctctcaat aaaaggatgt 1020
 ttcaaaggaa caacaacaaa aaaaaaaaaa aaaaaacgtt gggggaaca cagggcacia 1080
 agtgtcccgg gggaaattgt tttccgccac aatccaaat tcacaaaaac c 1131

<210> 141

<211> 887

<212> DNA

<213> Homo sapien

<400> 141

gcgtggccgc ggccgaggta cactgaatta ttcacagtaa tcgcttggtt ggggaaaggg 60
 ttagtaaatg ccaaaggaaa taccacaga aatctctac acagcttaga tgttgtgctg 120
 gcatttaagg cccatgagtg atggtccatt ctgcagcttt tcatgccatg cctttccttt 180
 gtgtgggggt ccacagatca gagtctgtct gtggcatcga ctctctatg tcctcattgt 240
 tcccacccat tgctgggatg tccacgttgg acttctcaaa agtggcccaa gaatctaagt 300
 gcaaaatctg tttggatttt tacaatddd tctaatctt ttacagtctt ggtcattcct 360
 attcaactg caattdtttt caatgacttg cctggtgtga atattdtttt aaagcatcca 420
 gtattaaaca aaaaaattta aacagctaaa aaaaaaaaaa aaaaaacaaa cggctgggcy 480
 aaaccagggc tcaataccgg ctccccgtgg tgctgaacac tggataactc cgcggtcac 540
 caattcccaa ccacaacata cgggcgagac aaggctgcac gcaaccggc acgcgcatgt 600
 cgcaggacac gtacggagc caagaacggg cagcaggacc acagagaacc agacgcaggc 660
 cgcgcacgtg gagcggaggg gtagaacga cagccgccgc gccgtgggca ggggccatgg 720
 cgcacacggg ccgacacgga agcggagccg cagcgacagc gagcagcacg cggggcgacg 780

115

gcgcggcgag gaggggagcg gcgcggggaa cggacgctgc agagaggcgg agggcggcga 840
 gccgcggcgc ggccgagccg aaggcgaccg caagcggcgg cggcggc 887

<210> 142
 <211> 2086
 <212> DNA
 <213> Homo sapien

<400> 142
 cgagccaaga attcggcacg aaaaacaaat acttcctgat cgatcccttg tcttgtttag 60
 tatgcttcct gaccattttt tacctaaca tttgtgttct tttcccgaga aggaaaatca 120
 acttctatcc tatctctacc cagcagaggc ccctgccccca ctttacacac aaaaccatct 180
 aactttttga tattctaaat gggggaaacc cctattttat aaccctcggg tacttttaat 240
 ctttagatga ggaactagag gagccactat gttcctctca gcaccatgat ttatgcctta 300
 gctaaggcct tcacttgggg aaggggaaga aggttgtttt caagcctgtg gcctcctgtc 360
 actcccacc cctggaaggc ccttcacttt tgggtgatgc ctagaggcct catggacagc 420
 agtcccttct gacaccaggt gagatatcat ctgggagggt cgcagccctc agttccctc 480
 atggctctct ctttcacttc cctccatgac accacctcat cgagttgaag atgttattga 540
 tgagtgcagt ggggtgatag tgcctccca aaattcatgt ccaccagaa attcagaatg 600
 caaccottatc tggaaataga atctttgcaa atgtgattag ttaagatgaa atcatactga 660
 gttaggatga acctgaaatc caatcactgg tgccttgta agaggaaagg tcacaaagag 720
 acagaggaga tacacagagg agcccatgta atgatgggta cggagactga cgtggcacia 780
 ctataagcca agaatgcca gggaggcca gctagcagaa gctagggaaa aacacagagg 840
 gattctcccc tggagccttt ggaggagtg tggccctgct gacaccttgg ttctggactt 900
 ctggcccca gaactgtgag aaaataaatt tctgtggttt aagccacaca gtttgtggtg 960
 ctctgacttc gtgagctttt ctgccatct gacagcctc gctgccttc ctccctgccc 1020
 accgtcctcc cgcccgtcc cagacctcc tcgctcctca tcccactcca ctctgtgag 1080
 tgctcctcca caccatggct gcaatcccca ccttaagctg gggactccca aaccccgact 1140
 tccccacagg gctcaggagg ctttctcca gccagcctca catttgact catgcttctc 1200
 ccccatgcca ccctcagcta cgctgaatta ttcacagtaa tcgcttggtt ggggaaaagg 1260
 ttagtaaatg ccaaaggaaa taccacaga aatctcctac acagcttaga tgttgctg 1320
 gcatttaagg cccatgagtg atggccatt ctgcagcttt tcatgccatg ccttctctt 1380
 gtgtgggggt ccacagatca gagtctgtct gtggcatcga cttccttatg tcctcattgt 1440

116

tcccacccat tgctgggatg tccacgttgg acttctcaaa agtggccaag aatctaagtg 1500
 caaaatctgt ttggattttt acaatttttt cctaattcttt tacagtcttg gtcattccta 1560
 tttcaactgc aatttttttc aatgacttgc ctggtgtgaa tattttttta aagcatccag 1620
 tattaacaa aaaaatttaa acagctaaaa aaaaaaaca aaaaacaaac ggctggcgga 1680
 aaccagggct caataccggc tccccgtggt gctgaacact ggtatactcc gcggttcacc 1740
 aattcccaac cacaacatac gggcgagaca aggctgcacg caaccoggca cgcgcatgtc 1800
 gcaggacacg tcacggagcc aagaacgggc agcaggacca cagagaacca gacgcaggcc 1860
 gcgcacgtgg agcggagggg tagaacggac agccgcccg cegtgggcag cggccatggc 1920
 gcacacgggc cgacacggaa gcggagccgc agcgacagcg agcagcacgc ggggcgacgg 1980
 cgcgccgagg aggggagcgg cgcggggaac ggacgctgca gagaggcgga gggcgccgag 2040
 ccgcgccgcg gccgagccga aggcgaccgc aagcggcggc ggcggc 2086

<210> 143
 <211> 676
 <212> DNA
 <213> Homo sapien

<400> 143
 gccgccgggc aggtactaaa taaaatgcaa aacatgtcac atcactcttc ttcattgggtt 60
 catgtcctct gtgggtcagg tcttccacat gtagagtaga ggtagggtat gttcacacct 120
 tcaatgacaa cctacacatt tctgctcaa caggtccaaa attgttccta ggtttcaaaag 180
 ttgttgtttg tttgtttttt tcctttttct tttttttttt tttttttgga gaagtggagt 240
 ttggctctgg ttggccccgg tgtggagtgt gcaagggcg gtgatctgcg gttcaccaac 300
 aaacctcgtg gtcctccgcg gttacaagg gcgattatc cgtggcctac aggcctcgcg 360
 agtatagccg tgggatataa tagggcagtg gcgcacacca gtgcccgagc ttaatttgtg 420
 ggtattttaa ggtagaagaa gcggggttct ctccccctt tgtgtgggtc tcgaggcgt 480
 ggactctggg aggcctcgcg tggaaccctc gaggggtgat ctcacacctg tgcgcttggg 540
 ggccttccca caaaagggtg gcctgggggg atttaccagg gcgtggcaga agcccaaact 600
 atgtggccg gggcgcacac aggggggttt cccaaaaggg tttttttaac cggattataa 660
 aagagggttt cgctag 676

<210> 144
 <211> 1260
 <212> DNA
 <213> Homo sapien

117

<400> 144
 taaacataca cacatcaaaa ataactcagc cacatgcaac aatacagaga atcttaaaga 60
 catagtatga gcaaaataat cagtacacac aaaattccac ttatacaaag ctcaaaaaca 120
 aaattaagca atatTTTTTTA gaaatgcact tataaatgat gactgacca ctatcaagga 180
 aagtatttaa cattgctctg aaagtcttg aaattcttga ttttccttct tcaatttcta 240
 caccatcac cagccccagt cttcccaac tactaaaca gcaccgtcat ccatttagca 300
 tttcaagcca gtgagaagtc atccttaatt ctgcttttct attaatttcc ctacttctaa 360
 tctattacgt gtcttattag atctaagatc aatatatttc ctgaatatgt ctatttatgt 420
 ccatttcaa cactaccact gaagtctaag ccattgtcac ctttcttctt ggattactgc 480
 aatagcctca cagcttccac tcttgaccac atacactcca ttctgcactc agccctcata 540
 gtgatcatta taaaggataa aatggtgtgg ccagttagct cagttggtta gatcatggta 600
 ctaataaaat gcaaaacatg tcacatcact cttcttcatg gttcatgtc ctctgtgggt 660
 caggtcttcc acatgtagag tagaggtagg gtatgttcc accttcaatg acaactacac 720
 atttctgctc caacaggctc aaaattgttc ctaggtttca aagttgttgt ttgtttgttt 780
 ttttcctttt tctttttttt tttttttttt ggagaagtgg agttttggct ctggttggcc 840
 caggcggtga gtgtgcaagt ggcggtgatc tgcggttcc caacaaacct cgtggtcctc 900
 cgcggtttac aagggcgatt attccgtggc ctacaggcct cgcgagtata gccgtgggat 960
 ataatagggc agtggcgcac accagtgcc gagcttaatt tgtgggtatt ttaaggtaga 1020
 agaagcgggg ttctctcccc cctttgtgtg ggtctcgagg gcgtggactc tgggaggcct 1080
 cgcgtggaac cctcgagggg tgatctcaca cctgtgcgct tgggggcctt cccacaaaag 1140
 gtgggcctgg ggggatttac cagggcgtgg cagaagccca aactatgtgg gccggggcgc 1200
 acacaggggg gtttcccaa agggtttttt taaccggtat taaaagagg gtttcgctag 1260

<210> 145

<211> 433

<212> DNA

<213> Homo sapien

<400> 145

cggccgccgg gcaggtactg gtggttgggt tcattagtgg atcacacaca gggttgtact 60
 tggcttghaa aatggtgcct cggatagggg gagtttggat aagtatgat gtatgtatga 120
 gttatagcaa aattaagtag attgaatcaa gtccatgcaa aagcagtaaa acagttatta 180
 attgttaatt ttttaaaaat taaaacgtta ataaaacagt ttgtaatgtt ttgctagtgt 240
 cttttataaa atgatgtaag ttacagtgga agtcttcaca ggacttgtgt ctttcctgga 300

118

actattgaaa tgtaatttag gatgatttga tcttccatct caagttgtca acatggctgt 360
 gtcattctgg cttacatatg ttttatttaa caaaattcta gtcaagggat aaggccttaa 420
 tgaagacaag ctt 433

<210> 146
 <211> 1791
 <212> DNA
 <213> Homo sapien

<400> 146
 ggaatgaaca aacaaacaaa aatccttgct ctctgggtgc ttacatttta gttgggagag 60
 ggacaaacaa gataagggaa atacatacct tagttaagaa caagtgccac agaggaaaag 120
 ccaggctgag gcagtgggtg tgaacatctt atacagggat gtccagaatc agggccttga 180
 agaaagccct gaaggcagcg tgtaccgagc aggaatgccc tgtggaggct gagcatttag 240
 gaagtgggaa cagccgggtc ggaggctctg gagggtgagg ggtgtcaaga aggccagcat 300
 ggctggagca gaaagcaggg cggggagggtg ggggaccagc tcacagggtc ctagagccag 360
 aatgagaagg gcttcttggc tggattacag gcgtgagcca ctggaacctg gccttgtttt 420
 gctttatttt ttctcttaca tgaagtaaag cgctttggtc aaacacacaa aaatactgcc 480
 ttgtactggt ggttggtttc attagtggat cacacacagt gttctacttg gcttgtaaaa 540
 tggtgccctg gataggggtga gtttggataa gtatgtatgt atgtatgagt tatagcaaaa 600
 ttaagtagat tgaatcaagt ccatgcaaaa gcaataaaac agttttaatt ttttaatttt 660
 ttaaaaatta aaactttaat aaaacagttt ttaatttttt gctaggttct tttaaaaaat 720
 gatgtaactt acatggaagt cttcacagga ctttttctt tcctggaact attgaaatgt 780
 aatttaggat gatttgatct tccatctcaa gttgtcaaca tggctgtgtc attctggctt 840
 acatatgttt tatttaacaa aattctagtc aagggataag ggcataatga agacaagctt 900
 cagttatgaa agtacaaact atttgtgtga ttaattttta aaaatgacat taagaagccc 960
 attgtaaaat aatatttgca gtcaaatggt ttttcttgct gtaagtcctg ttgtagctat 1020
 gtttagggta gtggttctca tctaccttgg agtgcataag acttacctag caggcttgtt 1080
 taaaaagttc agattcctag ctttgtaacc agggattgcc tcagggtgga tgggctgtgg 1140
 tcctggagtc atcactttta taaatagtgg ttcagagacc acagagagag actgcttcat 1200
 cgaatgggaa gtaccaagga gaaagtacaa ttcagtattg tctggaggca agtggacact 1260
 ttgtacctga ggtttagaat aggtggtctc ttgccagtac aatccccagg cgttttctgt 1320
 gttcagaagt agtaagaatg cctttaatc agaggattat ctaagctctt taaagctgtt 1380

119

tttctccatt tgtcatagtg ccttctctga aaaatgaatg tacaggatc ctatctteta 1440
 atgtaattag gattttttaa aagcaatctt gatagttttt cttttaaaaa gtaaaattca 1500
 gcaactgtgac ttgaaccccc aaatctttca catacaggtg aacattaag ccacaaataa 1560
 atatgacaga aagaagaaaa gatcctattc ctgtcattag ggactagtac ccattaactt 1620
 gaaccgactc ggcaagggtg caacatttct tggcacatcg tgcacacact atgttttgac 1680
 acgaggactt cccacttata aacaccggac cggggaatat ttcacatcgt ttaagtaatg 1740
 caccocgggc aaaaaggaga aaccctcatt caaaaaatct atcgccgtct a 1791

<210> 147
 <211> 349
 <212> DNA
 <213> Homo sapien

<400> 147
 ggaatgatcg atactatagg ggcattggtc atctaataca tgctcgagcg cgcgccagtt 60
 gtgatggatg cgtggctcgc gccgaggctc acgttttagct gactataatt ttaaatagcc 120
 ctatgtgaca agtggctact ttattggaca gtgtagatct aagattaatt cctcaactgt 180
 tttgcaactc acaaagacat acctctgagt tggcaaccag cagggtggat aacggggcag 240
 tggtgataaa atcaaagaat aggtaatgaa acaatcatcc agttaacaat cagcaagggt 300
 cttcagagcc taattaatgt ttaattctaa ataaattgca acaattaag 349

<210> 148
 <211> 848
 <212> DNA
 <213> Homo sapien

<400> 148
 agctgggatt acagacgccc accaccacac ccagctaatt tttgtatctt tagtagagat 60
 ggggtttcac catgttgacc gggctggctc tgaactccta acctcgtgat cctcctgctt 120
 cagcctccca aagtgcaggg attacagggtg tgagccactg cgcgccacat cccatttaac 180
 tttctgtctc tgtgactctg atgactctag gaacctcata taagtggaat aatataggat 240
 ttattctttt ttaaaaaatt tattttgaga tggagtctca ctctgtcact caggctggag 300
 tgcagtgact cgatctcggc tcaactgcaac ctccgccttc ctggcttaag caatttttgt 360
 gcctcagcct cccaagtatc tgagattaca ggcgtgtgcc accacacca gctatctttt 420
 attttttatt ttttagtagaa gatgggggtt cgccatgttg gccggactgg tctggaactc 480
 ctggcctcaa gtggctctcc cacctcggcc tctcaaagtg ctgggattac aggcgtgagc 540

120

caccacgttt agctgagtat aattttaaat agccctatgt gacaagtggc tactttattg 600
 gacagtgtag atctaagatt aattcctcaa ctgttttgca ctcaacaaag acatacctct 660
 gagttggcaa ccagcagggg ggataacggg ccagtggtga taaaatcaaa gaataggtaa 720
 tgaacaatc atccagttaa caatcagcaa ggttcttcag agcctaatta atgtttaatt 780
 ctaataaat tgcaacaatt aagaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 840
 actcggtc 848

<210> 149
 <211> 414
 <212> DNA
 <213> Homo sapien

<400> 149
 cagtggtagc cgcgacgcag gtaccacagc tcccagtgcc cattacctct atcatggatg 60
 ctgggtgact ttgggaagtc accacctctt cccaagcctg tttccatat cacagatgtg 120
 gggccatggc ctcgatgatg gtctccacag gtctttccac ctctgtgagt ccaagtcagg 180
 tcaatcagca aggaccaat ctctgaccct gggtcagctc ctcagaacca acccccagca 240
 tctctaaagc aaaagcctca cctcaagggc tgctcagaag agagcacctt cagcatgagt 300
 tgttgctgga aggatctaat aagctgtggt tctctgggaag tgggtgcttta cttagccctg 360
 tggacaactt ctctatgcat ctgtgtgagc agatgatcat tgtattacct tttta 414

<210> 150
 <211> 2088
 <212> DNA
 <213> Homo sapien

<400> 150
 ggtggcagtg atacatgttg gcaggctggt cttgatcctg actcaagtga tccgccgctt 60
 caacctcca aggtgctggg attacagggtg tgagccacca cacttgtgca gtatatcctc 120
 agtatgaaga tatttttata ttctctgtgt ctctggcttc agagtttcac ctgccacac 180
 agggtcctgt gctggcaact ggacttcccc ataagccttg ggtatcctgt gatgggctgt 240
 gtctccctga agattgtctg gcttgccac ttctccgtgc atgactctgg gtgtgagtct 300
 gtctaggaac aggagggaaa gttggactca gacagaaatc agatgcttcc atgtattcag 360
 ggcgcgcatt gtgagcagtg gagtatgagc cttgagggcc tcatggttgc agggcaggct 420
 tccctgcaga tgggtggcag cccctggtag aatgctggat ttctctggaa tctagaagtg 480
 ccatatttta gtggaaaggc atcagggctg tttgacagtg tgcgtctttc caatcccatg 540
 ttctccatt cgtgtgtctg ttataaaact gagtgaaggc tgctatgacc tgtgttcaact 600

121

ctggttacag ggaggtgcaa accattctgt ctcccagcct ttcttctctc tttgtgtgct 660
 cccagcactt ccttcttttc taacatggcc tggagagagt ctctctctcc ttgtctctgt 720
 ctcttaataa tagtttttaa cgtggacatc tcttccttgg tacagtgggt tttaaatcct 780
 gagaagaacc aagtcagggt ttttaaagca gactaaaagc atgaaattgc tttcagaaga 840
 atgtatatca tcgggaaaag tttgggggca gagtggggga atcaggcttt attcaaaaga 900
 aacagttgaa aacatggact tttctaccc aatgccatt tcacgactcc tctgagacta 960
 attgggaaac ggggaaattc ttggaathtt tttttaaga aacttttttg tgtttttttt 1020
 aattttaggt cacttattag tgaacctca ttttagatct gacattggta gatagatgga 1080
 tttaggcaaa tatgatgctg ttgtggggaa tccacgtggt tgacgttaga acctcccttc 1140
 tgcagactgt tgccgtcat ctaagcgaat tggaaatgct gagcttccat aagtcagctg 1200
 agttttaaag gtaaactgta tggctgaagt agtaaagcac ctgaccacaa aacctcttgt 1260
 aaaaacagcc ctgagtaggt atttcaggg ctccacaaag ttgcttatgg gaatcctgag 1320
 ctgcttttca ccatctcaag aagcctaaga agttatata ttaatcagg agacaaaaca 1380
 gttcaaagca taaggtccat ggtggtggaa aatggatgca agtgattcta agtttggtga 1440
 tttgtggata gcagagggat cgggacctct tggaggaacc ctgggtacca agctcccagg 1500
 cccttctct atcatggatg ctgggtgact ttgggaagtc accacctctt cccaagcctg 1560
 tttcccatat cacagatgtg gggccatggc ctcgatgatg gtctccacag gtctttccac 1620
 ctctgtgagt ccaagtcagg tcaatcagca aggacctatc tctgccctgg gtcagctcct 1680
 cagaaccaac cccagcatc tctaaagcaa aagcctcacc tcaagggtg ctcagaagag 1740
 agcaccttca gcatgagttg ttgctggaag atctaataag ctgtgtttcc tgggaagtgg 1800
 tgctttactt agccctgtgg acaacttctc tatgcatctg tgtgagcaga tgatcattgt 1860
 attacctttt atcggtagta agcttgaaa aataatttaa gaatacaatg gagaaatgta 1920
 aataagtatc tatgtaaatt tgtttaaat aaactgaatg tatttaattg tccatttata 1980
 tgttctttta tgtaacatgt agtttaataa agttcctggt tatgagagtc atgtttcatc 2040
 tcagcttctt ccaaaaaaaaa aaaaaaaaa agatctttaa ttaagcgg 2088

<210> 151

<211> 509

<212> DNA

<213> Homo sapien

<400> 151

cggactcccc ccgcgagcgc gctggcttcg cgtatcgggt tacttcttt ataaaaattt 60

122

ttataactta tgtggaatg ggatctcact atgttgetca gacttgtctt gaactcctgt 120
 tatagcacca cctcttacia gtgttgggca gctccgcttc tctcacttgt ctggaattct 180
 aaggcacttc cctgaagtgc tcacctgag ctaatatggg atagggctgg agagagaaca 240
 gaggtggata gcatgccaga atgaggtggg aagggtggga tcagcagcct ttgggaagga 300
 aagaagtatg agtcccaggg tattaacaag gtggaggggc aataaaattt attacatatt 360
 gggattcata ctaaagtgt agattttagc ttctcttgcc acaataaaca aaaaaaatgc 420
 cacaacacca cacaaaaaaaa aaaggtgctg ggaaccaggg ccaaactcc cggggtgaat 480
 gtttcccgcc atcaaattaa aacacacag 509

<210> 152
 <211> 560
 <212> DNA
 <213> Homo sapien

<400> 152
 ccagcctggg taacagatgt gagaccctgt ctgttaagag aatcagaaaa gagagagaaa 60
 gctagactta gctccaagtc tggagctttt ggggttttct tcctttataa aaatttttta 120
 acttatgtgg aaatgggatc tcactatggt gctcagactt gtcttgaact cctggttagc 180
 accacctctt acaagtgttg ggagctccg cttctctcac ttgtctggaa ttctaaggca 240
 cttccctgaa gtgctcatcc tgagctaata tgggataggg ctggagagag aacagaggtg 300
 gatagctgcc agaatgaggt ggaaggtgg ggatcagcag cctttgggaa ggaaagaagt 360
 atgagtccca gggattataa aggtggaggg gcaataaaat ttattacata tgggattcat 420
 actaaatgag tagatttttag cttctcttgc cacaataaac aaaaaaatg ccacaacacc 480
 acacaaaaaa aaaaggtgctg ggaaccagg gccaaactgc cccgggtgaa tgtttccgcg 540
 catcaaatta aaacacacag 560

<210> 153
 <211> 577
 <212> DNA
 <213> Homo sapien

<400> 153
 tgatgatata tggggcatgg tcctctagat gctgctcgag cggcgcagtg tgatggatgc 60
 gtggtcgcgg cgaggtacca cctgttcatt ggggaactgt gggaaacgga gccaacggac 120
 ctaagtgcc tttgacagtg agtttcatac catttcagta gtgtatttct ttcttaatct 180
 gaataacca gtatgatact ctcagacaca gaagaataaa gggagcaggt cattaacggt 240

123

ttctttttaa acctttatga tgacttcctt atgaattact gaacgaacac tggaatggga 300
 ctcaggatc ctgaggacat ctctcaactc tggccttagt tccccctctg taaaattagg 360
 gtgccaacta aatgatctac aaggtccctt ccagcgccgc cattctgtaa ttacatcatg 420
 tgtaactgta ttaaacatac acaagtgact gccaggcatg ggaatgtaac ttccgagtaa 480
 atgctttggg ttgttcagaa tacactatga acttctttcc aaagacgggt tgtggtaaat 540
 agtggatatt ttgattataa gaaatagagt ttccttg 577

<210> 154
 <211> 1138
 <212> DNA
 <213> Homo sapien

<400> 154
 aagaattcg gcacgaggaa agtgctggga ttacaagcat gagcccagcg cctggctgta 60
 tctttcattt tacccaagtc actttacca agtaagtaat taggggaaag cctgagtctt 120
 gtaccacctg ttcatttggg gaactgtggg aaacggagcc aacggacctt agtgcctttt 180
 gacagtgagt ttcataccat ttcagtagtg tatttcttcc ttaatctgaa taaaccagaa 240
 tgatactctc agcacagaag aataaagga gcgagtcatt aacgttttct ttttaacct 300
 ttatgatgac ttccttatga attactgaac gaacactgga atgggactca ggtatcctga 360
 ggacatctct caactctggc cttagtccc cctctgtaa attaggggtgc caactaatg 420
 atctacaagg tcccttccag cgccgccatt ctgtaattac atcatgtgta actgtattaa 480
 acatacacia gtgactgcca ggcatgggaa tgtaacttcc gagtaaatgc tttggtttgt 540
 tcagaataca ctatgaactt ctttccaag acgggtgtg gtaaatagtg gatattttga 600
 ttataagaaa tagagtttcc ttgaagctt agctggagat acagcaatag tgtgggttcc 660
 ctacaaatat cacagtgtat tcaaacatat ttttctatca aaaatcattt ttgtaaaagc 720
 tgtgtgtttt tatccaactt gtgataataa atgttcttta ttttagaata aaaaaaaaa 780
 aaaaaaaaa aaagaaaaa aaaggaaata aaaaaaaaa acaggagaca aagacaacgg 840
 cggcacgcaa caaccacatc gcggaaggcg acaagcgaac aaccagccc gagctcgtga 900
 aggcgagcca acatgaagga gcgcactatc caagacaggt agctgacata acagaagaga 960
 acaaaaaaaa gagacaagta gaacaaaaac aaagagaaga caggacacac gagaaaagca 1020
 ggtgtaatca gacgaacgac gcgacaaaca gagagacgtg caagcataaa atagcaaaa 1080
 ccaagagaca gcgacggaca cacgaagcaa gacgagcgac gccgagcaca gcagggat 1138

<210> 155

<211> 800
 <212> DNA
 <213> Homo sapien

<400> 155
 cgtggtcgcc ggccgagggt gccataggcc ccagaccaa ctagaccacc agcatgttca 60
 tgtccagacc tcggcagtgg cgtgcactgc ttgtgcacct cagttcctcc agtgttggtt 120
 tgtttgtttt ttaattcagc atcctgctgg ttttactttc caagcaagat ctggtgcgac 180
 tcccaaatgc gttttaatga gctcatcctt atttgccttt cttcttacgt attttgttgt 240
 atttaagat tgtgcaggag atattctaga aggcattaat ggtttgcatt caaacgatg 300
 tggtttgtcc aagttatttt ctgtctttat tactgagacg gattaatctc cttatTTTTT 360
 tcttgatgat ttgaagttgt aacagttgtc cagctattgc ttaataaaaat tttgcagatc 420
 aaaaaaaaaa aaaaaaaaaa aaaaaagggt gggggtaacc agggccaaga ggggtccctc 480
 ggggtcgaca attgggtcac ccgggtccat caatttcccc acaaacataa tacaggacat 540
 aggcacacac agcaaacgca cacagcacca agacagacaa ctacggcgag ctaaggacgc 600
 agagaagacg cggcaacgcg gaacgccccg agcaaggccg aggcaacaca ggagaggggc 660
 agcgcacgac ggccggagca cgagcaggaa agcaacgaag agagacaacg gacacacgcg 720
 agggcgaaga gaagagagca ggaacgacag gacaagcaca caaacgagcg gcaacagcag 780
 acccagacga aacagcgcga 800

<210> 156
 <211> 4632
 <212> DNA
 <213> Homo sapien

<400> 156
 atgtatgcag cagtggaaca tgggcctgtg ctttgcagtg actccaacat cctgtgcctg 60
 tcctggaagg ggcgtgtccc caagagtgag aaggagaagc ctgtgtgcag gagacgctac 120
 tatgaggaag gctggctggc cacgggcaac gggcgaggag tggttggggg gactttcacc 180
 tctagtcaact gtgcgagga caggagtact ccacagagga taaatttcaa cctccggggc 240
 cacaatagcg aggttgtgct ggtgaggtgg aatgagccct accagaaact ggccacgtgc 300
 gatgcggacg gaggcataatt cgtgtggatt cagtacgagg gcaggtggtc tgtggagctg 360
 gtcaacgacc gcggggcgca ggtgagtgat ttcacgtgga gccatgatgg aactcaagca 420
 cttatttcct atcgagatgg gtttgcctg gttgggtctg tcagtggaca aagacactgg 480
 tcatccgaaa tcaacttga aagtcaaatt acgtgtggca tatggactcc tgacgaccaa 540
 caggtgctgt ttggcacggc cgatgggcag gtgattgtca tggattgcca cggcagaatg 600

125

ctggcccaag	tcctcttgca	cgagtcagac	ggtgtcctcg	gcatgtcctg	gaactacccg	660
atcttcctgg	tggaggacag	cagcgagagc	gacacggact	cagatgacta	cgcccctccc	720
caagatggtc	cggcagcata	tcccatccca	gtgcagaaca	tcaagcctct	gctcacccgc	780
agcttcacct	cgggagacat	cagcttaatg	aacaactacg	atgacttgtc	tcccacggtc	840
atccgctcag	ggctgaaaga	ggtggtagcc	cagtggtgca	cacaggggga	cttgctggca	900
gtcgtggga	tggaacggca	gacccagctt	ggtgagcttc	ccaatgggcc	ccttctgaag	960
agtgccatgg	tcaagttcta	caatgttcgt	ggggagcaca	tcttcacact	ggacactctc	1020
gtgcagcgcc	ccatcatctc	catctgctgg	ggtcacccgg	attcgaggct	gttgatggca	1080
tcaggaccag	ccctgtacgt	ggtgcgtgtg	gagcacccgg	tgtccagcct	gcagctgctg	1140
tgccagcagg	ccatcgccag	caccttgctg	gaggacaagg	acgtcagcaa	gctgactctg	1200
cccccccgcc	tctgctccta	cctctccact	gccttcatcc	ccaccatcaa	gcccccaatt	1260
ccagatccga	acaacatgag	agactttgtc	agctacccat	cagccggcaa	cgagcggctg	1320
cactgcacca	tgaagcgcac	agaggacgac	ccggaggtgg	gcggcccgtg	ctacacgctc	1380
tacctggagt	acctgggctg	gcttgtgccc	atcctcaaag	ggcggcgcat	cagcaagctg	1440
cggccagagt	tcgtcatcat	ggacccgctg	acagatagca	aaccagatga	aatctatggg	1500
aacagcttga	tttctactgt	gatcgacagc	tgcaactgct	cagactccag	tgacattgag	1560
ctgagtgatg	actgggctgc	caagaaatct	ccaaaatct	ccagagctag	caaatcacc	1620
aaactcccaa	ggatcagcat	tgaggccctg	aagtcacca	agctgccccg	ggctgctcag	1680
gagctctccc	ggtccccacg	gttgccctg	cgcaagccct	ctgtgggctc	gccagcctg	1740
actcggagag	agtttccttt	tgaagacatc	actcagcaca	actatcttgc	tcaggtcacg	1800
tctaatatct	ggggaaccaa	atttaagatt	gtgggcttgg	ctgctttcct	gccaaccaac	1860
ctcggtgtag	taatctataa	aaccagcctc	ctgcatctcc	agccgcggca	gatgaccatt	1920
tatctcccag	aagttcggaa	aatttccatg	gactatatta	atctacctgt	cttcaacca	1980
aatgttttca	gtgaagatga	agatgattta	ccagtacag	gagcatctgg	tgtccctgag	2040
aacagccac	cttgtagcgt	gaacatccct	attgcaccga	tccacagctc	ggctcaggct	2100
atgtcccca	cgcagagcat	agggctgggt	cagtccttac	tggccaatca	gaatgtgcag	2160
ctagatgtcc	tgaccaacca	gacgacagct	gtagggacag	cagaacatgc	aggtgacagg	2220
tgccaccag	taaccagggt	ctccaaccgg	tactccaatc	ctggacaggt	gattttcgga	2280
agcgtggaaa	tgggcccgc	cattcagaac	ccccctccac	tgtccctgcc	tccccgcgcg	2340

126

caggggcca	tgcagctgtc	cacggtgggc	catggagacc	gagaccacga	acacctgcag	2400
aagtacgcca	aggccctgcy	gccaacaccg	cagctggcag	ctgaggggga	cgcagtggtc	2460
tttagtgccc	cccaggaggt	ccaggtgacg	aagataaacc	ctccaccccc	gtaccagga	2520
accatccccg	ctgccccac	cacagcagca	ccccgcccc	ctctgccgcc	cccacagccc	2580
ccagtggatg	tgtgctttaa	gaagggcgac	ttctccctct	accccacgtc	agtgcactac	2640
cagaccccc	tgggctatga	gaggatcacc	accttcgaca	gcagtggcaa	cgtggaggag	2700
gtgtgccggc	cccgccccg	gatgctgtgc	tcccagaaca	cctacaccct	ccccggcccc	2760
ggtagctctg	ccaccttgag	gctcacggcc	actgagaaga	aggctccctca	gccctgcagc	2820
agtgccaccc	tgaaccgcct	gaccgtccct	cgctactcca	tccccaccgg	ggaccacccc	2880
ccgtatcctg	aaattgccag	ccagctggcc	cagggggcgg	gggtgccca	gaggtccgac	2940
aatagcctca	tccacgctac	cctgcccagg	aacaaccgtg	aggctacgct	caagatggcc	3000
cagctggccg	acagccccg	ggccccctg	cagccccctg	ccaagtcca	gggcccggccc	3060
gggggggtg	tgacacagct	cccagcgcgg	ccccacctg	ccctgtacac	ctgcagtcat	3120
tgcagtggca	cagggcccag	ctcacagccc	ggagcctccc	tggcccatac	cgccagcgcc	3180
tccccgttgg	cctcccagtc	ctcctacagc	ctcctgagcc	cacccgacag	cgcccgcgac	3240
cgcaccgact	acgtcaactc	ggccttcacg	gaggacgagg	ccctgtcca	gcactgtcag	3300
cttgagaagc	ccttgaggca	ccctcccctg	cctgaagctg	ctgtcacctc	gaaacggcca	3360
cccccttacc	agtgggaccc	catgtgggtg	gaggacgttt	gggttctctca	agaaaggaca	3420
gcacagactt	cagggcccaa	ccccttaaaa	ctgtcctctc	tgatgctgag	tcagggccag	3480
cacctggagc	tgtcccgact	gcccttcac	tcccccaagt	ctcctgccag	ccccactgcc	3540
actttccaaa	caggctatgg	gatgggagtg	ccatatccag	gaagctataa	caacccccct	3600
ttgcctggag	tgaggctcc	ctgctctccc	aaagatgccc	tgtcccaca	gcagtttgca	3660
caacaggagc	ctgctgtggt	ccttcagccc	ctgtaccac	ccagcctctc	ctattgcacc	3720
ctgccccca	tgtaccagc	aagcagcagc	tgctctagtt	tacagctgcc	acctgtcgcc	3780
ttgcatccat	ggagttccta	cagcgcctgc	ccgcccctgc	agaacccccca	gggcactctc	3840
ccccaaagc	cacacttggt	ggtggagaag	ccccttgtgt	ccccaccacc	tgccgacctc	3900
caaagccact	tgggcacaga	ggtgatggta	gagactgcag	acaacttcca	ggaagtctct	3960
tcctgaccg	aaagcccagt	ccccagcgg	acagaaaaat	ttggaaagaa	gaaccggaag	4020
cgcctggaca	gccgagcaga	agaaggcagc	gttcaggcca	tactgaggg	caaagtgaag	4080
aaggaggcta	ggactttgag	tgactttaat	tcctaatct	ccagcccaca	cctggggaga	4140

127

gagaagaaga aagtgaagag tcagaaagac caactgaagt caaagaagtt gaataagaca 4200
 aacgagttcc aggacagctc cgagagcgag cctgagctgt tcatcagcgg ggatgagctc 4260
 atgaaccaga gccagggcag cagaaagggc tggaaaagca agcgtcccc acgggccgcc 4320
 ggcgagctgg aggaggcaa gtgccggcgg gccagtgaga aggaggacgg ggcggctgggc 4380
 agccaaggct tcgtgtacgt gatggccaac aagcagccgc tgtggaacga ggccaccag 4440
 gtctaccagc tggacttcgg ggggcgggtg acccaggagt ccgccaagaa cttccagatt 4500
 gagttagagg ggcggcaggt gatgcagttt ggacggattg atggcagtgc gtacattcta 4560
 gacttcagat atccgttctc agccgtgcag gcctttgcag ttgccctggc caacgtgact 4620
 cagcgctca aa 4632

<210> 157
 <211> 998
 <212> DNA
 <213> Homo sapien

<400> 157
 tgctgctcga gcgcgcgag tgtgatggat ccgccgggc aggtacctt tctctcaca 60
 ttggcagaat agcacgact agatgcctga ccttgagctc tagtctccc gtttaaactc 120
 taccttgggc agtaacgaca attattcctc attcaagtaa tttcaatgct gaaactgaac 180
 tctattacta atgccttcca atcagagttc ctgatgggga tgcctgtggg atggcccact 240
 aacctggggg acctaggcta gcattgggtg agttgggtaa ggaagatgat gcgtagttc 300
 ctgatagatg ctacgagatg tagtttgga tttcagttgt tgtccagtta tgattttcac 360
 tgggggttct gcagtcacag caagctgtgt atgaactagc tgtactagtg gatgacacac 420
 tataactaat caaactagac taaagacaca ctgaaaatct gcgttataac taacaagata 480
 tcactcatct gacacataac caccattaca ccttatggta cgtcaggatt cataaatagt 540
 actgctctga atgacttatg ggaaatggtg ccaactcaaaa gcaacttctt aacttgagga 600
 ataactcctt tgtagtttac tttctggtag tggttggtgc cttgtatcgg gatacagcta 660
 tattcttagc tcaaatgtct cttttggaga gcacagtagt taccctactg gtgagactga 720
 gaacctgagc tcatgagagg ccattccttc ctgggtgtcg gaccagggct ctgtgtcagg 780
 aaaaaccttc tgggtgacct ttgtagactc gtttcagggt gattccctct tatcttgcca 840
 gagtagaatt cgcagtcggt ggcccttccc ttcaccgta actcggcccc tctgggcagg 900
 ggcgggggtg cggctcttaa cgctggctcc gggtttgggg ggcgggggc ccgcaacgag 960
 ggttttgggc gggtcgccc cctccctcca acgggccg 998

<210> 158
 <211> 766
 <212> DNA
 <213> Homo sapien

<400> 158
 gggatgatcg ctcaactatag ggcgctggtc actagatgca tgccgagcgg cgccagggtga 60
 tggatcgagc ggccgcccgg gcaggtagcat gttcatgaat ttgtgctgaa taattacttg 120
 agtgtgaaat tgttatgtta tgcgatatat agtagtcaa tatagaagat aatgcaaaac 180
 aatttaaagt gattgtagca gttcgctgta ttctacagca gcaggattgt aggcagatta 240
 ctgtagttct cacagcgagc agcatgtgag attggccagt ccgctcaaat tcgtgccaat 300
 acttggata tgctatcttg tcaatttcta gacattctgg agagtgtgta gtacttgttc 360
 atcttgaca aattacactt aatagttatg tatccatttc tctaattttg ataacatttt 420
 acataagttt atcgttatga gatatgttct ttattttgaa gtgcttattg tccattttac 480
 attgggtcat ctgttattga attgtaaaca ttccttgaat atttaaataat gagtgtctgg 540
 tcagttttgg tcacaaatat cctcgttttt tcactttttg cccttttatt attctgaaaa 600
 tgccaagtga ttaaaattaa ttttactatt gttcaataaa caaaacaaaa aaaaaaaaaa 660
 aaaaacacaa aaaaacaaaa gcgcgggggg taaccggggg cccaaggggg tccccggggg 720
 acattggtct ccccggtcac aattcccccc aatcgacaaa cagggc 766

<210> 159
 <211> 1400
 <212> DNA
 <213> Homo sapien

<400> 159
 ctatgattag cttattaggc tttgtggttt atatgcatca gaaagagtaa gacttaattt 60
 tgtgtggaac aaataccctg gtgtagcatg tttcattaga atttgtttat agagatattg 120
 ccatagaaaa gttatTTTTT attagtaaag aatgctttgt atttcctttg tggcttctaa 180
 gtaccctttt ttggttatta tacctttatc cataagtatc tttaaatatt acaaaaaatta 240
 catattcttt taaatatTTT aaagatttat tatattcatt taggttttaa tccactttta 300
 attttttaga tgaaaagtaa gagaaaagta tataaatcat gagcaciaat tgaactaacc 360
 aaggtaacaa tcaatctgct caagaaatg agcatcacca ccacctcctc ctgcaactgtc 420
 caaatcagca cccagtagct ccaagcaaaa tgttactcac tacactgact tctaacacaa 480
 tagacttggt ttgtctgTTT tcaactatac aaaaatgaat catagagtat gtgttggTTT 540

129

gtatctggct cctttcacta aaattttggg ttataaaatt catccatgtg gttgaacaca 600
 gttgtagatt gttcatttta attgttttac agtattttatt gtgtgactaa aacactactt 660
 atttattcta taattgacag actttggggt gcttttgctt tgggagtata aacattttta 720
 tatctatgct ttaggtacat gttcatgaat ttgtgctgaa taattacttg agtgtgaaat 780
 tgttatgtta tgcgatata agtagtcaaa tatagaagat aatgcaaac aatttaaagt 840
 gattgtagca gtttgctgta ttctacagca gcagattgta gcagattact gtattctaca 900
 gcagcagcat gtgagattgc cagttgctca aattcgtgcc aatacttggg attttttacc 960
 ttttaatttt agacattctg gagagtgtgt agtaattttt catcttggaa aattacatta 1020
 aattagtatc catttctcta attttgataa cattttcata agtttattgt tattagatat 1080
 tttctttatt ttgaagtgtt tattgtccat ttacattgg gtcatctgtt attgaattgt 1140
 aaacattcct tgaatattta aatatgagt cttggtcagt ttttgtcaca aatatcctct 1200
 tttttcactt tttgcccttt tattattctg aaaatgcca ttgattaaaa ttaattttac 1260
 tattgttcaa taaacaaaac aaaaaaaaa aaaaaaaaaac acaaaaaaac aaaagcggg 1320
 ggggtaaccg ggggcccaag ggggtccccg ggggacattg gtctccccgg tcacaattcc 1380
 cccaatcgc acaacagggc 1400

<210> 160
 <211> 556
 <212> DNA
 <213> Homo sapien

<400> 160
 acctattcac cattccaacg tgaagaagct ctgcatgtag gaaagaataa ttaacacact 60
 tatagtctac tgcccatgta aggatcagct cgggctaaga ggccaaagat ggggtgacatc 120
 gtcatgctct gccttttatt ttttctttct taccactta gcttcctaatt tggaggaagg 180
 aggcgtggta aaggatatg aagactatgg tttaattaga ccagaaaaca ctgtcataat 240
 ctctgggctg cagtcagaat gtccagtttt gtctttgggc caagataagg gcagtgggat 300
 ttatgatgtg ttgtttatag tctgaaacta ctctggtgat caccagggtc agtttcttta 360
 atc gatggtt tccaagctgg cctaagtaca tttaagtaga gactgggctg ataaacatga 420
 ccagacgaga cataaagacc ctgttgggaa tgacattgaa ctctcaaagt caagatttct 480
 tacacaaatc tatcagctgg agaataatga gaggcagctg tggatatatgt gtgcaataa 540
 ggacattatg aagctt 556

<210> 161

<211> 1327
 <212> DNA
 <213> Homo sapien

<400> 161
 ggaagacctg attggaata gtcgaaagcc ttgatatgtg caaagaaaga accatttgat 60
 caaccagtt cttatacag gatactaact taaaatatag actcaagta tacgataatt 120
 caaacattta ttgtatttat actattctat atgtactttt ccaggaacca ggaatacaaa 180
 actgacatgt tctctgtaca gaggctcaga ctagtagaga acagttaggt acgccgttaa 240
 ttataaacta atatgtatca tcaattatgg gtttttatgg gggtttgga ggtggaaggg 300
 accagggaga gatgatgagt gatgatgggt atgtagtctt taggaggatg caattataac 360
 attgctcttc ctttcacgca ccacatgatt tagcaagtac ttcattatgg ctccaccatt 420
 aacatggtca atggcttctg gatactcaca gttcaggcac agtttctcct gaagattttt 480
 tacctctccc atctttaaga aattgtctgg atgtccatga aagatgctga cacttgtatt 540
 aattcattaa aaaacaccac cccctcctg aaataaacta aaaagtaatg aattcataga 600
 aaaaaatttc accaagattg aaactagaga atatacctag acttgcaact tgagctttga 660
 gaaatgtgta cctattcacc attccaacgt gaagaagctc tgcagtagga aaaataatta 720
 acacacttat agtctactgc ccatgtaagg atcagctccg gctaagaggc caaagatggg 780
 tgacatcggt atgctctgcc tttatTTTT ctttcttacc cacttagctt cctaattgga 840
 ggaaggaggc gtggtaaagg tatatgaaga ctatggttta attagaccag aaaacactgt 900
 cataatctct ggggtcatca gaatgtccag ttttgtcttt gggccaagat aagggcagtg 960
 ggatttatga tgtgttgttt atagtctgaa actactctgg tgatcaccag ggtcagtttc 1020
 tttaatgatg gtttccaact ggcctaatac attaagtaag actggctgat aacatgacca 1080
 gacagacata aagaccctgt tgggaatgac attgaaactct caaagtcaag atttcttaca 1140
 caaatctatc agctggagaa aatgaaggca gtgtgtgata tgtgtgcaaa taaggacatt 1200
 atgaagctta aatatggaat gtctcttggg cccccgatgt catctgtatt ctctttttct 1260
 tcttgtaacta tacccttgc ctgtaaataa aaggtttatt tgaaaaaaaa aaaaaaaaaa 1320
 gatcggc 1327

<210> 162
 <211> 318
 <212> DNA
 <213> Homo sapien

<400> 162
 ggttctccta aatgtcttaa cccatgttta tcttgttctg ctattccatg agcaaagaga 60

131

ataaagcaca aagctgtgag agtattaat atggacacta gatttacatt tccaacaaga 120
 aattcatctc cctccaaagt cccagaccag ggctagaatg tggttcattt ttaacaatca 180
 aagtggcaag atctgtttgg tgatcactgt aaaacaggaa acacagtaat gccttcatgt 240
 tgagggtgcta aaaggtcaag cttgggtaac aatgtccata gctggtctgg tgaatgtttc 300
 gtcaatcaaa tagtgaaa 318

<210> 163
 <211> 1042
 <212> DNA
 <213> Homo sapien

<400> 163
 acagtctgtt tcctccttca cccccagaac aaaaatcgaa cttctggttg gacagtgtca 60
 gatgtcactg aggtgacccc agcctgtttg cagttccaag tcttccgtgt aggcgtcact 120
 gctactggaa cttttagat gaggagcctg tatgatgatg tcctgaacat ttctatcctt 180
 tcctcacaca gagggaagct actgggaata tcagagacaa gctattatta aacaagtgtc 240
 tctagtccaa gacatctcct gtggcagggg aatgaggggg caggctgtat cagtgatatt 300
 ttataaact ctggttttag aaaaaattct tcagatggac gcattathtt aagactttaa 360
 cattttccaa aaccaactga atcttatccc ctccatttat cccctccag acaacttctaa 420
 tcaaggtcac catctccaac tcccccata gacagtaaaa atatggctgg agaattctac 480
 tgtaatagaa aaccaaggag atatggtaat ttgacagtgt gtttcctttc catccactag 540
 acaagaatac cccctccat tctttcctcc cctcagtcac cagaatgaag tgggctggaa 600
 aacagttggt ctggttctt tatagagact gattcccaca ttggatactg cctggaggcc 660
 ttgggatga atgagaagtt ctgctggttt ggatcagtag cagaagcagg taacacatca 720
 gggaaaccgt cagcctaaga taggagggga cagaaaatga tgaaagagtt tctgatacat 780
 ttatcagcta aattgctatg gtcacccca tgtctcctgt aatgtccaac actaaggaat 840
 taaactaagt aactacaac ctttgtgtct tgctctgacc ttggaccaat ggaatatact 900
 tcttatttca tattcagtgg ataagcaaat ctgcttcac cctgccttaa ctactcaag 960
 gtctctgtga tgcaactcag agttttcctc cttccctgca tagtcttctc ctccctagct 1020
 gcctttcaaa ttggtgaaaa tg 1042

<210> 164
 <211> 1120
 <212> DNA
 <213> Homo sapien

132

<400> 164
 gccgcctttt tttttttttt ttttttagaca agaaattatt ttagtccttt agtacagtct 60
 gtttcctcct tcacccccag aacaaaaatc gaacttctgg ttggacagcg tcagatgtca 120
 ctgaggtgac ccagcctgt ttgcagttcc aagtcttccg tgtaggcgctc actgctactg 180
 gaactttgta gatgaggagc ctgtatgatg atgtcctgaa catttctatc ctttcctcac 240
 acagagggaa gctactggga atatcagaga caagctatta ttaaacaagt gtctctagtc 300
 caagacatct cctgtggcag ggaaatgagg gggcaggctg tatcagtgat atttttataa 360
 actctggttt tagaaaaaat tcttcagatg gacgcattat ttaagactt taacattttc 420
 caaaaccaac tgaatcttat ccctccatt tatccccctc cagacacttc taatcaaggt 480
 caccatctcc aacttcccc atagacaata aaaatatggc tggagaattc tactgtaata 540
 gaaaaccaag gagatatagt aatttgacag tgtgtttcct ttccatccac tagacaagaa 600
 taccctctcc cattctttcc tcccctcagt caccagaatg aaggggctgg aaaacggtgg 660
 tctggttcct tttagagctg attccccatt ggatactgcc tggaggcctt ggggatgaat 720
 gagaagttct gcagtttgga tcagtagcag aagcaggtaa cacatcaggg aaccggctcag 780
 cctaagatag gaggggacag aaaatgatga aagagtttct gatacattta tcagctaaat 840
 tgctatggtc acccccatgt ctctgtaat gtccaaacct aaggaattaa ctaagtaaac 900
 taaaaccttt gtgttcttgc tctgacctg gacaatggaa ttcttcttat tttcattcag 960
 tggatagcaa atctgcttct tccctgcctt aactcactca aggtctctgt gatgcactcc 1020
 agagttttcc tccttcctg catagcttcc tcctccctag ctgcctttca aattggtgaa 1080
 aatgaagctt caggattatg aaaactagta cttaatgaag 1120

<210> 165
 <211> 810
 <212> DNA
 <213> Homo sapien

<400> 165
 agatcatgct cgagcggcgc agtgtgatgg attggctcgc gccgaggtag ttttttatgg 60
 cttacatctg tgcctggctg gccatcaagt ctgggtgcca ctgtttgaga tttggggctg 120
 tttctgcaa ctgatctctg ctacagataa ggcttccctc ctggaggcca aagccctggt 180
 taacgttaag agctctatga tgatgcaaac ttcagaggcg atcacctaac ataacaaaaa 240
 cctccccaga accagaacct gttttttcac caaaacctt ccgctgcttg aataagaatg 300
 tcttttcctt tcctaccaac tttgatgcca ctggccactg tgacataact tttacttagc 360

133

ggggtaaadc atagatggat tacttgaact gccaacacaa gactgctgga cgagggacag 420
 agctggatat gttagacaaa gatatacгаа cgacttggcg taatcactgg tcaatagctg 480
 acaccatgat gtgaaaagta gtaatcacgg ctcacaagta ccaacacaag atacagaaga 540
 caggagaaga ggaacaggaa aagaagaaac aacagagcac aaagagagaa caagcacaca 600
 acagacgaag gccacaagag cgaaggagga ccggacgcag caccagcaac agaggaacgg 660
 cacgcacaga agaacacaga caagaaaacg agaagaaacc acacgcacaa ctagccagaa 720
 tcagagacag aaaaacgcaa gacaggaggc agaagcagaa acacaagaaa accgaacacc 780
 aaaacaggca gcacaaacac gaagagaaag 810

<210> 166
 <211> 601
 <212> DNA
 <213> Homo sapien

<400> 166
 gaagtataac tatatgggcg aatgggtcct tagatgcagg ctcgagcggc gcagtgtgat 60
 ggatccgccc gggcaggtag tcagggtgta tatgatattc tgagctgaat aagtgcgagg 120
 agcagattat taagatctgc cattctgaaa cgctggctct tttctccttc ctatagtgca 180
 ccataaaatt ctgttgatca gattatatta catacatttg ggggagtgga gggacatgag 240
 ttaagtagcc cttcatgtat ttataatctc ttttctactg aatcaaatga cttagccatg 300
 accctgaatg gacctgtttt acttcaagtг agatgtctgc cttttatgaa ttgtatatgt 360
 gaatagagtt cgggggttgc caaaaatgca tacatgtatg taagtaaaat tttttatgaa 420
 gtagtctgtc aaatgtatca taaagtttat ttttctttta tacgtaaatc attaaaaata 480
 atcacatatt tttgaaaaaa aaaaaaaaaa aaaaaaaggг ggggggtatc tcggggccaa 540
 aaggggtccc gggggggaat tggttttccg gttcaaattt ccacaaattt gggagaaaat 600
 a 601

<210> 167
 <211> 1035
 <212> DNA
 <213> Homo sapien

<400> 167
 tggtcgcggc gaggtactgt aaatgtgatg gaaaacattg atgagaattt attggcagtt 60
 cagattgtgt tttcccaact taggctcttt attaattggt taaggttttc tccaaaaagg 120
 gcatttcaac aatgggaatt attttaaat ggttaaacca gtgggcacag attacktacc 180
 ttccttctct gctttgtgac tcaccagcag taacacacac aatccacatc ttgtgcacct 240

caaatgaaca gacttggttt ccttgctttc ttgacatttc catgactgtt tcacatacaa 300
 actattgggt gaggtttttc agctgttacc gaccacgtc ctgctgtctc tgtgtgggcc 360
 tacaaaaact gtccattccc acccctttgc tttgccattt gcaagagtct ggaattgtca 420
 ggtctcagct tcgaaaagtc ctggttccac tgacaggaca cattcttttag tgggaattaa 480
 gacctacaaa gtctagtttg tatgtaggta tgaagggat tttttaaata aagtggaaaa 540
 gctgtgaaca gcattagaac tctgtctatt tcttaatttt aaaatatgct gatatgcctt 600
 aaactgtagt tgtagatcct tgtcattttg ctgtttgaaa ataaccaatg tgttttctaa 660
 aactgtcgtg taatctactt tcattgttaa tgcagaattg tcatatatgt aagccgcatg 720
 ttgacatttt gtctttttta aactaaagta attgtattga tgtgaagcat atcatttttt 780
 caaatatgaa agtgatcact tagcaacatg cttggtaatt tggcatctgt taaggtagga 840
 gagtggtgaa cagataatct atgcataat cactagtgcc aagacataaa gcgggggaaa 900
 atatattttt acccaaacat taaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaggc 960
 tgggggtaac cggggccaaa ggggtcccgg ggtgaattgg ttttccgctc aaattcccc 1020
 atttttgggc aaacc 1035

<210> 168
 <211> 1666
 <212> DNA
 <213> Homo sapien

<400> 168
 ctgggtgatg aagtgagact ctccaaaaa aaaaagaaat tattaatccc tgcctgtgct 60
 ctacatagcc tcatgggcat cattggatag ctgagagggc ccttgattct ggcaaggcaa 120
 ataaagccag aatgagaaat taccatcttc tactagagaa aaccaagaga aaaattttta 180
 tgctaggatg cctttatgac caottaattt tttaatctta gtttaatggg ctctccctgg 240
 tgctaactgc tgacagtggc cacctctttt ttggggattg aggggcctac ataactagct 300
 ggccttacc catatctttt gttcaaacat aataccatct ttttgcttct tctgaacttt 360
 agatctccat aacacatgta ctgtagaatg tgatggaaaa gcattgatga gaatttattg 420
 gcagttcaga ttgtgttttc ccaacttagg ctctttatta attggttaag gttttctcca 480
 aaaaggcat ttcaacaatg ggaattattt aatgtaacag tgggcacaga ttacttatct 540
 tccttctctg ctttgtgact caccagcagt aacacacaca atccacatct tgtgcacctc 600
 aatgaacag acttggtttc cttgctttct tgacatttcc atgactgttt cacatacaaa 660
 ctattgggtg aggtttttca gctgttaccg acccaagtc tgcctgtctct gtgtggctct 720

135

acaaaaactg tccattccca cccctttgct ttgccatttg caagagtctg gaattgtcag 780
 gtctcagctt cgaaaagtcc tggttccact gacaggacac attccttagt gggaaattaag 840
 acctacaaag tctagtttgt atgtaggat gaaggaatt ttttaaataa attgaaaagc 900
 tgtgaacagc attagaactt tgtctatttc ttaattttta aatatgctga tatgccttaa 960
 actgtagttg tagatccttg tcattttgct gtttgaaaat aaccaatgtg ttttctaaaa 1020
 ctgtcgtgta atctactttc attgttaatg cagaattgtc atatatgtaa gctgcatggt 1080
 agacatttgt cttttttaa ctaaagtaat tgtattgatg tgaagcatat cattttttca 1140
 aatatgaaag tgatcactta gcaacatgct tggaatttg gcatctgtta aggtaggaga 1200
 gtggtgaaca gataatctat gcatatatca ctagtgccaa gacataaagc gggggaaaat 1260
 atatttttac ccaaacatta aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa caactgtggt 1320
 cggcgcgctt gtggccccgg aagaagagtc ttctcgtaga accatcgtgg tttgggcca 1380
 gcggggcccc aggagtagg gtgccacacg ggccaaaagc gtgtcccagg agacacccgg 1440
 gggcactaga acaacttagg gtgtgtgagg aatattttcg ctcaccccat gttacaaaaa 1500
 caaccgcgca gagggggcaa acagcaacag ggtttctgtg aaacaacaac ccccaaatgg 1560
 agggaagtcc tcgagaagga catacaggga aagcctaata caacagaggg aagatcccaa 1620
 ggaaaagcac tatcatataa ataattatcg ccgccggctg tgccggg 1666

<210> 169
 <211> 633
 <212> DNA
 <213> Homo sapien

<400> 169
 aaaacaacac ggaatgtcta cgactaacta tagggcccct ggtgtatcta gatgcatgct 60
 cgagccggcc gccatgatgt gactggatgt cgcggccgag gtacagagta tgtagtgggc 120
 atctgttgaa tgaatgcttt tcccagtacg cacgtgtatt catacaatat taatataatt 180
 agtcccctgg gcttacagat aaaaatgaaa cgcatcaacg tgcccagctg cagtgagacc 240
 caggtgttct tcctccacc ctagtgggcc cctgggcagg tctttttttt ttggtaacac 300
 tcaccaggtc tgttctgtag tcaatcatgt gatggactgt gtccgtgaaac tgtgcaggac 360
 actgttctca tagtgttcat tagcgacaga gtaaacatgt ttgccatgca agggttatatt 420
 ggcattctgca ttttaagtgt aatgttgaat caatgaaaag gtgttgatta agcagtagtt 480
 gtagatatgc taagtttttc aaattactaa tatcaagtgg agatggtttt tactttataa 540
 gggatttgct ttggatgtag cataaataat gggtttcctt ttttgtaac tgtaacatta 600

136

attggctggc aactttggta ttcccataga ctg 633

<210> 170
 <211> 563
 <212> DNA
 <213> Homo sapien

<400> 170
 gggaaggaag acatataggg cggaatgggt octagatgca tgtcgagcgg cgcagtgtga 60
 tggatcggcg ccgggcaggt acaaaaaata ggataaatgc ttgttttttt atttagcaat 120
 gtccaaaata atgaattgat ttcccagata tcctctaaag gtaaccaggg atttttttta 180
 ttttaattatc ttgaaccac atatttaaat atacgtagta tgctacaaac cattgcagtt 240
 aagtaccttt attgatgctt gagtggcca cttttctttt tttttttttt ggagacagag 300
 cctcgctctg tcaccaggc tggagtgcag gggcgtcatc tttgactcac ttgcaacctt 360
 ccttccttcc gtggggtgca ggcagattct cctgtgcctt acagcctccg agtttgctg 420
 ggatttacag ggcattgttg caagtttccc acattttcag tgagaaattc ctcaattggc 480
 ctccgtgagt ggtttgaaa ttgaccagc aattcttggg gtgggtgtat tagctatcta 540
 tggctggtgt aacaaattga cct 563

<210> 171
 <211> 682
 <212> DNA
 <213> Homo sapien

<400> 171
 gaaaaggttg gcagcaggtg cacgtgttat cagcctgatc atctatcacc tgatggtttt 60
 agcaatacct aatccgtga tatcatcaga ggttgcaaaa tgatgagatt caggtttttt 120
 ttttacataa ttatttgtca gaattattct gcaaatagct tctctttaac agtattcggg 180
 taccttgaaa tacaggttgt acaaaaaata ggataaatgc ttgttttttt atttagcaat 240
 gtccaaaata atgaattgat ttcccagat cctctaaagg taaccaggga ttttttttat 300
 ttaattatct tgaaccaca tatttaata tacgtagtat gctacaaacc attgcagtta 360
 atacctttat tgatgcttga gttgccact tttttctttt tttttttttg gagacagagc 420
 ctcgctctgt caccaggct ggagtgcagg ggcgtcatct ttgactcact tgcaaccttc 480
 cttccttccg tggggtgcag gcagattctc ctgtgcctta cagcctccga gtttgctgg 540
 gatttacagg gcattgttgc aagtttccca cattttcagt gagaaattcc tcaattggcc 600
 tccgtgagtg gtttggaat tgaccocaga attcttggag tgggtgtatt agctatctat 660

137

ggctggtgta acaaattgac ct

682

<210> 172
 <211> 75
 <212> PRT
 <213> Homo sapien

<400> 172

Met Gly Pro Arg Ser Arg Leu Trp Pro Ser Ser Pro Leu Trp Leu Val
 1 5 10 15

Gln Pro Leu Cys Thr Pro Gly Val Phe Thr Pro Gly Ala Asp Ser Ser
 20 25 30

His Cys Ser Ser Phe Leu Arg Glu Ile Thr Val Val Ile Ala Ala Gly
 35 40 45

Ala Asn Arg Leu Gly Leu Val Ser Cys Ala Phe Gly Gln Leu Leu Thr
 50 55 60

Arg Ser Ser Leu Lys Gln Trp Gly Gly Pro His
 65 70 75

<210> 173
 <211> 38
 <212> PRT
 <213> Homo sapien

<400> 173

Met Phe Pro Arg Leu Asp Ser Thr Ser Trp Pro Gln Gly Ile Leu Trp
 1 5 10 15

Ala Trp Thr Pro Lys Pro Leu Arg Leu Glu Val Cys Glu Pro Pro Ser
 20 25 30

Leu Pro Ser Leu Trp Ser
 35

<210> 174
 <211> 52
 <212> PRT
 <213> Homo sapien

<400> 174

Met Thr Leu Phe Ile Arg Cys Cys Thr Asn Tyr Gly Asn Leu Cys Gln
 1 5 10 15

138

Tyr Phe Asn Val Cys Trp Ile Ile Thr Asp Ile Phe Ile Ile Leu Met
 20 25 30

Ser Thr Asn Leu Phe Ile Leu Ile Ala Arg Val Ser Leu Gly Ser Lys
 35 40 45

His His Leu Gly
 50

<210> 175
 <211> 37
 <212> PRT
 <213> Homo sapien

<400> 175

Met Ala Gly Ser Gly Lys Val Pro Ile Thr Thr Thr Tyr Lys Pro Pro
 1 5 10 15

Thr Asn Ser Asn Ala Ile His Leu Pro Thr Pro Ile Ile Arg Lys Ala
 20 25 30

Gly Phe Thr Gly Ile
 35

<210> 176
 <211> 88
 <212> PRT
 <213> Homo sapien

<400> 176

Met Gly Leu Thr Leu Lys Ser Leu Cys Asp Ser Lys Met Asn Cys Gln
 1 5 10 15

Ser Asn Val Pro Leu Met Lys Asp Pro Ile Thr Leu Gln His Val Cys
 20 25 30

Ile Gln Arg Thr Tyr Leu Arg Leu Ser Phe Gly His Gly Gly Arg Leu
 35 40 45

Leu Leu Lys Thr Tyr Gln Ser Pro Leu Trp Arg Ser Ala Asp Arg Pro
 50 55 60

His Asp Leu Gly Asn Gly Leu Leu Val Ile Trp Asp Cys Leu Gly Leu
 65 70 75 80

139

Cys Asn Gly Thr Trp Gly Gln Asn
85

<210> 177
<211> 61
<212> PRT
<213> Homo sapien

<400> 177

Met Asp His Lys Ser Ala Asn His Ser Ser Ala Leu Leu Lys Met Leu
1 5 10 15

Leu Ala Gly Gly Met Ser Leu Pro Glu Val Pro Glu Gly Leu Thr Pro
20 25 30

Thr Pro Ser Ser Gln Thr His Leu Ser Lys Gly Lys Gly Arg Asn Leu
35 40 45

Glu Lys Ser Tyr Phe His Asn His Ser Leu Arg Glu Pro
50 55 60

<210> 178
<211> 198
<212> PRT
<213> Homo sapien

<400> 178

Met Thr Pro Ile His Leu Ile Cys Ser Pro Ser His Glu Leu Gln Asp
1 5 10 15

Thr Thr His Pro Gln Pro Gln Arg Glu Cys Gln Arg Phe Ser Thr His
20 25 30

Gly Ala Gln Thr Thr Gln Cys Ala Thr His His His Pro Tyr Ile Ser
35 40 45

Gly Ala Ala Thr Arg Thr Tyr Leu Arg His Val Ala Pro Asp Tyr Ser
50 55 60

Ala Pro Leu Met Ala Pro Pro Thr Asn Thr Arg Leu Ala Pro Ala Ser
65 70 75 80

Leu Gln Pro Thr His Leu Arg Pro Pro Leu Ala Arg His Pro Leu Thr
85 90 95

Ala Asp Cys Arg Thr His Gln Leu Thr Asp Thr Arg Pro Leu His Pro

140

100

105

110

Arg Pro Ile Thr Ser Arg Thr Pro Gln Pro Leu Pro Ser His Thr His
 115 120 125

Gly Leu His His Thr Arg Pro Pro His Thr Ala Thr Gly Cys Pro Tyr
 130 135 140

Leu Ser Thr Ser Arg Pro Leu Pro Pro Leu His Thr Arg Ser Ile His
 145 150 155 160

Pro Asp Asn Pro His Cys Thr Thr Pro His His Ser Pro Ser Lys Pro
 165 170 175

Ser Thr Thr Thr His Gln Gln Ser Pro Ala Pro Thr Pro Asn Lys Pro
 180 185 190

His Pro Arg Arg Ala Ser
 195

<210> 179
 <211> 20
 <212> PRT
 <213> Homo sapien

<400> 179

Met Ile Gly Ile Thr Trp Cys Phe Glu Leu Ile His Pro Thr Leu Glu
 1 5 10 15

Leu Thr Ala Thr
 20

<210> 180
 <211> 107
 <212> PRT
 <213> Homo sapien

<400> 180

Met Gly Ala Ser Gly Pro Glu Arg Glu Asp Arg Asn Ser Glu Asn Gly
 1 5 10 15

Val Glu Lys Lys Asn Val Lys Glu Leu His Glu Glu His Met Ala Glu
 20 25 30

Lys Lys Glu Leu Gln Glu Glu Asn Gln Arg Leu Gln Gly Leu Pro Val
 35 40 45

141

Ser Gly Ser Glu Glu Gly Arg Leu Pro Val Pro Ser Ala Arg Ser Ser
 50 55 60

Thr Leu Arg Ala Ser Cys Arg Asn Glu Leu Gly Ser Leu Leu Pro Gly
 65 70 75 80

Gly Glu Thr Ser Leu Gly Leu Lys Glu Gly His Arg Thr Lys Gly Ala
 85 90 95

Arg Gly Gly His Arg Glu Asp Pro Gln Glu Lys
 100 105

<210> 181
 <211> 27
 <212> PRT
 <213> Homo sapien

<400> 181

Met Ser Thr His Ser Val His Ser Thr Gly Leu Pro Phe Tyr Lys Leu
 1 5 10 15

Ser Leu Thr Ser Leu Ser Ser Met Thr Leu Val
 20 25

<210> 182
 <211> 40
 <212> PRT
 <213> Homo sapien

<400> 182

Cys Phe Glu Lys Met Leu Asn Arg Leu Gly Ala Val Ala His Val Cys
 1 5 10 15

Asn Pro Ser Thr Leu Gly Gly Arg Gly Gly Trp Ile Met Arg Ser Gly
 20 25 30

Val Arg Asp Gln Pro Gly Gln His
 35 40

<210> 183
 <211> 26
 <212> PRT
 <213> Homo sapien

<400> 183

142

Met Arg Lys Gln Ala Phe Asp Leu Leu Glu Ser Thr Ala Gln Lys Ser
 1 5 10 15

Leu Val Pro Ile Phe Glu Phe Pro Lys Gln
 20 25

<210> 184
 <211> 39
 <212> PRT
 <213> Homo sapien

<400> 184

Met Lys Glu Glu Gly Arg Leu Leu Thr Val Ala Glu Gly Arg Gln Gly
 1 5 10 15

Pro Ser Cys Ser Ser His Ile Asn Ser Lys Lys Pro Ser Gln Gln Asn
 20 25 30

Lys Ser Ile Phe Asn Ser Ser
 35

<210> 185
 <211> 76
 <212> PRT
 <213> Homo sapien

<400> 185

Met Val Glu Pro Ala Leu Ser Gly Cys Gln Gln Arg Lys Gly Gly Tyr
 1 5 10 15

Ser Ser Glu Arg Gln Ser Gln Pro Thr Gln Gly Gly Gln Gly Val Arg
 20 25 30

Pro Gln Thr Tyr Ser Pro Ala Asp Leu Thr Val Arg Pro Ser Cys Ser
 35 40 45

Gly Thr Gly Asn Ala Gln Ala Glu Ile Ala Leu Leu His Thr Tyr Asn
 50 55 60

Thr Thr Leu Glu Asn Asn Leu Glu Trp Phe Thr Leu
 65 70 75

<210> 186
 <211> 35
 <212> PRT
 <213> Homo sapien

143

<400> 186

Met Arg Gln Pro Cys Leu Ala Ile Pro Glu Ala Ser Ala Ser Leu Ile
 1 5 10 15

Cys Arg Cys His Arg His Phe Thr Tyr Ser His Leu Met Ala Arg Phe
 20 25 30

Leu Leu Leu
 35

<210> 187

<211> 76

<212> PRT

<213> Homo sapien

<400> 187

Met Phe Phe Ala Leu Met Gly Ile Cys Pro Gly Thr Leu Pro Pro Gly
 1 5 10 15

Pro Pro Leu Pro Arg Trp Pro Pro Pro Val Phe Cys Phe Phe Phe Phe
 20 25 30

Phe Phe Gly Phe Phe Phe Cys Cys Phe Thr Val Lys Leu Phe Ile Glu
 35 40 45

Gln Ile Glu Asp Asn Asp Ile Cys Phe Tyr Tyr Arg Ser Leu Pro Ser
 50 55 60

Ser Tyr Ile Ile Asp Thr Tyr Tyr Glu Thr Cys Ile
 65 70 75

<210> 188

<211> 173

<212> PRT

<213> Homo sapien

<400> 188

Met Ile Gly Cys Ser Leu Leu Val Ala Cys Leu Cys Cys Leu Val Gln
 1 5 10 15

Ser Phe Arg Ala Met Phe Ser Cys Phe Ser Gly Leu Ser Leu Cys Leu
 20 25 30

Met Leu Pro Leu Trp Cys Val Cys Pro Thr Val Cys Ala Phe Phe Cys
 35 40 45

144

Gly Tyr Leu Leu Phe Phe Ser Leu Arg His Ala Ala Cys Gly Cys Leu
 50 55 60

Leu Val Cys Leu Ser Cys Leu Ala Leu Pro Ser Gly Pro Ile Leu Ser
 65 70 75 80

Phe Ser Phe Cys Leu Arg Val Val Ser Ser Val Arg Val Ala Cys Ala
 85 90 95

Arg Ser Ala Ala Val Leu Leu Leu Arg Gly Val Pro Pro Pro Ser Leu
 100 105 110

Arg Thr Leu Ser Leu Ile Ala Ser Thr Ala Thr Arg Leu Ser Phe Val
 115 120 125

Phe Leu Phe Ser Leu Pro Arg Gly Leu Leu Cys Val Gly Gly Ser Gly
 130 135 140

Ser Val Leu Gly Ser Leu Val Arg Arg Ala Gln Ser Val Gly Leu Arg
 145 150 155 160

Asp Phe Val Ser Val Leu Gln Val Val Leu Thr Cys Leu
 165 170

<210> 189
 <211> 29
 <212> PRT
 <213> Homo sapien

<400> 189

Met Val Leu Tyr Ser Glu Gly His Gln His Gly Pro His Leu Leu Asn
 1 5 10 15

Met Glu Asn Gln Asn Leu Asn Glu Leu Pro Leu Lys Gly
 20 25

<210> 190
 <211> 122
 <212> PRT
 <213> Homo sapien

<400> 190

Phe Phe Ala Asp Glu Val Ser Arg Leu Ser Pro Gly Leu Glu Cys Ser
 1 5 10 15

145

Gly Val Ile Ser Ala His Cys Asn Phe His Leu Leu Gly Ser Ser Ser
 20 25 30

Ser Pro Ala Ser Ala Ser Gln Val Ala Glu Ile Thr Gly Ala Cys His
 35 40 45

Pro Thr Trp Leu Ile Phe Val Ile Leu Val Glu Thr Gly Phe His His
 50 55 60

Val Gly Gln Ala Asp Ala Leu Leu Thr Ser Gly Asp Pro Pro Phe Ser
 65 70 75 80

Ala Pro Lys Val Leu Gly Ile Thr Gly Val Ser His Arg Ala Arg Pro
 85 90 95

Ala Asn Thr Phe Ala Leu Thr Thr Leu Gly Leu Leu Tyr Lys Ile Val
 100 105 110

Met Ile Ala Met Glu Val Leu Pro Val Pro
 115 120

<210> 191
 <211> 11
 <212> PRT
 <213> Homo sapien

<400> 191

Met Trp Arg Ala Lys Gln Tyr Asp Leu Gln Thr
 1 5 10

<210> 192
 <211> 28
 <212> PRT
 <213> Homo sapien

<400> 192

Met Met Phe Ser Leu Ser Gln Lys Gly Ser Ala Ala Val Gln Ser Pro
 1 5 10 15

Ser Thr Leu Ser Thr Pro Thr Phe Ser Ile Ser Tyr
 20 25

<210> 193
 <211> 48
 <212> PRT
 <213> Homo sapien

146

<400> 193

Met Asp Ser Gly Ala Arg Ala Gly Lys Pro Leu Leu Asp Pro Val Cys
 1 5 10 15

Leu Pro Ala Trp Ser Leu Cys Leu Gln Pro Cys Leu Tyr Ser Ser Leu
 20 25 30

Pro Pro His Gln Pro Pro Leu Ala Ser Pro Tyr Arg Leu Ser Lys Lys
 35 40 45

<210> 194

<211> 1138

<212> PRT

<213> Homo sapien

<400> 194

Met Gly Asp Phe Ala Ala Pro Ala Ala Ala Asn Gly Ser Ser Ile
 1 5 10 15

Cys Ile Asn Ser Ser Leu Asn Ser Ser Leu Gly Gly Ala Gly Ile Gly
 20 25 30

Val Asn Asn Thr Pro Asn Ser Thr Pro Ala Ala Pro Ser Ser Asn His
 35 40 45

Pro Ala Ala Gly Gly Cys Gly Gly Ser Gly Gly Pro Gly Gly Gly Ser
 50 55 60

Ala Ala Val Pro Lys His Ser Thr Val Val Glu Arg Leu Arg Gln Arg
 65 70 75 80

Ile Glu Gly Cys Arg Arg His His Val Asn Cys Glu Asn Arg Tyr Gln
 85 90 95

Gln Ala Gln Val Glu Gln Leu Glu Leu Glu Arg Arg Asp Thr Val Ser
 100 105 110

Leu Tyr Gln Arg Thr Leu Glu Gln Arg Ala Lys Lys Ser Gly Ala Gly
 115 120 125

Thr Gly Lys Gln Gln His Pro Ser Lys Pro Gln Gln Asp Ala Glu Ala
 130 135 140

Ala Ser Ala Glu Gln Arg Asn His Thr Leu Ile Met Leu Gln Glu Thr
 145 150 155 160

147

Val Lys Arg Lys Leu Glu Gly Ala Arg Ser Pro Leu Asn Gly Asp Gln
 165 170 175

Gln Asn Gly Ala Cys Asp Gly Asn Phe Ser Pro Thr Ser Lys Arg Ile
 180 185 190

Arg Lys Asp Ile Ser Ala Gly Met Glu Ala Ile Asn Asn Leu Pro Ser
 195 200 205

Asn Met Pro Leu Pro Ser Ala Ser Pro Leu His Gln Leu Asp Leu Lys
 210 215 220

Pro Ser Leu Pro Leu Gln Asn Ser Gly Thr His Thr Pro Gly Leu Leu
 225 230 235 240

Glu Asp Leu Ser Lys Asn Gly Arg Leu Pro Glu Ile Lys Leu Pro Val
 245 250 255

Asn Gly Cys Ser Asp Leu Glu Asp Ser Phe Thr Ile Leu Gln Ser Lys
 260 265 270

Asp Leu Lys Gln Glu Pro Leu Asp Asp Pro Thr Cys Ile Asp Thr Ser
 275 280 285

Glu Thr Ser Leu Ser Asn Gln Asn Lys Leu Phe Ser Asp Ile Asn Leu
 290 295 300

Asn Asp Gln Glu Trp Gln Glu Leu Ile Asp Glu Leu Ala Asn Thr Val
 305 310 315 320

Pro Glu Asp Asp Ile Gln Asp Leu Phe Asn Glu Asp Phe Glu Glu Lys
 325 330 335

Lys Glu Pro Glu Phe Ser Gln Pro Ala Thr Glu Thr Pro Leu Ser Gln
 340 345 350

Glu Ser Ala Ser Val Lys Ser Asp Pro Ser His Ser Pro Phe Ala His
 355 360 365

Val Ser Met Gly Ser Pro Gln Ala Arg Pro Ser Ser Ser Gly Pro Pro
 370 375 380

Phe Ser Thr Val Ser Thr Ala Thr Ser Leu Pro Ser Val Ala Ser Thr

149

Lys Asn Pro Leu Met Pro Tyr Ile Gln Gln Gln Gln Gln Gln Gln
625 630 635 640

Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Pro Pro Pro Pro Gln Leu
645 650 655

Gln Ala Pro Arg Ala His Leu Ser Glu Asp Gln Lys Arg Leu Leu Leu
660 665 670

Met Lys Gln Lys Gly Val Met Asn Gln Pro Met Ala Tyr Ala Ala Leu
675 680 685

Pro Ser His Gly Gln Glu Gln His Pro Val Gly Leu Pro Arg Thr Thr
690 695 700

Gly Pro Met Gln Ser Ser Val Pro Pro Gly Ser Gly Gly Met Val Ser
705 710 715 720

Gly Ala Ser Pro Ala Gly Pro Gly Phe Leu Gly Ser Gln Pro Gln Ala
725 730 735

Ala Ile Met Lys Gln Met Leu Ile Asp Gln Arg Ala Gln Leu Ile Glu
740 745 750

Gln Gln Lys Gln Gln Phe Leu Arg Glu Gln Arg Gln Gln Gln Gln
755 760 765

Gln Gln Gln Gln Ile Leu Ala Glu Gln Gln Leu Gln Gln Ser His Leu
770 775 780

Pro Arg Gln His Leu Gln Pro Gln Arg Asn Pro Tyr Pro Val Gln Gln
785 790 795 800

Val Asn Gln Phe Gln Gly Ser Pro Gln Asp Ile Ala Ala Val Arg Ser
805 810 815

Gln Ala Ala Leu Gln Ser Met Arg Thr Ser Arg Leu Met Ala Gln Asn
820 825 830

Ala Gly Met Met Gly Ile Gly Pro Ser Gln Asn Pro Gly Thr Met Ala
835 840 845

Thr Ala Ala Ala Gln Ser Glu Met Gly Leu Ala Pro Tyr Ser Thr Thr
850 855 860

150

Pro Thr Ser Gln Pro Gly Met Tyr Asn Met Ser Thr Gly Met Thr Gln
865 870 875 880

Met Leu Gln His Pro Asn Gln Ser Gly Met Ser Ile Thr His Asn Gln
885 890 895

Ala Gln Gly Pro Arg Gln Pro Ala Ser Gly Gln Gly Val Gly Met Val
900 905 910

Ser Gly Phe Gly Gln Ser Met Leu Val Asn Ser Ala Ile Thr Gln Gln
915 920 925

His Pro Gln Met Lys Gly Pro Val Gly Gln Ala Leu Pro Arg Pro Gln
930 935 940

Ala Pro Pro Arg Leu Gln Ser Leu Met Gly Thr Val Gln Gln Gly Ala
945 950 955 960

Gln Ser Trp Gln Gln Arg Ser Leu Gln Gly Met Pro Gly Arg Thr Ser
965 970 975

Gly Glu Leu Gly Pro Phe Asn Asn Gly Ala Ser Tyr Pro Leu Gln Ala
980 985 990

Gly Gln Pro Arg Leu Thr Lys Gln His Phe Pro Gln Gly Leu Ser Gln
995 1000 1005

Ser Val Val Asp Ala Asn Thr Gly Thr Val Arg Thr Leu Asn Pro
1010 1015 1020

Ala Ala Met Gly Arg Gln Met Met Pro Ser Leu Pro Gly Gln Gln
1025 1030 1035

Gly Thr Ser Gln Ala Arg Pro Met Val Met Ser Gly Leu Ser Gln
1040 1045 1050

Gly Val Pro Gly Met Pro Ala Phe Ser Gln Pro Pro Ala Gln Gln
1055 1060 1065

Gln Ile Pro Ser Gly Ser Phe Ala Pro Ser Ser Gln Ser Gln Ala
1070 1075 1080

Tyr Glu Arg Asn Ala Pro Gln Asp Val Ser Tyr Asn Tyr Ser Gly
1085 1090 1095

151

Asp Gly Ala Gly Gly Ser Phe Pro Gly Leu Pro Asp Gly Ala Asp
 1100 1105 1110

Leu Val Asp Ser Ile Ile Lys Gly Gly Pro Gly Asp Glu Trp Met
 1115 1120 1125

Gln Glu Leu Asp Glu Leu Phe Gly Asn Pro
 1130 1135

<210> 195
 <211> 30
 <212> PRT
 <213> Homo sapien

<400> 195

Met Gln Leu Pro Leu Ser His Lys Arg Lys Lys Gln Tyr Ser Phe Tyr
 1 5 10 15

Val Phe Asp Thr Asn Ile Lys His Asn Ser Val Leu Val His
 20 25 30

<210> 196
 <211> 46
 <212> PRT
 <213> Homo sapien

<400> 196

Met Lys Ile Tyr Phe Lys Ile Leu Leu Met Phe Leu Lys Lys Tyr Phe
 1 5 10 15

Leu Arg Phe His Leu Met Lys Thr Met Lys Tyr Ser Val Phe Tyr Ser
 20 25 30

Thr Thr Arg Gln Met Trp Ser Ile Pro Phe Val Phe Phe Phe
 35 40 45

<210> 197
 <211> 18
 <212> PRT
 <213> Homo sapien

<400> 197

Met Leu Glu Ala Gly Ile Ser Phe Lys Val Arg Leu Gln Lys Trp Lys
 1 5 10 15

152

Gln Ile

<210> 198
 <211> 132
 <212> PRT
 <213> Homo sapien

<400> 198

Met Phe Tyr Ser Ile Leu Ala Met Leu Arg Asp Arg Gly Ala Leu Gln
 1 5 10 15

Asp Leu Met Asn Met Leu Glu Leu Asp Ser Ser Gly His Leu Asp Gly
 20 25 30

Pro Gly Gly Ala Ile Leu Lys Lys Leu Gln Gln Asp Ser Asn His Ala
 35 40 45

Trp Phe Asn Pro Lys Asp Pro Ile Leu Tyr Leu Leu Glu Ala Ile Met
 50 55 60

Val Leu Ser Asp Phe Gln His Asp Leu Leu Ala Cys Ser Met Glu Lys
 65 70 75 80

Arg Ile Leu Leu Gln Gln Gln Glu Leu Val Arg Ser Ile Leu Glu Pro
 85 90 95

Asn Phe Arg Tyr Pro Trp Ser Ile Pro Phe Thr Leu Lys Pro Glu Leu
 100 105 110

Leu Ala Pro Leu Gln Ser Glu Gly Leu Ala Ser Pro Met Ala Ala Gly
 115 120 125

Gly Val Trp Pro
 130

<210> 199
 <211> 226
 <212> PRT
 <213> Homo sapien

<400> 199

Pro Pro Lys His Leu Lys Ser Lys Phe Gly Gly Met Arg Lys Ala Asp
 1 5 10 15

Asp Asp Leu Ile Leu Leu Leu Gly Arg Ile Glu Glu Pro Phe Trp Gln

153

20

25

30

Asn Phe Lys His Leu Gln Glu Val Phe Gln Lys Ile Lys Thr Leu
 35 40 45

Ala Gln Leu Ser Lys Asp Val Gln Asp Val Met Phe Tyr Ser Ile Leu
 50 55 60

Ala Met Leu Arg Asp Arg Gly Ala Leu Gln Asp Leu Met Asn Met Leu
 65 70 75 80

Glu Leu Asp Ser Ser Gly His Leu Asp Gly Pro Gly Gly Ala Ile Leu
 85 90 95

Lys Lys Leu Gln Gln Asp Ser Asn His Ala Trp Phe Asn Pro Lys Asp
 100 105 110

Pro Ile Leu Tyr Leu Leu Glu Ala Ile Met Val Leu Ser Asp Phe Gln
 115 120 125

His Asp Leu Leu Ala Cys Ser Met Glu Lys Arg Ile Leu Leu Gln Gln
 130 135 140

Gln Glu Leu Val Arg Ser Ile Leu Glu Pro Asn Phe Arg Tyr Pro Trp
 145 150 155 160

Ser Ile Pro Phe Thr Leu Lys Pro Glu Leu Leu Ala Pro Leu Gln Ser
 165 170 175

Glu Gly Leu Ala Ile Thr Tyr Gly Leu Leu Glu Glu Cys Gly Leu Arg
 180 185 190

Thr Glu Leu Asp Asn Pro Arg Ser Thr Trp Asp Val Glu Ala Lys Met
 195 200 205

Pro Leu Ser Ala Leu Tyr Gly Thr Leu Ser Leu Leu Gln Gln Leu Ala
 210 215 220

Glu Ala
 225

<210> 200
 <211> 37
 <212> PRT
 <213> Homo sapien

154

<400> 200

Met Ala Lys His Lys Gly Gly Tyr Gly Lys Tyr Trp Val Thr Leu Ile
 1 5 10 15

Ile Gly Leu Asn Ala Thr Asn Asn Ile Ile Ile Val Leu Thr Tyr Phe
 20 25 30

Phe Arg Leu Leu Ser
 35

<210> 201

<211> 28

<212> PRT

<213> Homo sapien

<400> 201

Met Val His Lys Ser Tyr Phe Thr Thr Leu Ser Leu Val Ile Leu Gly
 1 5 10 15

Val Trp Pro Cys Lys Ala Ser Ser Gln Arg Phe Cys
 20 25

<210> 202

<211> 77

<212> PRT

<213> Homo sapien

<400> 202

Met Gly Ser Val Cys Val Cys Phe His Arg Ser Thr Thr Ser Glu Val
 1 5 10 15

Ser Leu His Leu Cys Ile Phe Thr Ser Gln Gly Gln Gly Pro Gly Asn
 20 25 30

Leu Arg Gly Ser His Ser Phe Ser Leu Pro Gln Thr Met Pro Leu Pro
 35 40 45

Pro Ile Ser Leu Gly Gln Glu Ser Gly Phe Cys Phe Pro Tyr Phe Phe
 50 55 60

Phe Pro Arg His Trp Glu Ala Ser Gly Glu Gln His Gln
 65 70 75

<210> 203

<211> 70

155

<212> PRT

<213> Homo sapien

<400> 203

Met Gly Pro Pro Leu Pro Leu Gly Gly Trp Ser Ser Asp Leu Leu Ala
 1 5 10 15

Gln Lys Val Leu Phe Phe His Leu Leu Cys Leu Asn Glu Ser Ser Trp
 20 25 30

Thr Tyr Thr Pro Leu Ser Asp Glu Arg Ala Arg Leu Arg Arg Cys Ala
 35 40 45

Gly His Leu Leu Arg Ile His Val Gly Ser Ala Ala Pro Gly Gly Gly
 50 55 60

Ser Thr Ser Ala Ala Thr
 65 70

<210> 204

<211> 37

<212> PRT

<213> Homo sapien

<400> 204

Met Ser Lys Lys Lys Asp Gln Asp Leu Cys Leu Lys Ile Glu Met His
 1 5 10 15

Thr Ala Ala Ala Gln Lys Leu Arg Pro Ala Ser Lys Leu His Glu Ala
 20 25 30

Leu Val Lys Thr Asp
 35

<210> 205

<211> 87

<212> PRT

<213> Homo sapien

<400> 205

Met Pro Ser Val Ala Gln Gly Pro Val Pro Trp His Leu Gly Ser Arg
 1 5 10 15

Ser Ala Val Ala Val Phe Glu Phe Leu Val Met Phe Glu Gln Arg Pro
 20 25 30

156

Tyr Val Cys Ile Leu His Trp Ala Pro Gln Ile Thr Trp Pro Ile Leu
 35 40 45

Arg Arg Gly Val Ser His Leu Gln Ser Pro Lys Ser Pro Leu Glu Val
 50 55 60

Phe Leu Asn Glu Arg Thr Glu Ala Phe Leu Lys Ser Ser Val Gly Glu
 65 70 75 80

Thr Val His His His Thr Gln
 85

<210> 206
 <211> 46
 <212> PRT
 <213> Homo sapien

<400> 206

Met Ser Pro Gly Thr Ala Met Ala Leu Gly Ala Pro Thr Leu Phe Phe
 1 5 10 15

Phe Phe Phe Phe Phe Phe Tyr Asn Gln Pro Ile Arg Asp Leu Ser
 20 25 30

Ile Asn Lys Pro Leu Phe Ile Ile Arg Asn Trp Leu Thr Gln
 35 40 45

<210> 207
 <211> 91
 <212> PRT
 <213> Homo sapien

<400> 207

Met Ser Ser Pro Gln Ser Ile Glu His Asn His Asp Ser His Glu Leu
 1 5 10 15

Pro Thr Pro Pro Ala Ala Ser Ala Gln Arg Glu Ser Pro Leu Gln Val
 20 25 30

Cys Leu Ile Ala Glu Pro Ile Phe Phe Leu Pro Gly Gln Gln Leu Leu
 35 40 45

Ser Ser Met Ser Arg His Trp Cys Ser Leu Gly Trp Ala Pro Val Thr
 50 55 60

Pro Met Glu Ile Leu Asp Gly Cys Tyr Arg Thr Gly Leu Asp Val Arg

158

<210> 210
 <211> 58
 <212> PRT
 <213> Homo sapien

<400> 210

Met Val Ile Phe Tyr Ser Ser Pro Ser Gln Asp Ser Ala Leu Ile Tyr
 1 5 10 15

Tyr Ile Pro Phe Ile Leu Leu Tyr Arg Leu Leu Ser Glu Thr His Val
 20 25 30

Gln Ile Arg Asp Lys Ile Leu Lys His Ile Thr Pro Ser Leu Val Phe
 35 40 45

Ser Ile Gln Ile Leu Arg Asn Ser Cys Tyr
 50 55

<210> 211
 <211> 37
 <212> PRT
 <213> Homo sapien

<400> 211

Met Asn Leu Tyr Leu Lys Met Lys Thr Ile Pro Lys Lys Thr Cys Met
 1 5 10 15

Ser Lys Thr Glu Leu Phe Leu Pro Phe Thr Pro Lys Tyr Leu Lys Leu
 20 25 30

Asn Leu Ser His Phe
 35

<210> 212
 <211> 99
 <212> PRT
 <213> Homo sapien

<400> 212

Phe Phe Phe Phe Leu Arg Trp Ser Leu Ala Leu Ser Pro Arg Leu Glu
 1 5 10 15

Cys Ser Gly Val Ile Ser Thr His Cys Asn Leu Cys Phe Pro Gly Ser
 20 25 30

159

Ser Asp Ser Arg Ala Ser Pro Thr Phe Gln Val Ala Trp Ile Thr Gly
 35 40 45

Val Arg His His Ser Trp Leu Ile Phe Val Leu Leu Val Glu Thr Gly
 50 55 60

Phe His His Val Val Gln Ala Val Glu Leu Leu Thr Ser Arg Asp Pro
 65 70 75 80

Pro Ala Ser Ala Ser Gln Ser Ala Ala Ile Ile Gly Val Asn His Cys
 85 90 95

Ala Arg Pro

<210> 213
 <211> 43
 <212> PRT
 <213> Homo sapien

<400> 213

Met Gln Glu Phe Thr Trp Leu Phe Glu Lys Glu Asn Phe Lys Val Ser
 1 5 10 15

Gly Trp Thr Glu Ser His Glu Ala Arg Ser Leu Leu Thr Ala Arg Ser
 20 25 30

Leu Glu Lys Gln Val Ser Gly Ser Phe Thr Ser
 35 40

<210> 214
 <211> 61
 <212> PRT
 <213> Homo sapien

<400> 214

Met Ala Val Asp Phe Tyr Asn Phe Val Thr Lys Leu Val Val Thr Thr
 1 5 10 15

Gly Tyr Leu Arg Ile Ser Phe Leu Ala Tyr Lys Phe Phe Ser Phe Pro
 20 25 30

Phe Leu Asp Ser Leu Ser Leu Cys Cys Pro Gly Leu Glu Cys Ser Gly
 35 40 45

Val Ile Pro Ala His Tyr Asn Leu Tyr Leu Pro Gly Arg

160

50

55

60

<210> 215
 <211> 127
 <212> PRT
 <213> Homo sapien
 <400> 215

Ser Gln Asn Ile Phe Phe Gly Val Ala Ile Phe Phe Phe Ser Phe Phe
 1 5 10 15

Arg Gln Ser Leu Ser Leu Val Ala Gln Ala Arg Val Gln Trp Arg Asp
 20 25 30

Pro Gly Ser Leu Gln Pro Leu Pro Pro Gly Phe Lys Arg Phe Leu Gly
 35 40 45

Leu Ser Leu Pro Ser Ser Ala Gly Tyr Arg Arg Ala Pro Pro Pro Cys
 50 55 60

Pro Ala Leu Leu Tyr Phe Ala Val Glu Thr Gly Phe His His Val Gly
 65 70 75 80

Gln Ala Gly Leu Glu Leu Leu Thr Ser Gly Asn Pro Ala Ala Ser Ala
 85 90 95

Ser Gln Ser Ala Gly Ile Thr Gly Thr Ser His Cys Thr Gln Pro Tyr
 100 105 110

Tyr His Lys Ser Ser Ala Cys Trp Tyr Leu Ile Arg Phe Tyr Leu
 115 120 125

<210> 216
 <211> 13
 <212> PRT
 <213> Homo sapien
 <400> 216

Met Glu Cys Ser Ser Leu Ala Glu Phe Lys Pro Val Phe
 1 5 10

<210> 217
 <211> 100
 <212> PRT
 <213> Homo sapien
 <400> 217

161

Pro Gln Gln Thr Leu Lys Arg Ile Gln Gln Val Leu Ile Lys Cys Cys
 1 5 10 15

Leu Ala Phe Tyr Leu Phe Leu Phe Phe Phe Leu Arg Trp Ser Leu
 20 25 30

Ala Leu Leu Pro Ser Leu Lys Cys Ser Gly Val Ile Ser Ala His Cys
 35 40 45

Asn Leu Arg Leu Pro Gly Leu Gly Asp Ser Leu Ala Ser Ala Ser Arg
 50 55 60

Val Ala Gly Met Thr Thr Gly Thr Cys His His Ala Gln Leu Ile Phe
 65 70 75 80

Val Phe Leu Val Glu Thr Gly Phe Cys Val Ser Gln Asp Gly Leu Asp
 85 90 95

Leu Leu Ile Ser
 100

<210> 218
 <211> 46
 <212> PRT
 <213> Homo sapien

<400> 218

Met Glu Gly Gly Glu Met Ser Thr Gln Val Glu Asn Arg Ser Glu Gly
 1 5 10 15

Thr Ile Pro Ile Gln Thr Thr Cys Lys Ser His Asn Lys Ala Pro His
 20 25 30

Cys Thr Glu Leu Arg His Lys Gln Arg Phe Pro Thr Asp Gly
 35 40 45

<210> 219
 <211> 72
 <212> PRT
 <213> Homo sapien

<400> 219

Ile Ser Phe Ile Phe Phe Ser Glu Ala Cys Gln Val Glu Val Arg Leu
 1 5 10 15

162

Leu Leu Ala Tyr Asn Ser Ser Ala Arg Ile Pro Lys Cys Pro Trp Met
 20 25 30

Glu Gly Gly Glu Met Ser Pro Gln Val Glu Thr Ser Ile Glu Gly Thr
 35 40 45

Ile Pro Phe Ser Lys Pro Val Lys Val Tyr Ile Met Pro Lys Pro Ala
 50 55 60

Arg Arg Pro Lys Pro Ala Arg Arg
 65 70

<210> 220
 <211> 41
 <212> PRT
 <213> Homo sapien

<400> 220

Met Glu Cys Lys Val Ile Lys Cys Ser Cys Phe His Leu Glu Gly Cys
 1 5 10 15

Gly Pro Glu Gly Lys Arg Ser Pro Lys Tyr Pro Pro Pro Trp Cys Ser
 20 25 30

Ser Leu Cys Leu Val Pro Ala Arg Ala
 35 40

<210> 221
 <211> 30
 <212> PRT
 <213> Homo sapien

<400> 221

Met Asn Ser Phe Gly Tyr Met Thr Pro Ser Lys Phe Phe Lys Lys Glu
 1 5 10 15

Ile Thr Phe Lys Thr Thr Tyr Ile Phe Cys Phe Cys Leu Arg
 20 25 30

<210> 222
 <211> 22
 <212> PRT
 <213> Homo sapien

<400> 222

Met Leu Gln Ile Gly His Leu Leu Ser Met His Ser Leu Asp Lys Asn
 1 5 10 15

163

Ile Gly Gln Val Gly Met
20

<210> 223
<211> 18
<212> PRT
<213> Homo sapien

<400> 223

Met Ser Asp Arg Val Val Ala Leu Leu Glu Val Phe Phe Pro Phe Gln
1 5 10 15

Arg Glu

<210> 224
<211> 133
<212> PRT
<213> Homo sapien

<400> 224

Met Gly Asn Ser Ile Asp Thr Val Arg Tyr Gly Lys Glu Ser Asp Leu
1 5 10 15

Gly Asp Val Ser Glu Glu His Gly Glu Trp Asn Lys Glu Ser Ser Asn
20 25 30

Asn Glu Gln Asp Asn Ser Leu Leu Glu Gln Tyr Leu Thr Ser Val Gln
35 40 45

Gln Leu Glu Asp Ala Asp Glu Arg Thr Asn Phe Asp Thr Glu Thr Arg
50 55 60

Asp Ser Lys Leu His Ile Ala Cys Phe Pro Val Gln Leu Asp Thr Leu
65 70 75 80

Ser Asp Gly Ala Ser Val Asp Glu Ser His Gly Ile Ser Pro Pro Leu
85 90 95

Gln Gly Glu Ile Ser Gln Thr Gln Glu Asn Ser Lys Leu Asn Ala Glu
100 105 110

Val Gln Gly Gln Gln Pro Glu Cys Asp Ser Thr Phe Gln Leu Leu His
115 120 125

Val Gly Val Thr Val
130

<210> 225
<211> 50
<212> PRT
<213> Homo sapien

<400> 225

Met Arg Asn Ser Ser Pro Ile Leu Thr Pro Ala Leu Phe Ser Phe His
1 5 10 15

Met Tyr Ile Gly Pro Leu Ile Arg Ile Phe Lys Lys Phe Pro Arg Pro
20 25 30

Pro Asn Leu Thr Ile Asp Asp Pro Leu Ser Leu Phe Arg Arg Asn Tyr
35 40 45

Ile Gly
50

<210> 226
<211> 43
<212> PRT
<213> Homo sapien

<400> 226

Met His Ser Phe Phe Leu Ser Met Leu Cys Pro Glu Ala Leu Arg Val
1 5 10 15

Leu Leu Lys Gln Ala Ala Gly Leu Leu Arg Glu Ile Lys Gly Phe Ile
20 25 30

Ser Thr Thr Arg Cys Gln Asn Leu His Phe Glu
35 40

<210> 227
<211> 99
<212> PRT
<213> Homo sapien

<400> 227

Met Leu Glu Arg Arg Ser Val Met Asp Arg Arg Arg Ala Gly Asn Ser
1 5 10 15

Pro Pro Arg Ile Glu Lys Cys Leu Leu Gly Arg Glu Glu Gly Glu Ala

165

20 25 30

Gly Ala Gly Pro Ser Pro Gly Ser Leu Leu Gly Pro Gln Lys Ala Leu
 35 40 45

Asn Gln Ala Pro Ser Leu Gln Gly Lys Pro Arg Pro Gln Pro Asp Asn
 50 55 60

Leu Glu Gly Arg Lys Ser Gln Thr Leu Gly Leu Phe Phe Gly Gly Ile
 65 70 75 80

Ile Gly Phe Phe Phe Phe Met Phe Leu Leu Glu Phe Cys Leu Leu Ala
 85 90 95

Asn Ser Val

<210> 228
 <211> 44
 <212> PRT
 <213> Homo sapien

<400> 228

Met Lys Ser Ile Gln Leu Lys Phe Ser Tyr Ile Ile Glu Pro Gln Leu
 1 5 10 15

Asn Gly Met Asn Gly Ile Gly Asn Leu Leu Glu Met Ile Phe Met Ile
 20 25 30

Thr Phe Val Val Ile Pro Phe Ser Trp Leu Arg Phe
 35 40

<210> 229
 <211> 41
 <212> PRT
 <213> Homo sapien

<400> 229

Tyr Phe Pro Leu Gln Ile Trp Ile Ser Glu Asp Ser Asn Asn Ile Glu
 1 5 10 15

Ala Val Asn Gln Trp Lys Glu Thr Val Ile Asn Pro Glu Lys Val Val
 20 25 30

Ile Arg Trp His Lys Leu Asn Pro Ser
 35 40

166

<210> 230
 <211> 48
 <212> PRT
 <213> Homo sapien
 <400> 230

Met Leu Lys Gly His Tyr Gln Tyr Gly Met Glu Asp Leu Ser Phe His
 1 5 10 15

Thr Phe Ser Ser Ser Phe Leu Asn Phe Leu Leu Leu Phe Leu Leu Ser
 20 25 30

Cys Met Val Ala Pro Phe Pro Phe Leu Leu Ser Val Pro Ser Lys Gln
 35 40 45

<210> 231
 <211> 108
 <212> PRT
 <213> Homo sapien
 <400> 231

Phe Leu Lys Arg Gln Ser Ile Ser Leu Leu Pro Gln Leu Glu Cys Ser
 1 5 10 15

Gly Thr Ile Ile Val His His Thr Leu Glu Leu Leu Gly Lys Gly Ser
 20 25 30

Ser Leu Ala Ser Ala Ser Gln Val Ala Arg Tyr Thr Gly Met Cys Tyr
 35 40 45

His Ala Trp Leu Ile Lys Lys Ile Phe Leu Glu Met Arg Ser Cys Cys
 50 55 60

Val Ala Gln Ala Gly Leu Lys Leu Leu Gly Ser Asn Asn Pro Pro Thr
 65 70 75 80

Leu Ala Ser Gln Ser Ala Gly Ile Thr Gly Val Ser His Ser Thr Ala
 85 90 95

Pro Tyr Leu Gln Ile Leu Asn Gln Ala Ile Ala Ile
 100 105

<210> 232
 <211> 64
 <212> PRT

167

<213> Homo sapien

<400> 232

Met Ser Pro Arg Ala Pro Phe Ala Pro Gly Cys Pro Gln Pro Leu Val
 1 5 10 15

Val Phe Tyr Val Cys Phe Phe Phe Phe Leu Ile Phe Cys Phe Val Lys
 20 25 30

Lys His His Tyr Met Phe Leu Tyr Pro Arg Leu Lys Thr Phe Gly Asn
 35 40 45

Leu Ile Ser Asn Ile Lys Ile Gln Ile Lys Thr His Ser Thr Ile Pro
 50 55 60

<210> 233

<211> 35

<212> PRT

<213> Homo sapien

<400> 233

Met Cys Val Asn Ala Ser Thr Val Gly Gln Met Cys Glu Asn Glu Leu
 1 5 10 15

Lys His Met Leu Arg Ile Lys Val Asn Arg Arg Asn Phe Glu Arg Phe
 20 25 30

Pro Leu Met
 35

<210> 234

<211> 72

<212> PRT

<213> Homo sapien

<400> 234

Met Asn Ile Phe Pro Trp Ala Gly Gly Pro Trp Ser Leu Pro Gln Ala
 1 5 10 15

Arg Tyr Arg Ala Pro Ala Cys Ala Pro Thr Asn His Gly Lys Gln Arg
 20 25 30

Arg Pro Pro His Leu Lys Ser Trp Pro Val Val Val Ser Ser Val Phe
 35 40 45

Leu Leu Ser Glu Gln Asn Val Leu Lys Leu Glu Leu Thr Lys Val Lys

168

50

55

60

Ser Ser Lys Thr Thr Tyr Ala Thr
65 70

<210> 235
<211> 1163
<212> PRT
<213> Homo sapien

<400> 235

Met Asp Arg Asn Arg Glu Ala Glu Met Glu Leu Arg Arg Gly Pro Ser
1 5 10 15

Pro Thr Arg Ala Gly Arg Gly His Glu Val Asp Gly Asp Lys Ala Thr
20 25 30

Cys His Thr Cys Cys Ile Cys Gly Lys Ser Phe Pro Phe Gln Ser Ser
35 40 45

Leu Ser Gln His Met Arg Lys His Thr Gly Glu Lys Pro Tyr Lys Cys
50 55 60

Pro Tyr Cys Asp His Arg Ala Ser Gln Lys Gly Asn Leu Lys Ile His
65 70 75 80

Ile Arg Ser His Arg Thr Gly Thr Leu Ile Gln Gly His Glu Pro Glu
85 90 95

Ala Gly Glu Ala Pro Leu Gly Glu Met Arg Ala Ser Glu Gly Leu Asp
100 105 110

Ala Cys Ala Ser Pro Thr Lys Ser Ala Ser Ala Cys Asn Arg Leu Leu
115 120 125

Asn Gly Ala Ser Gln Ala Asp Gly Ala Arg Val Leu Asn Gly Ala Ser
130 135 140

Gln Ala Asp Ser Gly Arg Val Leu Leu Arg Ser Ser Lys Lys Gly Ala
145 150 155 160

Glu Gly Ser Ala Cys Ala Pro Gly Glu Ala Lys Ala Ala Val Gln Cys
165 170 175

Ser Phe Cys Lys Ser Gln Phe Glu Arg Lys Lys Asp Leu Glu Leu His

169

		180						185						190			
Val	His	Gln	Ala	His	Lys	Pro	Phe	Lys	Cys	Arg	Leu	Cys	Ser	Tyr	Ala		
		195					200					205					
Thr	Leu	Arg	Glu	Glu	Ser	Leu	Leu	Ser	His	Ile	Glu	Arg	Asp	His	Ile		
	210					215					220						
Thr	Ala	Gln	Gly	Pro	Gly	Ser	Gly	Glu	Ala	Cys	Val	Glu	Asn	Gly	Lys		
225					230					235					240		
Pro	Glu	Leu	Ser	Pro	Gly	Glu	Phe	Pro	Cys	Glu	Val	Cys	Gly	Gln	Ala		
				245					250					255			
Phe	Ser	Gln	Thr	Trp	Phe	Leu	Lys	Ala	His	Met	Lys	Lys	His	Arg	Gly		
			260					265					270				
Ser	Phe	Asp	His	Gly	Cys	His	Ile	Cys	Gly	Arg	Arg	Phe	Lys	Glu	Pro		
		275					280					285					
Trp	Phe	Leu	Lys	Asn	His	Met	Lys	Ala	His	Gly	Pro	Lys	Thr	Gly	Ser		
	290					295					300						
Lys	Asn	Arg	Pro	Lys	Ser	Glu	Leu	Asp	Pro	Ile	Ala	Thr	Ile	Asn	Asn		
305					310					315					320		
Val	Val	Gln	Glu	Glu	Val	Ile	Val	Ala	Gly	Leu	Ser	Leu	Tyr	Glu	Val		
				325					330					335			
Cys	Ala	Lys	Cys	Gly	Asn	Leu	Phe	Thr	Asn	Leu	Asp	Ser	Leu	Asn	Ala		
			340					345					350				
His	Asn	Ala	Ile	His	Arg	Arg	Val	Glu	Ala	Ser	Arg	Thr	Arg	Ala	Pro		
		355					360					365					
Ala	Glu	Glu	Gly	Ala	Glu	Gly	Pro	Ser	Asp	Thr	Lys	Gln	Phe	Phe	Leu		
	370					375					380						
Gln	Cys	Leu	Asn	Leu	Arg	Pro	Ser	Ala	Ala	Gly	Asp	Ser	Cys	Pro	Gly		
385					390					395					400		
Thr	Gln	Ala	Gly	Arg	Arg	Val	Ala	Glu	Leu	Asp	Pro	Val	Asn	Ser	Tyr		
				405					410					415			

170

Gln Ala Trp Gln Leu Ala Thr Arg Gly Lys Val Ala Glu Pro Ala Glu
 420 425 430

Tyr Leu Lys Tyr Gly Ala Trp Asp Glu Ala Leu Ala Gly Asp Val Ala
 435 440 445

Phe Asp Lys Asp Arg Arg Glu Tyr Val Leu Val Ser Gln Glu Lys Arg
 450 455 460

Lys Arg Glu Gln Asp Ala Pro Ala Ala Gln Gly Pro Pro Arg Lys Arg
 465 470 475 480

Ala Ser Gly Pro Gly Asp Pro Ala Pro Ala Gly His Leu Asp Pro Arg
 485 490 495

Ser Ala Ala Arg Pro Asn Arg Arg Ala Ala Ala Thr Thr Gly Gln Gly
 500 505 510

Lys Ser Ser Glu Cys Phe Glu Cys Gly Lys Ile Phe Arg Thr Tyr His
 515 520 525

Gln Met Val Leu His Ser Arg Val His Arg Arg Ala Arg Arg Glu Arg
 530 535 540

Asp Ser Asp Gly Asp Arg Ala Ala Arg Ala Arg Cys Gly Ser Leu Ser
 545 550 555 560

Glu Gly Asp Ser Ala Ser Gln Pro Ser Ser Pro Gly Ser Ala Cys Ala
 565 570 575

Ala Ala Asp Ser Pro Gly Ser Gly Leu Ala Asp Glu Ala Ala Glu Asp
 580 585 590

Ser Gly Glu Glu Gly Ala Pro Glu Pro Ala Pro Gly Gly Gln Pro Arg
 595 600 605

Arg Cys Cys Phe Ser Glu Glu Val Thr Ser Thr Glu Leu Ser Ser Gly
 610 615 620

Asp Gln Ser His Lys Met Gly Asp Asn Ala Ser Glu Arg Asp Thr Gly
 625 630 635 640

Glu Ser Lys Ala Gly Ile Ala Ala Ser Val Ser Ile Leu Glu Asn Ser
 645 650 655

171

Ser Arg Glu Thr Ser Arg Arg Gln Glu Gln His Arg Phe Ser Met Asp
660 665 670

Leu Lys Met Pro Ala Phe His Pro Lys Gln Glu Val Pro Val Pro Gly
675 680 685

Asp Gly Val Glu Phe Pro Ser Ser Thr Gly Ala Glu Gly Gln Thr Gly
690 695 700

His Pro Ala Glu Lys Leu Ser Asp Leu His Asn Lys Glu His Ser Gly
705 710 715 720

Gly Gly Lys Arg Ala Leu Ala Pro Asp Leu Met Pro Leu Asp Leu Ser
725 730 735

Ala Arg Ser Thr Arg Asp Asp Pro Ser Asn Lys Glu Thr Ala Ser Ser
740 745 750

Leu Gln Ala Ala Leu Val Val His Pro Cys Pro Tyr Cys Ser His Lys
755 760 765

Thr Tyr Tyr Pro Glu Val Leu Trp Met His Lys Arg Ile Trp His Arg
770 775 780

Val Ser Cys Asn Ser Val Ala Pro Pro Trp Ile Gln Pro Asn Gly Tyr
785 790 795 800

Lys Ser Ile Arg Ser Asn Leu Val Phe Leu Ser Arg Ser Gly Arg Thr
805 810 815

Gly Pro Pro Pro Ala Leu Gly Gly Lys Glu Cys Gln Pro Leu Leu Leu
820 825 830

Ala Arg Phe Thr Arg Thr Gln Val Pro Gly Gly Met Pro Gly Ser Lys
835 840 845

Ser Gly Ser Ser Pro Leu Gly Val Val Thr Lys Ala Ala Ser Met Pro
850 855 860

Lys Asn Lys Glu Ser His Ser Gly Gly Pro Cys Ala Leu Trp Ala Pro
865 870 875 880

Gly Pro Asp Gly Tyr Arg Gln Thr Lys Pro Cys His Gly Gln Glu Pro
885 890 895

172

His Gly Ala Ala Thr Gln Gly Pro Leu Ala Lys Pro Arg Gln Glu Ala
 900 905 910

Ser Ser Lys Pro Val Pro Ala Pro Gly Gly Gly Gly Phe Ser Arg Ser
 915 920 925

Ala Thr Pro Thr Pro Thr Val Ile Ala Arg Ala Gly Ala Gln Pro Ser
 930 935 940

Ala Asn Ser Lys Pro Val Glu Lys Phe Gly Val Pro Pro Ala Gly Ala
 945 950 955 960

Gly Phe Ala Pro Thr Asn Lys His Ser Ala Pro Asp Ser Leu Lys Ala
 965 970 975

Lys Phe Ser Ala Gln Pro Gln Gly Pro Pro Pro Ala Lys Gly Glu Gly
 980 985 990

Gly Ala Pro Pro Leu Pro Pro Arg Glu Pro Pro Ser Lys Ala Ala Gln
 995 1000 1005

Glu Leu Arg Thr Leu Ala Thr Cys Ala Ala Gly Ser Arg Gly Asp
 1010 1015 1020

Ala Ala Leu Gln Ala Gln Pro Gly Val Ala Gly Ala Pro Pro Val
 1025 1030 1035

Leu His Ser Ile Lys Gln Glu Pro Val Ala Glu Gly His Glu Lys
 1040 1045 1050

Arg Leu Asp Ile Leu Asn Ile Phe Lys Thr Tyr Ile Pro Lys Asp
 1055 1060 1065

Phe Ala Thr Leu Tyr Gln Gly Trp Gly Val Ser Gly Pro Gly Leu
 1070 1075 1080

Glu His Arg Gly Thr Leu Arg Thr Gln Ala Arg Pro Gly Glu Phe
 1085 1090 1095

Val Cys Ile Glu Cys Gly Lys Ser Phe His Gln Pro Gly His Leu
 1100 1105 1110

Arg Ala His Met Arg Ala His Ser Val Val Phe Glu Ser Asp Gly

173

1115 1120 1125

Pro Arg Gly Ser Glu Val His Thr Thr Ser Ala Asp Ala Pro Lys
 1130 1135 1140

Gln Gly Arg Asp His Ser Asn Thr Gly Thr Val Gln Thr Val Pro
 1145 1150 1155

Leu Arg Lys Gly Thr
 1160

<210> 236
 <211> 55
 <212> PRT
 <213> Homo sapien

<400> 236

Met Cys Val Phe Cys Gly Phe Phe Cys Ser Arg Phe Val Arg Glu Met
 1 5 10 15

Trp Gly Asn Phe Gly Pro Lys Thr Asn Phe Thr Pro Gly Thr Pro Phe
 20 25 30

Cys Pro Trp Leu Ser Pro Asn Leu Phe Cys Leu Val Val Val Trp Phe
 35 40 45

Tyr Arg Leu Leu Ile Phe Tyr
 50 55

<210> 237
 <211> 156
 <212> PRT
 <213> Homo sapien

<400> 237

Met Pro Met Glu Gly His Thr Leu Cys Met Arg Ile Arg Gly Ser Trp
 1 5 10 15

Leu Ala Ala Arg Leu Pro Val Met Pro Phe Glu Gly Asp Val Gly Pro
 20 25 30

Trp Val Arg Met Lys Val Phe Ile Cys His Ser Ser Ser Pro Gln Val
 35 40 45

Ala Ile His Leu Gly Gly Gly Arg Glu Gly Ser Ala Leu Ala Ile Val
 50 55 60

174

Tyr Pro Ala Ser Leu Arg Phe Ile Asp Leu His Lys Arg Leu Cys Ser
65 70 75 80

Gly Lys Gly Arg Gly Pro Gln Lys Gly Ala Trp Gln Asp Arg Trp Met
85 90 95

Leu Tyr Gly His Met Glu Ile Thr Pro Ser Ser Leu Ala Pro Ala Ser
100 105 110

Ala Ser Arg Pro Leu His Gly Val Arg Cys Phe Cys Ala Cys Cys Pro
115 120 125

Thr Ser Leu His Ser Arg Ala Leu Ile Asn His Phe Asp Pro Pro Leu
130 135 140

Ala Glu Gly Ser Pro Leu Tyr Arg Val Gln Ser Leu
145 150 155

<210> 238
<211> 86
<212> PRT
<213> Homo sapien

<400> 238

Met Met Asn Phe Leu Cys Leu Asn Phe Arg Asp Ile Trp Cys Asp Phe
1 5 10 15

His Leu Tyr Leu Met Leu Pro Leu Leu Pro Ser Leu Leu Asn Thr Ser
20 25 30

Lys Asn Ser Glu His Ile Leu Ile Pro Pro Val Phe Tyr Phe Tyr Asp
35 40 45

Leu Asp Ile Leu His His Lys Ile Pro Pro Asn Trp Asp Tyr Val Phe
50 55 60

Glu Val Ile His Phe Thr Ile Ile Thr Thr Ile Thr Ile Ile Phe Ile
65 70 75 80

Val Cys Phe Val Pro Gly
85

<210> 239
<211> 289

175

<212> PRT

<213> Homo sapien

<400> 239

Ala Asp Leu Ser Phe Ile Glu Asp Thr Val Ala Phe Pro Glu Lys Glu
1 5 10 15

Glu Asp Glu Glu Glu Glu Glu Gly Val Glu Trp Gly Tyr Glu Glu
20 25 30

Gly Val Glu Trp Gly Leu Val Phe Pro Asp Ala Asn Gly Glu Tyr Gln
35 40 45

Ser Pro Ile Asn Leu Asn Ser Arg Glu Ala Arg Tyr Asp Pro Ser Leu
50 55 60

Leu Asp Val Arg Leu Ser Pro Asn Tyr Val Val Cys Arg Asp Cys Glu
65 70 75 80

Val Thr Asn Asp Gly His Thr Ile Gln Val Ile Leu Lys Ser Lys Ser
85 90 95

Val Leu Ser Gly Gly Pro Leu Pro Gln Gly His Glu Phe Glu Leu Tyr
100 105 110

Glu Val Arg Phe His Trp Gly Arg Glu Asn Gln Arg Gly Ser Glu His
115 120 125

Thr Val Asn Phe Lys Ala Phe Pro Met Glu Leu His Leu Ile His Trp
130 135 140

Asn Ser Thr Leu Phe Gly Ser Ile Asp Glu Ala Val Gly Lys Pro His
145 150 155 160

Gly Ile Ala Ile Ile Ala Leu Phe Val Gln Ile Gly Lys Glu His Val
165 170 175

Gly Leu Lys Ala Val Thr Glu Ile Leu Gln Asp Ile Gln Tyr Lys Gly
180 185 190

Lys Ser Lys Thr Ile Pro Cys Phe Asn Pro Asn Thr Leu Leu Pro Asp
195 200 205

Pro Leu Leu Arg Asp Tyr Trp Val Tyr Glu Gly Ser Leu Thr Ile Pro
210 215 220

176

Pro Cys Ser Glu Gly Val Thr Trp Ile Leu Phe Arg Tyr Pro Leu Thr
 225 230 235 240

Ile Ser Gln Leu Gln Ile Glu Glu Phe Arg Arg Leu Arg Thr His Val
 245 250 255

Lys Gly Ala Glu Leu Val Glu Gly Cys Asp Gly Ile Leu Gly Asp Asn
 260 265 270

Phe Arg Pro Thr Gln Pro Leu Ser Asp Arg Val Ile Arg Ala Ala Phe
 275 280 285

Gln

<210> 240
 <211> 59
 <212> PRT
 <213> Homo sapien

<400> 240

Met Cys Gln Ile Asp Arg Gln Asp Leu Val Leu Leu Lys Leu Val Ile
 1 5 10 15

Tyr Cys Ser Arg His Leu Lys Gly Trp Arg Arg Ser Glu His Tyr Val
 20 25 30

Pro Ala Arg Ala Ser Ile Thr Leu Arg Arg Ser Thr Ser His Leu Val
 35 40 45

Ala Arg Ser Pro Asn Met Ser Ser Ser Gly Val
 50 55

<210> 241
 <211> 41
 <212> PRT
 <213> Homo sapien

<400> 241

Met Leu Leu Asn Gly Leu His Asn Pro Ala Leu Lys His Leu Arg Asp
 1 5 10 15

Leu Cys Lys Thr Phe Pro Trp Ser Leu Cys Phe Ser His Ile Asn Gln
 20 25 30

177

Leu Ala Tyr Phe Ser His Ser Pro Ser
 35 40

<210> 242
 <211> 80
 <212> PRT
 <213> Homo sapien

<400> 242

Met Asn Cys Leu Tyr Pro Ser Pro Met Cys Phe Tyr Arg Ser Cys Leu
 1 5 10 15

Val His Phe Val Ala Asp Leu Leu Gly Asp Phe Thr Glu Gly Lys Val
 20 25 30

Ser Ser Lys Leu Tyr Asp Asp Phe Met Leu Ile Asp Leu Leu Ser Ser
 35 40 45

Gly Ser Trp Glu Thr His Ser Ala Ile Ser Leu Leu Ser Tyr Phe Ser
 50 55 60

Tyr Asp Ala Gln Pro Pro Lys Ala Thr Arg Glu Gln Tyr Arg Val Pro
 65 70 75 80

<210> 243
 <211> 45
 <212> PRT
 <213> Homo sapien

<400> 243

Glu Arg Pro Gly Met Leu Asp Phe Thr Gly Lys Ala Lys Trp Asp Ala
 1 5 10 15

Trp Asn Glu Leu Lys Gly Thr Ser Lys Glu Asp Ala Met Lys Ala Tyr
 20 25 30

Ile Asn Lys Val Glu Glu Leu Lys Lys Lys Tyr Gly Ile
 35 40 45

<210> 244
 <211> 24
 <212> PRT
 <213> Homo sapien

<400> 244

Met Cys Leu Asn Phe Ser Phe Asn Tyr Leu Ile Pro Phe Ala Gln Glu

178

1 5 10 15

Ile Thr Ile Ser Leu Phe Phe Phe
20

<210> 245
<211> 69
<212> PRT
<213> Homo sapien

<400> 245

Leu Phe Phe Gln Leu Phe Asp Thr Phe Cys Pro Arg Asp Tyr Tyr Leu
1 5 10 15

Ser Leu Phe Phe Phe Ser Phe Lys Thr Glu Cys Cys Ser Val Thr Gln
20 25 30

Val Gly Val Gln Trp His Asn Ser Ala Ser Leu Gln Pro Leu Pro Pro
35 40 45

Arg Leu Lys Arg Ser Ser His Leu Ser Leu Pro Ser Ser Trp Asp His
50 55 60

Arg His Ile Pro Pro
65

<210> 246
<211> 39
<212> PRT
<213> Homo sapien

<400> 246

Met Glu Thr Lys His His Ser His Lys Lys Ser Asn Ser Ile Leu Asn
1 5 10 15

His Trp Lys Val Thr Ile Pro Leu Tyr Ser Phe Pro Lys Leu Phe Val
20 25 30

Ala Lys Ser Tyr Arg Lys Glu
35

<210> 247
<211> 93
<212> PRT
<213> Homo sapien

<400> 247

179

Leu Leu Gln Ala Leu Lys Lys Ile Phe Phe Leu Asn Ser Leu Thr Leu
 1 5 10 15

Ser Pro Arg Leu Glu Ala Ser Asn Val Ile Ser Ala His Cys Asn Leu
 20 25 30

His Ser Arg Val Ala Gly Ile Thr Asp Met His His His Pro Gln Leu
 35 40 45

Ile Phe Val Phe Leu Val Glu Thr Gly Phe Arg His Val Gly Gln Ala
 50 55 60

Gly Leu Ala Leu Leu Ala Leu Arg Asp Pro Pro Pro Leu Ala Phe Gln
 65 70 75 80

Ser Ala Gly Ile Thr Gly Val Ser His Cys Thr Trp Pro
 85 90

<210> 248
 <211> 51
 <212> PRT
 <213> Homo sapien

<400> 248

Met Phe Phe Phe Phe Val Phe Phe Phe Phe Leu Phe Ala Arg Phe Ser
 1 5 10 15

Arg Asn Val Gly Asp Leu Trp Ala Gly Lys Pro Phe Pro Pro Gly His
 20 25 30

Val Leu Pro Arg Tyr Pro His Leu Phe Phe Phe Phe Phe Phe Cys
 35 40 45

Phe Ile Thr
 50

<210> 249
 <211> 62
 <212> PRT
 <213> Homo sapien

<400> 249

Met Asn Phe Thr Leu Ala Ile Phe His Tyr Phe Ser Leu Ser Gln Met
 1 5 10 15

180

Ser Val Leu Met Arg Gln Leu Ala Leu Thr Gly Ala Thr Leu Met Cys
 20 25 30

His Leu Pro Thr Phe Asn Phe Trp Val Lys Ala Glu Arg Glu Lys Leu
 35 40 45

Met Asp Phe Ser Phe Ser Arg Arg Asp Lys Asn Gln Leu His
 50 55 60

<210> 250
 <211> 190
 <212> PRT
 <213> Homo sapien
 <400> 250

Met Lys Leu Gln Leu Arg Ile Lys Ser Leu Thr Gln Asn Arg Thr Thr
 1 5 10 15

Thr Trp Lys Leu Asn Asn Leu Leu Leu Asn Asp Tyr Trp Val Asn Lys
 20 25 30

Lys Ile Lys Ala Glu Ile Asn Lys Phe Phe Glu Thr Ile Glu Asn Lys
 35 40 45

Asp Thr Met Tyr Gln Asn Thr Ala Lys Ala Val Phe Arg Gly Lys Phe
 50 55 60

Ile Ala Leu Asn Thr His Ile Arg Asn Trp Glu Ile Pro Lys Ile Asn
 65 70 75 80

Val Leu Thr Ser Gln Leu Lys Glu Leu Glu Lys Arg Glu Gln Thr His
 85 90 95

Ser Lys Gln Glu Ile Thr Lys Ile Ile Ala Glu Leu Lys Glu Ile Glu
 100 105 110

Thr Gln Lys Ala Leu Gln Lys Ile Ser Asp Ser Arg Ser Trp Phe Phe
 115 120 125

Glu Lys Ile Asn Lys Thr Asp Arg Leu Leu Ala Arg Ile Ile Lys Lys
 130 135 140

Lys Arg Glu Lys Asn Gln Ile Asp Thr Ile Lys Asn Asp Lys Gly Asp
 145 150 155 160

181

Ile Thr Thr Asn Pro Thr Glu Ile Gln Thr Ala Ile Arg Glu Cys Tyr
 165 170 175

Gln His Leu Tyr Ile Asn Lys Leu Glu Asn Leu Glu Glu Ile
 180 185 190

<210> 251
 <211> 132
 <212> PRT
 <213> Homo sapien

<400> 251

Met Pro Val Leu Ser Pro Pro Leu His Met Pro Tyr Pro Ala Ala Lys
 1 5 10 15

Leu Asp Ser Val Leu Pro Asp Lys Thr Trp Tyr Trp His Leu Tyr Ala
 20 25 30

Ser Val Cys Leu Pro Ser Thr Phe Lys Lys Pro Leu Gln Ser Ala Asp
 35 40 45

Thr Lys Lys Gln Ser His Thr Cys Ser Lys Ser Ala Cys Phe Pro Leu
 50 55 60

Ile Ser Ala Ser Cys Gln Arg His Cys Leu Thr Ser Ser Ser Leu Leu
 65 70 75 80

Ser Ile Cys Val Pro His Lys Thr Leu Arg Asp Ser Ala Ser Tyr Val
 85 90 95

Tyr Gly Leu Trp Val Phe Ile Ser Thr Val Pro Cys Leu Thr Leu Ser
 100 105 110

Pro Cys Gly Glu Tyr Thr His Pro Thr Pro Thr Val Pro Cys Thr Ser
 115 120 125

Val Ala Ala Gln
 130

<210> 252
 <211> 30
 <212> PRT
 <213> Homo sapien

<400> 252

Met Gln Phe Arg Ile His Ala Ser Phe Ser Val Lys Trp Arg Ser Tyr

182

1 5 10 15

Ser Phe Asn Ser Glu Asn Ser Gln Leu Asn Lys Gln Pro Leu
 20 25 30

<210> 253
 <211> 49
 <212> PRT
 <213> Homo sapien

<400> 253

Met Arg Val Val Trp Gly Trp Arg Cys Gly Cys Val Gly Val Leu Val
 1 5 10 15

Leu Val Val Gly Gly Cys Val Glu Trp Ala Val Val Phe Gly Val Cys
 20 25 30

Val Gly Cys Val Val Trp Val Gly Arg Trp Trp Cys Asp Val Val Val
 35 40 45

Trp

<210> 254
 <211> 54
 <212> PRT
 <213> Homo sapien

<400> 254

Met Lys Lys Ser Val Ser Cys Cys Ser Ser Leu Trp Val Ser Leu Ser
 1 5 10 15

Lys Asp Glu Asn Ala Glu Val Gly Arg Gly Asp Ser Leu Leu Gly Thr
 20 25 30

Gly Arg Cys Gly Leu Pro Ile Thr Arg Leu Lys Leu Thr Ser Leu Pro
 35 40 45

Ser Ser Pro Thr Val Val
 50

<210> 255
 <211> 1088
 <212> PRT
 <213> Homo sapien

<400> 255

183

Asp Asp Ser Leu Ile Ser Ser Ala Thr Ala Ile Met Glu Ala Val Val
 1 5 10 15
 Arg Glu Trp Ile Leu Leu Glu Lys Gly Ser Ile Glu Ser Leu Arg Thr
 20 25 30
 Phe Leu Leu Thr Tyr Val Leu Gln Arg Pro Asn Leu Gln Lys Tyr Val
 35 40 45
 Arg Glu Gln Ile Leu Leu Ala Val Ala Val Ile Val Lys Arg Gly Ser
 50 55 60
 Leu Asp Lys Ser Ile Asp Cys Lys Ser Ile Phe His Glu Val Ser Gln
 65 70 75 80
 Leu Ile Ser Ser Gly Asn Pro Thr Val Gln Thr Leu Ala Cys Ser Ile
 85 90 95
 Leu Thr Ala Leu Leu Ser Glu Phe Ser Ser Ser Lys Thr Ser Asn
 100 105 110
 Ile Gly Leu Ser Met Glu Phe His Gly Asn Cys Lys Arg Val Phe Gln
 115 120 125
 Glu Glu Asp Leu Arg Gln Ile Phe Met Leu Thr Val Glu Val Leu Gln
 130 135 140
 Glu Phe Ser Arg Arg Glu Asn Leu Asn Ala Gln Met Ser Ser Val Phe
 145 150 155 160
 Gln Arg Tyr Leu Ala Leu Ala Asn Gln Val Leu Ser Trp Asn Phe Leu
 165 170 175
 Pro Pro Asn Leu Gly Arg His Tyr Ile Ala Met Phe Glu Ser Ser Gln
 180 185 190
 Asn Val Leu Leu Lys Pro Thr Glu Ser Leu Arg Glu Thr Leu Leu Asp
 195 200 205
 Ser Arg Val Met Glu Leu Phe Phe Thr Val His Arg Lys Ile Arg Glu
 210 215 220
 His Ser Asp Met Ala Gln Asp Ser Leu Gln Cys Leu Ala Gln Leu Ala
 225 230 235 240

184

Ser Leu His Gly Pro Ile Phe Pro Asp Glu Gly Ser Gln Val Asp Tyr
 245 250 255

Leu Ala His Phe Ile Glu Gly Leu Leu Asn Thr Ile Asn Gly Ile Glu
 260 265 270

Ile Glu Asp Ser Glu Ala Val Gly Ile Ser Ser Ile Ile Ser Asn Leu
 275 280 285

Ile Thr Val Phe Pro Arg Asn Val Leu Thr Ala Ile Pro Ser Glu Leu
 290 295 300

Phe Ser Ser Phe Val Asn Cys Leu Thr His Leu Thr Cys Ser Phe Gly
 305 310 315 320

Arg Ser Ala Ala Leu Glu Glu Val Leu Asp Lys Asp Asp Met Val Tyr
 325 330 335

Met Glu Ala Tyr Asp Lys Leu Leu Glu Ser Trp Leu Thr Leu Val Gln
 340 345 350

Asp Asp Lys His Phe His Lys Gly Phe Phe Thr Gln His Ala Val Gln
 355 360 365

Val Phe Asn Ser Tyr Ile Gln Cys His Leu Ala Ala Pro Asp Gly Thr
 370 375 380

Arg Asn Leu Thr Ala Asn Gly Val Ala Ser Arg Glu Glu Glu Glu Ile
 385 390 395 400

Ser Glu Leu Gln Glu Asp Asp Arg Asp Gln Phe Ser Asp Gln Leu Ala
 405 410 415

Ser Val Gly Met Leu Gly Arg Ile Ala Ala Glu His Cys Ile Pro Leu
 420 425 430

Leu Thr Ser Leu Leu Glu Glu Arg Val Thr Arg Leu His Gly Gln Leu
 435 440 445

Gln Arg His Gln Gln Gln Leu Leu Ala Ser Pro Gly Ser Ser Thr Val
 450 455 460

Asp Asn Lys Met Leu Asp Asp Leu Tyr Glu Asp Ile His Trp Leu Ile

186

Lys Gln Gln Tyr Trp Thr Glu Val Leu Gln Pro Leu Gln Gln Arg Phe
 705 710 715 720

Leu Arg Val Ile Asn Gln Glu Asn Phe Gln Gln Met Cys Gln Gln Glu
 725 730 735

Glu Val Lys Gln Glu Ile Thr Ala Thr Leu Glu Ala Leu Cys Gly Ile
 740 745 750

Ala Glu Ala Thr Gln Ile Asp Asn Val Ala Ile Leu Phe Asn Phe Leu
 755 760 765

Met Asp Phe Leu Thr Asn Cys Ile Gly Leu Met Glu Val Tyr Lys Asn
 770 775 780

Thr Pro Glu Thr Val Asn Leu Ile Ile Glu Val Phe Val Glu Val Ala
 785 790 795 800

His Lys Gln Ile Cys Tyr Leu Gly Glu Ser Lys Ala Met Asn Leu Tyr
 805 810 815

Glu Ala Cys Leu Thr Leu Leu Gln Val Tyr Ser Lys Asn Asn Leu Gly
 820 825 830

Arg Gln Arg Ile Asp Val Thr Ala Glu Glu Glu Gln Tyr Gln Asp Leu
 835 840 845

Leu Leu Ile Met Glu Leu Leu Thr Asn Leu Leu Ser Lys Glu Phe Ile
 850 855 860

Asp Phe Ser Asp Thr Asp Glu Val Phe Arg Gly His Glu Pro Gly Gln
 865 870 875 880

Ala Ala Asn Arg Ser Val Ser Ala Ala Asp Val Val Leu Tyr Gly Val
 885 890 895

Asn Leu Ile Leu Pro Leu Met Ser Gln Asp Leu Leu Lys Phe Pro Thr
 900 905 910

Leu Cys Asn Gln Tyr Tyr Lys Leu Ile Thr Phe Ile Cys Glu Ile Phe
 915 920 925

Pro Glu Lys Ile Pro Gln Leu Pro Glu Asp Leu Phe Lys Ser Leu Met
 930 935 940

187

Tyr Ser Leu Glu Leu Gly Met Thr Ser Met Ser Ser Glu Val Cys Gln
 945 950 955 960

Leu Cys Leu Glu Ala Leu Thr Pro Leu Ala Glu Gln Cys Ala Lys Ala
 965 970 975

Gln Glu Thr Asp Ser Pro Leu Phe Leu Ala Thr Arg His Phe Leu Lys
 980 985 990

Leu Val Phe Asp Met Leu Val Leu Gln Lys His Asn Thr Glu Met Thr
 995 1000 1005

Thr Ala Ala Gly Glu Ala Phe Tyr Thr Leu Val Cys Leu His Gln
 1010 1015 1020

Ala Glu Tyr Ser Glu Leu Val Glu Thr Leu Leu Ser Ser Gln Gln
 1025 1030 1035

Asp Pro Val Ile Tyr Gln Arg Leu Ala Asp Ala Phe Asn Lys Leu
 1040 1045 1050

Thr Ala Ser Ser Thr Pro Pro Thr Leu Asp Arg Lys Gln Lys Met
 1055 1060 1065

Ala Phe Leu Lys Ser Leu Glu Glu Phe Met Ala Asn Val Gly Gly
 1070 1075 1080

Leu Leu Cys Val Lys
 1085

<210> 256
 <211> 78
 <212> PRT
 <213> Homo sapien

<400> 256

Met Val Leu Met Thr Ser Ser Gly Gln Pro Ser Cys Pro Gly Ile Met
 1 5 10 15

Ala Cys Gln His Ser Leu Cys Pro Pro Asn Leu Arg Pro Arg Met Arg
 20 25 30

Ser Cys Gln His Asn Ile His Pro Phe Glu Gln Met Glu Ser Gly Thr
 35 40 45

188

Leu Thr Gln Pro Ser Val Leu Asn Asn Thr Ala Ile Ile Ala Thr Trp
 50 55 60

Leu Ser Arg Gln Cys Lys Pro Ser Glu Ser Ala Glu Leu Phe
 65 70 75

<210> 257
 <211> 595
 <212> PRT
 <213> Homo sapien

<400> 257

Val Gln Lys Thr Asn Gln Cys Leu Gln Gly Gln Ser Leu Lys Thr Ser
 1 5 10 15

Leu Thr Leu Lys Val Asp Arg Gly Ser Glu Glu Thr Tyr Arg Pro Glu
 20 25 30

Phe Pro Ser Thr Lys Gly Leu Val Arg Ser Leu Ala Glu Gln Phe Gln
 35 40 45

Arg Met Gln Gly Val Ser Met Arg Asp Ser Thr Gly Phe Lys Asp Arg
 50 55 60

Ser Leu Ser Gly Ser Leu Arg Lys Asn Ser Ser Pro Ser Asp Ser Lys
 65 70 75 80

Pro Pro Phe Ser Gln Gly Gln Glu Lys Gly His Trp Pro Trp Ala Lys
 85 90 95

Gln Gln Ser Ser Leu Glu Gly Gly Asp Arg Pro Leu Ser Trp Glu Glu
 100 105 110

Ser Thr Glu His Ser Ser Leu Ala Leu Asn Ser Gly Leu Pro Asn Gly
 115 120 125

Glu Thr Ser Ser Gly Gly Gln Pro Arg Leu Ala Glu Pro Asp Ile Tyr
 130 135 140

Gln Glu Lys Leu Ser Gln Val Arg Asp Val Arg Ser Lys Asp Leu Gly
 145 150 155 160

Ser Ser Thr Asp Leu Gly Thr Ser Leu Pro Leu Asp Ser Trp Val Asn
 165 170 175

189

Ile Thr Arg Phe Cys Asp Ser Gln Leu Lys His Gly Ala Pro Arg Pro
 180 185 190

Gly Met Lys Ser Ser Pro His Asp Ser His Thr Cys Val Thr Tyr Pro
 195 200 205

Glu Arg Asn His Ile Leu Leu His Pro His Trp Asn Gln Asp Thr Glu
 210 215 220

Gln Glu Thr Ser Glu Leu Glu Ser Leu Tyr Gln Ala Ser Leu Gln Ala
 225 230 235 240

Ser Gln Ala Gly Cys Ser Gly Trp Gly Gln Gln Asp Thr Ala Trp His
 245 250 255

Pro Leu Ser Gln Thr Gly Ser Ala Asp Gly Met Gly Arg Arg Leu His
 260 265 270

Ser Ala His Asp Pro Gly Leu Ser Lys Thr Ser Thr Ala Glu Met Glu
 275 280 285

His Gly Leu His Glu Ala Arg Thr Val Arg Thr Ser Gln Ala Thr Pro
 290 295 300

Cys Arg Gly Leu Ser Arg Glu Cys Gly Glu Asp Glu Gln Tyr Ser Ala
 305 310 315 320

Glu Asn Leu Arg Arg Ile Ser Arg Ser Leu Ser Gly Thr Val Val Ser
 325 330 335

Glu Arg Glu Glu Ala Pro Val Ser Ser His Ser Phe Asp Ser Ser Asn
 340 345 350

Val Arg Lys Pro Leu Glu Thr Gly His Arg Cys Ser Ser Ser Ser Ser
 355 360 365

Leu Pro Val Ile His Asp Pro Ser Val Phe Leu Leu Gly Pro Gln Leu
 370 375 380

Tyr Leu Pro Gln Pro Gln Phe Leu Ser Pro Asp Val Leu Met Pro Thr
 385 390 395 400

Met Ala Gly Glu Pro Asn Arg Leu Pro Gly Thr Ser Arg Ser Val Gln
 405 410 415

190

Gln Phe Leu Ala Met Cys Asp Arg Gly Glu Thr Ser Gln Gly Ala Lys
 420 425 430

Tyr Thr Gly Arg Thr Leu Asn Tyr Gln Ser Leu Pro His Arg Ser Arg
 435 440 445

Thr Asp Asn Ser Trp Ala Pro Trp Ser Glu Thr Asn Gln His Ile Gly
 450 455 460

Thr Arg Phe Leu Thr Thr Pro Gly Cys Asn Pro Gln Leu Thr Tyr Thr
 465 470 475 480

Ala Thr Leu Pro Glu Arg Ser Lys Gly Leu Gln Val Pro His Thr Gln
 485 490 495

Ser Trp Ser Asp Leu Phe His Ser Pro Ser His Pro Pro Ile Val His
 500 505 510

Pro Val Tyr Pro Pro Ser Ser Ser Leu His Val Pro Leu Arg Ser Ala
 515 520 525

Trp Asn Ser Asp Pro Val Pro Gly Ser Arg Thr Pro Gly Pro Arg Arg
 530 535 540

Val Asp Met Pro Pro Asp Asp Asp Trp Arg Gln Ser Ser Tyr Ala Ser
 545 550 555 560

His Ser Gly His Arg Arg Thr Val Gly Glu Gly Phe Leu Phe Val Leu
 565 570 575

Ser Asp Ala Pro Arg Arg Glu Gln Ile Arg Ala Arg Val Leu Gln His
 580 585 590

Ser Gln Trp
 595

<210> 258
 <211> 55
 <212> PRT
 <213> Homo sapien

<400> 258

Met Thr Val Met Ile Leu Leu Phe Lys Lys Asn Pro Asn Cys Tyr Phe
 1 5 10 15

191

Asp Leu Tyr Asp Leu Thr Leu Asn His Gly Ser Ile Thr Met Met Phe
 20 25 30

Lys Thr Leu Ile Asp Ser Thr Cys Phe Lys Asn Ser Gln Ile Pro Ser
 35 40 45

Ala Phe Ile Ile Arg Asp Arg
 50 55

<210> 259
 <211> 43
 <212> PRT
 <213> Homo sapien

<400> 259

Met Met Leu Thr Met Glu Phe Lys Asn Lys Gln Gln His Phe Val Val
 1 5 10 15

Ser Thr Gly Val Gly Val Glu Glu Leu Gln Arg His His Gly Asn Lys
 20 25 30

Ser Leu Pro Arg Ile Ser Gly Pro Arg Asn Leu
 35 40

<210> 260
 <211> 75
 <212> PRT
 <213> Homo sapien

<400> 260

Met Ala Tyr Arg Met Lys Arg Gly Thr Arg Asn Pro Cys Gly Arg Gly
 1 5 10 15

Leu Asp Leu Lys Gln Cys Pro Leu Trp Leu Leu Leu Pro Trp Leu Thr
 20 25 30

Gly Phe Leu Asp His Val His Phe Thr Gly Pro Trp Asp Leu His Leu
 35 40 45

Leu Ala Ser Pro Ala Gly Leu Ile Pro Ala Arg Ala Pro Ser Phe Leu
 50 55 60

Leu Met Val Phe Arg Trp Pro Asp His Gly Lys
 65 70 75

193

Glu Lys Leu Met Cys Ser Ser Ser Arg Ser
 210 215

<210> 262
 <211> 104
 <212> PRT
 <213> Homo sapien

<400> 262

Met Pro Ser Phe Phe Cys Phe Ser Ile Ser Leu Ile Arg Asp Trp Lys
 1 5 10 15

Val Ser Ile Arg Ser Asn Thr Asp Phe Ile Val Ile Gly Thr Asn Cys
 20 25 30

Ser Pro Thr Thr Pro Tyr Ser Ala Ser Ser Ile Thr Leu Leu Cys Glu
 35 40 45

Ile Leu Arg Asn Gly Leu Pro Leu Gln Gly Leu Asn Leu Pro Tyr Leu
 50 55 60

Arg Phe Glu Ser Ser Val Leu Phe Cys Ile Cys Phe Lys Tyr Leu Gly
 65 70 75 80

Ser Val Thr His Ala Asn Met Thr Cys Pro Val Gln Ala Thr Leu Gly
 85 90 95

Ile His Ile Ser His Val Ser Ser
 100

<210> 263
 <211> 260
 <212> PRT
 <213> Homo sapien

<400> 263

Glu Lys Lys Lys Lys Met Lys Asn Glu Asn Ala Asp Lys Leu Leu Lys
 1 5 10 15

Ser Glu Lys Gln Met Lys Lys Ser Glu Lys Lys Ser Lys Gln Glu Lys
 20 25 30

Glu Lys Ser Lys Lys Lys Lys Gly Gly Lys Thr Glu Gln Asp Gly Tyr
 35 40 45

194

Gln Lys Pro Thr Asn Lys His Phe Thr Gln Ser Pro Lys Lys Ser Val
 50 55 60

Ala Asp Leu Leu Gly Ser Phe Glu Gly Lys Arg Arg Leu Leu Leu Ile
 65 70 75 80

Thr Ala Pro Lys Ala Glu Asn Asn Met Tyr Val Gln Gln Arg Asp Glu
 85 90 95

Tyr Leu Glu Ser Phe Cys Lys Met Ala Thr Arg Lys Ile Ser Val Ile
 100 105 110

Thr Ile Phe Gly Pro Val Asn Asn Ser Thr Met Lys Ile Asp His Phe
 115 120 125

Gln Leu Asp Asn Glu Lys Pro Met Arg Val Val Asp Asp Glu Asp Leu
 130 135 140

Val Asp Gln Arg Leu Ile Ser Glu Leu Arg Lys Glu Tyr Gly Met Thr
 145 150 155 160

Tyr Asn Asp Phe Phe Met Val Leu Thr Asp Val Asp Leu Arg Val Lys
 165 170 175

Gln Tyr Tyr Glu Val Pro Ile Thr Met Lys Ser Val Phe Asp Leu Ile
 180 185 190

Asp Thr Phe Gln Ser Arg Ile Lys Asp Met Glu Lys Gln Lys Lys Glu
 195 200 205

Gly Ile Val Cys Lys Glu Asp Lys Lys Gln Ser Leu Glu Asn Phe Leu
 210 215 220

Ser Arg Phe Arg Trp Arg Arg Arg Leu Leu Val Ile Ser Ala Pro Asn
 225 230 235 240

Asp Glu Asp Trp Ala Tyr Ser Gln Gln Leu Ser Ala Leu Ser Gly Gln
 245 250 255

Ala Cys Thr Leu
 260

<210> 264
 <211> 62
 <212> PRT

195

<213> Homo sapien

<400> 264

Met Ser Gly Phe Ile Tyr Val Leu Glu Lys Asp His Leu Lys Lys Ile
 1 5 10 15

Asn Thr Phe Ser Thr Thr Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys
 20 25 30

Arg Arg Gly Gly Glu Pro Gly Ala Gln Ser Gly Pro Arg Gly Ala Asn
 35 40 45

Trp Val Leu Pro Ala His Ile Pro Pro Lys Tyr Trp His Thr
 50 55 60

<210> 265

<211> 89

<212> PRT

<213> Homo sapien

<400> 265

Met Leu Gln Leu Asn Thr Arg Phe Tyr Phe Leu Ser Asn Cys Gly Phe
 1 5 10 15

Val Phe Ile Tyr His Pro Leu Phe Ile Pro Phe Leu Thr His Thr Leu
 20 25 30

Cys Arg Ala Ser Gly Ile Tyr Tyr Ser Thr Val Cys Leu Cys Lys Arg
 35 40 45

Leu Ser Val Leu Ala Ser Thr Tyr Glu Arg Met His Ala Lys Phe Cys
 50 55 60

Leu Ser Met Pro Gly Leu Ile Ser Leu Lys Gln Asn Asp Leu Arg Val
 65 70 75 80

Pro Ser Met Leu Phe Ile Leu Pro Asn
 85

<210> 266

<211> 38

<212> PRT

<213> Homo sapien

<400> 266

Met Thr Ser Arg Trp Leu Asn Phe Ser Cys Leu Trp Cys Phe Gly Pro

196

1 5 10 15
 Asn Ser Thr Gly Gln His His Asp His Met Glu Thr Tyr Phe Trp Lys
 20 25 30

Gln Asn Phe Asn Phe Ile
 35

<210> 267
 <211> 111
 <212> PRT
 <213> Homo sapien
 <400> 267

Asn Asp Leu Asp Arg Tyr Asn Pro Leu Ser Ser Gln Arg Leu Val Arg
 1 5 10 15

Asn Ala Leu Ala His Val Gly Ala Lys Glu Arg Glu Leu Ser Trp Ala
 20 25 30

His Ser Glu Ser Phe Ala Ala Leu Cys Arg Tyr Gly Lys Arg Glu Phe
 35 40 45

Lys Ile Gly Gly Glu Leu Arg Ile Gly Lys Gln Pro Tyr Arg Leu Gln
 50 55 60

Ile Gln Leu Ser Ala Gln Arg Ser His Thr Leu Glu Phe Gln Ser Leu
 65 70 75 80

Glu Asp Leu Ile Met Gly Glu Ala Thr Gln Arg Pro Arg Ser Gly Ala
 85 90 95

Arg Pro Val Leu Gln Glu Leu Ala Thr His Leu His Pro Ala Glu
 100 105 110

<210> 268
 <211> 60
 <212> PRT
 <213> Homo sapien
 <400> 268

Met Val Asn Thr Val Leu Leu Ser Leu Lys Ile Ser Leu Phe Cys Pro
 1 5 10 15

His Gln Leu Phe Tyr Cys Ser Val Leu Arg Lys Pro Asn Ser Cys Val
 20 25 30

Phe Phe Pro Ser Leu Leu Ile Leu Ser Cys Val Pro Ser Gly Lys Cys
 35 40 45

His Tyr Phe Leu Asp Ile Leu Asn Leu Leu Phe Leu
 50 55 60

<210> 269
 <211> 72
 <212> PRT
 <213> Homo sapien
 <400> 269

Met Cys Leu Cys Ile Leu Val Ser Lys Leu Arg Thr Ser Asp Glu Leu
 1 5 10 15

Pro Val Val Pro Ser Tyr Cys Arg Arg Leu Glu Val Arg Gly Ile Ser
 20 25 30

Ala Ser Thr Arg Glu Ala Glu Val Ala Ser Glu Pro Thr Ile Met Thr
 35 40 45

Ala Cys Thr Pro Ser Leu Ala Thr Val Arg Glu Leu Leu Ser Gln Ile
 50 55 60

Lys Arg Lys Gln Ser Leu Leu Ser
 65 70

<210> 270
 <211> 152
 <212> PRT
 <213> Homo sapien
 <400> 270

Gly Ser Leu Gly Gly Glu Pro Gly Val Ser Cys Leu Lys Met His Ser
 1 5 10 15

Asp Ala Ala Ala Val Asn Phe Gln Leu Asn Ser His Leu Ser Thr Leu
 20 25 30

Ala Asn Ile His Lys Ile Tyr His Thr Leu Asn Lys Leu Asn Leu Thr
 35 40 45

Glu Asp Ile Gly Gln Asp Asp His Gln Thr Gly Ser Leu Arg Ser Cys
 50 55 60

198

Ser Ser Ser Asp Cys Phe Asn Lys Val Met Pro Pro Arg Lys Lys Arg
65 70 75 80

Arg Pro Ala Ser Gly Asp Asp Leu Ser Ala Lys Lys Ser Arg His Asp
85 90 95

Ser Met Tyr Arg Lys Tyr Asp Ser Thr Arg Ile Lys Thr Glu Glu Glu
100 105 110

Ala Phe Ser Ser Lys Arg Cys Leu Glu Trp Phe Tyr Glu Tyr Ala Gly
115 120 125

Thr Asp Asp Val Val Gly Pro Glu Gly Met Glu Lys Phe Cys Glu Asp
130 135 140

Ile Gly Val Glu Pro Glu Asn Val
145 150

<210> 271
<211> 52
<212> PRT
<213> Homo sapien

<400> 271

Met Glu Pro His Ile Met Lys Phe Asn Ser His Val Lys Thr Phe Cys
1 5 10 15

Ile Val Gly Cys Gln Lys Tyr Leu Pro Lys Leu Ser Phe Asp Leu Ser
20 25 30

Glu Trp Gly Trp Leu Leu Pro Ile Leu Gln Phe Val Ser Gln Ala Trp
35 40 45

Arg Asn Gln Ala
50

<210> 272
<211> 449
<212> PRT
<213> Homo sapien

<400> 272

Met Val Met Glu Lys Pro Ser Pro Leu Leu Val Gly Arg Glu Phe Val
1 5 10 15

199

Arg Gln Tyr Tyr Thr Leu Leu Asn Lys Ala Pro Glu Tyr Leu His Arg
 20 25 30

Phe Tyr Gly Arg Asn Ser Ser Tyr Val His Gly Gly Val Asp Ala Ser
 35 40 45

Gly Lys Pro Gln Glu Ala Val Tyr Gly Gln Asn Asp Ile His His Lys
 50 55 60

Val Leu Ser Leu Asn Phe Ser Glu Cys His Thr Lys Ile Arg His Val
 65 70 75 80

Asp Ala His Ala Thr Leu Ser Asp Gly Val Val Val Gln Val Met Gly
 85 90 95

Leu Leu Ser Asn Ser Gly Gln Pro Glu Arg Lys Phe Met Gln Thr Phe
 100 105 110

Val Leu Ala Pro Glu Gly Ser Val Pro Asn Lys Phe Tyr Val His Asn
 115 120 125

Asp Met Phe Arg Tyr Glu Asp Glu Val Phe Gly Asp Ser Glu Pro Glu
 130 135 140

Leu Asp Glu Glu Ser Glu Asp Glu Val Glu Glu Glu Gln Glu Glu Arg
 145 150 155 160

Gln Pro Ser Pro Glu Pro Val Gln Glu Asn Ala Asn Ser Gly Tyr Tyr
 165 170 175

Glu Ala His Pro Val Thr Asn Gly Ile Glu Glu Pro Leu Glu Glu Ser
 180 185 190

Ser His Glu Pro Glu Pro Glu Pro Glu Ser Glu Thr Lys Thr Glu Glu
 195 200 205

Leu Lys Pro Gln Val Glu Glu Lys Asn Leu Glu Glu Leu Glu Glu Lys
 210 215 220

Ser Thr Thr Pro Pro Pro Ala Glu Pro Val Ser Leu Pro Gln Glu Pro
 225 230 235 240

Pro Lys Pro Arg Val Glu Ala Lys Pro Glu Val Gln Ser Gln Pro Pro
 245 250 255

200

Arg Val Arg Glu Gln Arg Pro Arg Glu Arg Pro Gly Phe Pro Pro Arg
 260 265 270

Gly Pro Arg Pro Gly Arg Gly Asp Met Glu Gln Asn Asp Ser Asp Asn
 275 280 285

Arg Arg Ile Ile Arg Tyr Pro Asp Ser His Gln Leu Phe Val Gly Asn
 290 295 300

Leu Pro His Asp Ile Asp Glu Asn Glu Leu Lys Glu Phe Phe Met Ser
 305 310 315 320

Phe Gly Asn Val Val Glu Leu Arg Ile Asn Thr Lys Gly Val Gly Gly
 325 330 335

Lys Leu Pro Asn Phe Gly Phe Val Val Phe Asp Asp Ser Glu Pro Val
 340 345 350

Gln Arg Ile Leu Ile Ala Lys Pro Ile Met Phe Arg Gly Glu Val Arg
 355 360 365

Leu Asn Val Glu Glu Lys Lys Thr Arg Ala Ala Arg Glu Arg Glu Thr
 370 375 380

Arg Gly Gly Gly Asp Asp Arg Arg Asp Ile Arg Arg Asn Asp Arg Gly
 385 390 395 400

Pro Gly Gly Pro Arg Gly Ile Val Gly Gly Gly Met Met Arg Asp Arg
 405 410 415

Asp Gly Arg Gly Pro Pro Pro Arg Gly Gly Met Ala Gln Lys Leu Gly
 420 425 430

Ser Gly Arg Gly Thr Gly Gln Met Glu Gly Arg Phe Thr Gly Gln Arg
 435 440 445

Arg

<210> 273

<211> 63

<212> PRT

<213> Homo sapien

<400> 273

201

Met Cys Cys Asp Val Ser Glu Arg Ala Glu Phe Arg Leu Val Ser Ala
 1 5 10 15

Arg Cys Ser Phe Ser His Pro Arg Thr Val Ala Arg Leu Leu Leu Arg
 20 25 30

His Pro Gly Gln Leu Pro Leu Pro Phe Gln Trp Gly Leu Thr Trp Leu
 35 40 45

Pro Ser Leu Ala Ala Asn Arg Arg Ala Pro Gln His Ser Arg Ser
 50 55 60

<210> 274
 <211> 60
 <212> PRT
 <213> Homo sapien

<400> 274

Met Asp Pro Gly Arg Tyr Cys Leu Val Leu Gln Glu Leu Met Gln Phe
 1 5 10 15

His Ser Glu Ala Cys Lys Ile Leu Asn Phe Arg Asp Asn Arg Pro Asp
 20 25 30

Thr Phe Leu Ile Ser Phe Tyr Ser Leu Met Ser Asn Asn Thr Ile Phe
 35 40 45

Lys Asn Met Val Leu Ile Cys Leu Ala Ser Asn Leu
 50 55 60

<210> 275
 <211> 111
 <212> PRT
 <213> Homo sapien

<400> 275

Lys Leu Ile Val Tyr Pro Pro Pro Pro Ala Lys Gly Gly Ile Ser Val
 1 5 10 15

Thr Asn Glu Asp Leu His Cys Leu Asn Glu Gly Glu Phe Leu Asn Asp
 20 25 30

Val Ile Ile Asp Phe Tyr Leu Lys Tyr Leu Val Leu Glu Lys Leu Lys
 35 40 45

202

Lys Glu Asp Ala Asp Arg Ile His Ile Phe Ser Ser Phe Phe Tyr Lys
50 55 60

Arg Leu Asn Gln Arg Glu Arg Arg Asn His Glu Thr Thr Asn Leu Ser
65 70 75 80

Ile Gln Gln Lys Arg His Gly Arg Val Lys Thr Trp Thr Arg His Val
85 90 95

Asp Ile Phe Glu Lys Asp Phe Ile Phe Val Pro Leu Asn Glu Ala
100 105 110

<210> 276
<211> 97
<212> PRT
<213> Homo sapien

<400> 276

Met Ser Gln Asp Thr Ser Arg Ser Gln Glu Arg Ala Ala Gly Pro Gln
1 5 10 15

Arg Thr Arg Arg Arg Pro Arg Thr Trp Ser Gly Gly Val Glu Pro Thr
20 25 30

Ala Ala Ala Pro Trp Ala Ala Ala Met Ala His Thr Gly Arg His Gly
35 40 45

Ser Gly Ala Ala Ala Thr Ala Ser Ser Thr Arg Gly Asp Gly Ala Ala
50 55 60

Arg Arg Gly Ala Ala Arg Gly Thr Asp Ala Ala Glu Arg Arg Arg Ala
65 70 75 80

Ala Ser Arg Gly Ala Ala Glu Pro Lys Ala Thr Ala Ser Gly Gly Gly
85 90 95

Gly

<210> 277
<211> 76
<212> PRT
<213> Homo sapien

<400> 277

Met Gly Ser Cys Pro Leu Trp Val Arg Ser Ser Thr Cys Arg Val Glu

203

1 5 10 15
 Val Gly Tyr Val His Thr Phe Asn Asp Asn Leu His Ile Ser Ala Pro
 20 25 30
 Thr Gly Pro Lys Leu Phe Leu Gly Phe Lys Val Val Val Cys Leu Phe
 35 40 45
 Phe Ser Phe Phe Phe Phe Phe Phe Phe Gly Glu Val Glu Phe Gly
 50 55 60
 Ser Gly Trp Pro Arg Cys Gly Val Cys Lys Gly Arg
 65 70 75

<210> 278
 <211> 20
 <212> PRT
 <213> Homo sapien

<400> 278

Met Glu Asp Gln Ile Ile Leu Asn Tyr Ile Ser Ile Val Pro Gly Lys
 1 5 10 15

Thr Gln Val Leu
 20

<210> 279
 <211> 24
 <212> PRT
 <213> Homo sapien

<400> 279

Met Val His Leu Met His Ala Arg Ala Arg Ala Ser Cys Asp Gly Cys
 1 5 10 15

Val Val Ala Ala Glu Val His Val
 20

<210> 280
 <211> 101
 <212> PRT
 <213> Homo sapien

<400> 280

Leu Phe Phe Phe Lys Lys Phe Ile Leu Arg Trp Ser Leu Thr Leu Ser
 1 5 10 15

204

Leu Arg Leu Glu Cys Ser Asp Ser Ile Ser Ala His Cys Asn Leu Arg
 20 25 30

Leu Pro Gly Leu Ser Asn Phe Cys Ala Ser Ala Ser Gln Val Ser Glu
 35 40 45

Ile Thr Gly Val Cys His His Thr Gln Leu Phe Phe Ile Phe Tyr Phe
 50 55 60

Ala Ala Lys Met Gly Phe Arg His Val Gly Arg Thr Gly Leu Glu Leu
 65 70 75 80

Leu Ala Ser Ser Gly Pro Pro Thr Ser Ala Ser Gln Ser Ala Gly Ile
 85 90 95

Thr Gly Val Ser His
 100

<210> 281
 <211> 43
 <212> PRT
 <213> Homo sapien

<400> 281

Met Trp Gly His Gly Leu Asp Asp Gly Leu His Arg Ser Phe His Leu
 1 5 10 15

Cys Glu Ser Lys Ser Gly Gln Ser Ala Arg Thr Gln Ser Leu Thr Leu
 20 25 30

Gly Gln Leu Leu Arg Thr Asn Pro Gln His Leu
 35 40

<210> 282
 <211> 46
 <212> PRT
 <213> Homo sapien

<400> 282

Met Ala Gly Asn Ile His Pro Gly Thr Phe Gly Pro Gly Ser Pro His
 1 5 10 15

Leu Phe Phe Leu Cys Gly Val Val Ala Phe Phe Leu Phe Ile Val Ala
 20 25 30

205

Arg Glu Ala Lys Ile Tyr Ser Phe Ser Met Asn Pro Asn Met
 35 40 45

<210> 283
 <211> 70
 <212> PRT
 <213> Homo sapien

<400> 283

Met Pro Gly Ser His Leu Cys Met Phe Asn Thr Val Thr His Asp Val
 1 5 10 15

Ile Thr Glu Trp Arg Arg Trp Lys Gly Pro Cys Arg Ser Phe Ser Trp
 20 25 30

His Pro Asn Phe Thr Glu Gly Glu Leu Arg Pro Glu Leu Arg Asp Val
 35 40 45

Leu Arg Ile Pro Glu Ser His Ser Ser Val Arg Ser Val Ile His Lys
 50 55 60

Glu Val Ile Ile Lys Val
 65 70

<210> 284
 <211> 49
 <212> PRT
 <213> Homo sapien

<400> 284

Met Ser Ser Ser Leu Phe Ala Phe Leu Leu Thr Tyr Phe Val Val Phe
 1 5 10 15

Lys Asp Cys Ala Gly Asp Ile Leu Glu Gly Ile Asn Gly Leu His Ser
 20 25 30

Lys Arg Cys Gly Leu Ser Lys Leu Phe Ser Val Phe Ile Thr Glu Thr
 35 40 45

Asp

<210> 285
 <211> 1544
 <212> PRT
 <213> Homo sapien

206

<400> 285

Met Tyr Ala Ala Val Glu His Gly Pro Val Leu Cys Ser Asp Ser Asn
 1 5 10 15

Ile Leu Cys Leu Ser Trp Lys Gly Arg Val Pro Lys Ser Glu Lys Glu
 20 25 30

Lys Pro Val Cys Arg Arg Arg Tyr Tyr Glu Glu Gly Trp Leu Ala Thr
 35 40 45

Gly Asn Gly Arg Gly Val Val Gly Val Thr Phe Thr Ser Ser His Cys
 50 55 60

Arg Arg Asp Arg Ser Thr Pro Gln Arg Ile Asn Phe Asn Leu Arg Gly
 65 70 75 80

His Asn Ser Glu Val Val Leu Val Arg Trp Asn Glu Pro Tyr Gln Lys
 85 90 95

Leu Ala Thr Cys Asp Ala Asp Gly Gly Ile Phe Val Trp Ile Gln Tyr
 100 105 110

Glu Gly Arg Trp Ser Val Glu Leu Val Asn Asp Arg Gly Ala Gln Val
 115 120 125

Ser Asp Phe Thr Trp Ser His Asp Gly Thr Gln Ala Leu Ile Ser Tyr
 130 135 140

Arg Asp Gly Phe Val Leu Val Gly Ser Val Ser Gly Gln Arg His Trp
 145 150 155 160

Ser Ser Glu Ile Asn Leu Glu Ser Gln Ile Thr Cys Gly Ile Trp Thr
 165 170 175

Pro Asp Asp Gln Gln Val Leu Phe Gly Thr Ala Asp Gly Gln Val Ile
 180 185 190

Val Met Asp Cys His Gly Arg Met Leu Ala His Val Leu Leu His Glu
 195 200 205

Ser Asp Gly Val Leu Gly Met Ser Trp Asn Tyr Pro Ile Phe Leu Val
 210 215 220

Glu Asp Ser Ser Glu Ser Asp Thr Asp Ser Asp Asp Tyr Ala Pro Pro

207

225					230										235						240	
Gln	Asp	Gly	Pro	Ala	Ala	Tyr	Pro	Ile	Pro	Val	Gln	Asn	Ile	Lys	Pro							
				245					250					255								
Leu	Leu	Thr	Val	Ser	Phe	Thr	Ser	Gly	Asp	Ile	Ser	Leu	Met	Asn	Asn							
			260					265					270									
Tyr	Asp	Asp	Leu	Ser	Pro	Thr	Val	Ile	Arg	Ser	Gly	Leu	Lys	Glu	Val							
	275						280					285										
Val	Ala	Gln	Trp	Cys	Thr	Gln	Gly	Asp	Leu	Leu	Ala	Val	Ala	Gly	Met							
	290					295					300											
Glu	Arg	Gln	Thr	Gln	Leu	Gly	Glu	Leu	Pro	Asn	Gly	Pro	Leu	Leu	Lys							
305					310					315					320							
Ser	Ala	Met	Val	Lys	Phe	Tyr	Asn	Val	Arg	Gly	Glu	His	Ile	Phe	Thr							
				325					330					335								
Leu	Asp	Thr	Leu	Val	Gln	Arg	Pro	Ile	Ile	Ser	Ile	Cys	Trp	Gly	His							
			340					345					350									
Arg	Asp	Ser	Arg	Leu	Leu	Met	Ala	Ser	Gly	Pro	Ala	Leu	Tyr	Val	Val							
		355					360					365										
Arg	Val	Glu	His	Arg	Val	Ser	Ser	Leu	Gln	Leu	Leu	Cys	Gln	Gln	Ala							
	370					375					380											
Ile	Ala	Ser	Thr	Leu	Arg	Glu	Asp	Lys	Asp	Val	Ser	Lys	Leu	Thr	Leu							
385					390					395					400							
Pro	Pro	Arg	Leu	Cys	Ser	Tyr	Leu	Ser	Thr	Ala	Phe	Ile	Pro	Thr	Ile							
				405					410					415								
Lys	Pro	Pro	Ile	Pro	Asp	Pro	Asn	Asn	Met	Arg	Asp	Phe	Val	Ser	Tyr							
			420					425				430										
Pro	Ser	Ala	Gly	Asn	Glu	Arg	Leu	His	Cys	Thr	Met	Lys	Arg	Thr	Glu							
		435					440				445											
Asp	Asp	Pro	Glu	Val	Gly	Gly	Pro	Cys	Tyr	Thr	Leu	Tyr	Leu	Glu	Tyr							
	450					455					460											

208

Leu Gly Gly Leu Val Pro Ile Leu Lys Gly Arg Arg Ile Ser Lys Leu
465 470 475 480

Arg Pro Glu Phe Val Ile Met Asp Pro Arg Thr Asp Ser Lys Pro Asp
485 490 495

Glu Ile Tyr Gly Asn Ser Leu Ile Ser Thr Val Ile Asp Ser Cys Asn
500 505 510

Cys Ser Asp Ser Ser Asp Ile Glu Leu Ser Asp Asp Trp Ala Ala Lys
515 520 525

Lys Ser Pro Lys Ile Ser Arg Ala Ser Lys Ser Pro Lys Leu Pro Arg
530 535 540

Ile Ser Ile Glu Ala Arg Lys Ser Pro Lys Leu Pro Arg Ala Ala Gln
545 550 555 560

Glu Leu Ser Arg Ser Pro Arg Leu Pro Leu Arg Lys Pro Ser Val Gly
565 570 575

Ser Pro Ser Leu Thr Arg Arg Glu Phe Pro Phe Glu Asp Ile Thr Gln
580 585 590

His Asn Tyr Leu Ala Gln Val Thr Ser Asn Ile Trp Gly Thr Lys Phe
595 600 605

Lys Ile Val Gly Leu Ala Ala Phe Leu Pro Thr Asn Leu Gly Ala Val
610 615 620

Ile Tyr Lys Thr Ser Leu Leu His Leu Gln Pro Arg Gln Met Thr Ile
625 630 635 640

Tyr Leu Pro Glu Val Arg Lys Ile Ser Met Asp Tyr Ile Asn Leu Pro
645 650 655

Val Phe Asn Pro Asn Val Phe Ser Glu Asp Glu Asp Asp Leu Pro Val
660 665 670

Thr Gly Ala Ser Gly Val Pro Glu Asn Ser Pro Pro Cys Thr Val Asn
675 680 685

Ile Pro Ile Ala Pro Ile His Ser Ser Ala Gln Ala Met Ser Pro Thr
690 695 700

209

Gln Ser Ile Gly Leu Val Gln Ser Leu Leu Ala Asn Gln Asn Val Gln
705 710 715 720

Leu Asp Val Leu Thr Asn Gln Thr Thr Ala Val Gly Thr Ala Glu His
725 730 735

Ala Gly Asp Arg Cys His Pro Val Thr Gln Val Ser Asn Arg Tyr Ser
740 745 750

Asn Pro Gly Gln Val Ile Phe Gly Ser Val Glu Met Gly Arg Ile Ile
755 760 765

Gln Asn Pro Pro Pro Leu Ser Leu Pro Pro Pro Pro Gln Gly Pro Met
770 775 780

Gln Leu Ser Thr Val Gly His Gly Asp Arg Asp His Glu His Leu Gln
785 790 795 800

Lys Ser Ala Lys Ala Leu Arg Pro Thr Pro Gln Leu Ala Ala Glu Gly
805 810 815

Asp Ala Val Val Phe Ser Ala Pro Gln Glu Val Gln Val Thr Lys Ile
820 825 830

Asn Pro Pro Pro Pro Tyr Pro Gly Thr Ile Pro Ala Ala Pro Thr Thr
835 840 845

Ala Ala Pro Pro Pro Pro Leu Pro Pro Pro Gln Pro Pro Val Asp Val
850 855 860

Cys Leu Lys Lys Gly Asp Phe Ser Leu Tyr Pro Thr Ser Val His Tyr
865 870 875 880

Gln Thr Pro Leu Gly Tyr Glu Arg Ile Thr Thr Phe Asp Ser Ser Gly
885 890 895

Asn Val Glu Glu Val Cys Arg Pro Arg Thr Arg Met Leu Cys Ser Gln
900 905 910

Asn Thr Tyr Thr Leu Pro Gly Pro Gly Ser Ser Ala Thr Leu Arg Leu
915 920 925

Thr Ala Thr Glu Lys Lys Val Pro Gln Pro Cys Ser Ser Ala Thr Leu
930 935 940

210

Asn Arg Leu Thr Val Pro Arg Tyr Ser Ile Pro Thr Gly Asp Pro Pro
 945 950 955 960

Pro Tyr Pro Glu Ile Ala Ser Gln Leu Ala Gln Gly Arg Gly Ala Ala
 965 970 975

Gln Arg Ser Asp Asn Ser Leu Ile His Ala Thr Leu Arg Arg Asn Asn
 980 985 990

Arg Glu Ala Thr Leu Lys Met Ala Gln Leu Ala Asp Ser Pro Arg Ala
 995 1000 1005

Pro Leu Gln Pro Leu Ala Lys Ser Lys Gly Gly Pro Gly Gly Val
 1010 1015 1020

Val Thr Gln Leu Pro Ala Arg Pro Pro Pro Ala Leu Tyr Thr Cys
 1025 1030 1035

Ser Gln Cys Ser Gly Thr Gly Pro Ser Ser Gln Pro Gly Ala Ser
 1040 1045 1050

Leu Ala His Thr Ala Ser Ala Ser Pro Leu Ala Ser Gln Ser Ser
 1055 1060 1065

Tyr Ser Leu Leu Ser Pro Pro Asp Ser Ala Arg Asp Arg Thr Asp
 1070 1075 1080

Tyr Val Asn Ser Ala Phe Thr Glu Asp Glu Ala Leu Ser Gln His
 1085 1090 1095

Cys Gln Leu Glu Lys Pro Leu Arg His Pro Pro Leu Pro Glu Ala
 1100 1105 1110

Ala Val Thr Leu Lys Arg Pro Pro Pro Tyr Gln Trp Asp Pro Met
 1115 1120 1125

Leu Gly Glu Asp Val Trp Val Pro Gln Glu Arg Thr Ala Gln Thr
 1130 1135 1140

Ser Gly Pro Asn Pro Leu Lys Leu Ser Ser Leu Met Leu Ser Gln
 1145 1150 1155

Gly Gln His Leu Asp Val Ser Arg Leu Pro Phe Ile Ser Pro Lys

211

1160						1165						1170		
Ser	Pro	Ala	Ser	Pro	Thr	Ala	Thr	Phe	Gln	Thr	Gly	Tyr	Gly	Met
1175						1180					1185			
Gly	Val	Pro	Tyr	Pro	Gly	Ser	Tyr	Asn	Asn	Pro	Pro	Leu	Pro	Gly
1190						1195					1200			
Val	Gln	Ala	Pro	Cys	Ser	Pro	Lys	Asp	Ala	Leu	Ser	Pro	Thr	Gln
1205						1210					1215			
Phe	Ala	Gln	Gln	Glu	Pro	Ala	Val	Val	Leu	Gln	Pro	Leu	Tyr	Pro
1220						1225					1230			
Pro	Ser	Leu	Ser	Tyr	Cys	Thr	Leu	Pro	Pro	Met	Tyr	Pro	Gly	Ser
1235						1240					1245			
Ser	Thr	Cys	Ser	Ser	Leu	Gln	Leu	Pro	Pro	Val	Ala	Leu	His	Pro
1250						1255					1260			
Trp	Ser	Ser	Tyr	Ser	Ala	Cys	Pro	Pro	Met	Gln	Asn	Pro	Gln	Gly
1265						1270					1275			
Thr	Leu	Pro	Pro	Lys	Pro	His	Leu	Val	Val	Glu	Lys	Pro	Leu	Val
1280						1285					1290			
Ser	Pro	Pro	Pro	Ala	Asp	Leu	Gln	Ser	His	Leu	Gly	Thr	Glu	Val
1295						1300					1305			
Met	Val	Glu	Thr	Ala	Asp	Asn	Phe	Gln	Glu	Val	Leu	Ser	Leu	Thr
1310						1315					1320			
Glu	Ser	Pro	Val	Pro	Gln	Arg	Thr	Glu	Lys	Phe	Gly	Lys	Lys	Asn
1325						1330					1335			
Arg	Lys	Arg	Leu	Asp	Ser	Arg	Ala	Glu	Glu	Gly	Ser	Val	Gln	Ala
1340						1345					1350			
Ile	Thr	Glu	Gly	Lys	Val	Lys	Lys	Glu	Ala	Arg	Thr	Leu	Ser	Asp
1355						1360					1365			
Phe	Asn	Ser	Leu	Ile	Ser	Ser	Pro	His	Leu	Gly	Arg	Glu	Lys	Lys
1370						1375					1380			

212

Lys Val Lys Ser Gln Lys Asp Gln Leu Lys Ser Lys Lys Leu Asn
 1385 1390 1395

Lys Thr Asn Glu Phe Gln Asp Ser Ser Glu Ser Glu Pro Glu Leu
 1400 1405 1410

Phe Ile Ser Gly Asp Glu Leu Met Asn Gln Ser Gln Gly Ser Arg
 1415 1420 1425

Lys Gly Trp Lys Ser Lys Arg Ser Pro Arg Ala Ala Gly Glu Leu
 1430 1435 1440

Glu Glu Ala Lys Cys Arg Arg Ala Ser Glu Lys Glu Asp Gly Arg
 1445 1450 1455

Leu Gly Ser Gln Gly Phe Val Tyr Val Met Ala Asn Lys Gln Pro
 1460 1465 1470

Leu Trp Asn Glu Ala Thr Gln Val Tyr Gln Leu Asp Phe Gly Gly
 1475 1480 1485

Arg Val Thr Gln Glu Ser Ala Lys Asn Phe Gln Ile Glu Leu Glu
 1490 1495 1500

Gly Arg Gln Val Met Gln Phe Gly Arg Ile Asp Gly Ser Ala Tyr
 1505 1510 1515

Ile Leu Asp Phe Gln Tyr Pro Phe Ser Ala Val Gln Ala Phe Ala
 1520 1525 1530

Val Ala Leu Ala Asn Val Thr Gln Arg Leu Lys
 1535 1540

<210> 286
 <211> 56
 <212> PRT
 <213> Homo sapien

<400> 286

Met Gly Asn Gly Ala Thr Gln Lys Gln Leu Pro Asn Leu Arg Asn Asn
 1 5 10 15

Ser Phe Val Val Tyr Phe Leu Val Leu Val Gly Ala Leu Tyr Arg Asp
 20 25 30

213

Thr Ala Ile Phe Leu Ala Gln Met Ser Leu Leu Glu Ser Thr Val Val
 35 40 45

Ile Leu Leu Val Arg Leu Arg Thr
 50 55

<210> 287
 <211> 77
 <212> PRT
 <213> Homo sapien

<400> 287

Met Leu Leu Ala Val Arg Thr Thr Val Ile Cys Leu Gln Ser Cys Cys
 1 5 10 15

Cys Arg Ile Gln Arg Thr Ala Thr Ile Thr Leu Asn Cys Phe Ala Leu
 20 25 30

Ser Ser Ile Phe Asp Tyr Tyr Ile Ser His Asn Ile Thr Ile Ser His
 35 40 45

Ser Ser Asn Tyr Ser Ala Gln Ile His Glu His Val Pro Ala Arg Ala
 50 55 60

Ala Ala Arg Ser Ile Thr Trp Arg Arg Ser Ala Cys Ile
 65 70 75

<210> 288
 <211> 45
 <212> PRT
 <213> Homo sapien

<400> 288

Met Tyr Leu Gly Gln Leu Gly Asn His Arg Leu Lys Lys Leu Thr Leu
 1 5 10 15

Val Ile Thr Arg Val Val Ser Asp Tyr Lys Gln His Ile Ile Asn Pro
 20 25 30

Thr Ala Leu Ile Leu Ala Gln Arg Gln Asn Trp Thr Phe
 35 40 45

<210> 289
 <211> 44
 <212> PRT
 <213> Homo sapien

214

<400> 289

Met Lys Ala Leu Leu Cys Phe Leu Phe Tyr Ser Asp His Gln Thr Asp
 1 5 10 15

Leu Ala Thr Leu Ile Val Lys Asn Glu Pro His Ser Ser Pro Gly Leu
 20 25 30

Gly Leu Trp Arg Glu Met Asn Phe Leu Leu Glu Met
 35 40

<210> 290

<211> 50

<212> PRT

<213> Homo sapien

<400> 290

Met Phe Arg Thr Ser Ser Tyr Arg Leu Leu Ile Tyr Lys Val Pro Val
 1 5 10 15

Ala Val Thr Pro Thr Arg Lys Thr Trp Asn Cys Lys Gln Ala Gly Val
 20 25 30

Thr Ser Val Thr Ser Asp Thr Val Gln Pro Glu Val Arg Phe Leu Phe
 35 40 45

Trp Gly
 50

<210> 291

<211> 44

<212> PRT

<213> Homo sapien

<400> 291

Met Ser Gln Trp Pro Val Ala Ser Lys Leu Val Gly Lys Glu Lys Thr
 1 5 10 15

Phe Leu Phe Lys Gln Arg Lys Gly Phe Gly Glu Lys Thr Gly Ser Gly
 20 25 30

Ser Gly Glu Val Phe Val Met Leu Gly Asp Arg Leu
 35 40

<210> 292

<211> 61

<212> PRT

215

<213> Homo sapien

<400> 292

Met Val His Tyr Arg Lys Glu Lys Lys Thr Ser Val Ser Glu Trp Gln
 1 5 10 15

Ile Leu Ile Ile Cys Ser Ser His Leu Phe Ser Ser Glu Asn His Ile
 20 25 30

Thr Pro Glu Tyr Leu Pro Gly Arg Ile His His Thr Ala Pro Leu Glu
 35 40 45

Pro Ala Ser Lys Asp Pro Phe Ala His Ile Val Ile Leu
 50 55 60

<210> 293

<211> 112

<212> PRT

<213> Homo sapien

<400> 293

Met Gly Ile Ile Leu Asn Trp Leu Asn Gln Trp Ala Gln Ile Thr Tyr
 1 5 10 15

Leu Pro Ser Leu Leu Cys Asp Ser Pro Ala Val Thr His Thr Ile His
 20 25 30

Ile Leu Cys Thr Ser Asn Glu Gln Thr Trp Phe Pro Cys Phe Leu Asp
 35 40 45

Ile Ser Met Thr Val Ser His Thr Asn Tyr Trp Val Arg Phe Phe Ser
 50 55 60

Cys Tyr Arg Pro Thr Ser Cys Cys Leu Cys Val Val Leu Gln Lys Leu
 65 70 75 80

Ser Ile Pro Thr Pro Leu Leu Cys His Leu Gln Glu Ser Gly Ile Val
 85 90 95

Arg Ser Gln Leu Arg Lys Val Leu Val Pro Leu Thr Gly His Ile Leu
 100 105 110

<210> 294

<211> 55

<212> PRT

<213> Homo sapien

216

<400> 294

Met Arg Phe Ile Phe Ile Cys Lys Pro Arg Gly Leu Ile Ile Leu Ile
 1 5 10 15

Leu Tyr Glu Tyr Thr Cys Val Leu Gly Lys Ala Phe Ile Gln Gln Met
 20 25 30

Pro Thr Thr Tyr Ser Val Pro Arg Pro Arg His Pro Val Thr Ser Trp
 35 40 45

Arg Pro Ala Arg Ala Cys Ile
 50 55

<210> 295

<211> 77

<212> PRT

<213> Homo sapien

<400> 295

Met Leu Glu Leu Pro Thr Phe Ser Phe Phe Phe Gly Asp Arg Ala
 1 5 10 15

Ser Leu Cys His Pro Gly Trp Ser Ala Gly Ala Ser Ser Leu Thr His
 20 25 30

Leu Gln Pro Ser Phe Leu Pro Trp Gly Ala Gly Arg Phe Ser Cys Ala
 35 40 45

Leu Gln Pro Pro Ser Leu Ala Gly Ile Tyr Arg Ala Leu Leu Gln Val
 50 55 60

Ser His Ile Phe Ser Glu Lys Phe Leu Asn Trp Pro Pro
 65 70 75

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US02/04197

A. CLASSIFICATION OF SUBJECT MATTER
 IPC(7) :C07H 21/02, 21/04
 US CL :536/23.1, 24.3
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 U.S. : 536/23.1, 24.3

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 Search the USPTO's database of nucleic acid and protein sequences.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	VINCENT et al. Oligonucleotides as short as 7-mers can be used for PCR amplification. DNA and Cell Biology. 1994, Vol. 13, No. 1, pages 75-82, note the octomer in the legend of Figure 5 and the heptamer in the legend of Figure 5 and compare to positions 60-67 of SEQ ID NO: 1.	1-5 and 7-8
X	SOMMER et al. Minimal homology requirements for PCR primers. Nucleic Acids Research. 1989, Vol. 17, No. 16, page 6749, note the first primer listed in Table I and compare the 3 nucleotides on its 3' end to the complementary nucleotides at positions 252-254 of SEQ ID NO: 1.	1-5 and 7-8

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Z" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search: 31 MAY 2002
 Date of mailing of the international search report: 28 JUN 2002

Name and mailing address of the ISA/US Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231
 Facsimile No. (703) 305-9230
 Authorized officer: 
 ETHAN WHISENANT, PH.D.
 Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US02/04197**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-5 and 7-8, SEQ ID NO: 1

Remark on Protest

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/04197

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

1. This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

I. Claim(s) 1-5, 7-8, drawn to polynucleotides, vectors comprising the polynucleotide, methods of introducing the polynucleotide to host cells, and host cells comprising the polynucleotide. This group comprises at least 171 different embodiments. See the requirement to elect a single sequence below.

II. Claim(s) 10-11, drawn to an isolated polypeptide. This group comprises at least 171 different embodiments. See the requirement to elect a single sequence below.

III. Claim(s) 12, drawn to antibodies. This group comprises at least 171 different embodiments. See the requirement to elect a single sequence below.

IV. Claim(s) 9, drawn to a method of synthesizing a polypeptide, classified in class 435, subclass 69.1. This group comprises at least 171 different embodiments. See the requirement to elect a single sequence below.

V. Claim(s) 6 and 14, drawn to a diagnostic method of using the polynucleotide of Claim 1. This group comprises at least 171 different embodiments. See the requirement to elect a single sequence below.

VI. Claim(s) 13-14, drawn to a diagnostic method of using a polypeptide. This group comprises at least 171 different embodiments. See the requirement to elect a single sequence below.

VII. Claim(s) 15, drawn to a kit for determining the presence of the nucleic acid molecule of claim 1 or the polypeptide of Claim 11. This group comprises at least 171 different embodiments. See the requirement to elect a single sequence below.

VIII. Claim(s) 16, drawn to a therapeutic method of using a polypeptide. This group comprises at least 171 different embodiments. See the requirement to elect a single sequence below.

IX. Claim(s) 17, drawn to a vaccine comprising the polypeptide of Claim 11 or a polynucleotide encoding said polypeptide of Claim 11. This group comprises at least 171 different embodiments. See the requirement to elect a single sequence below.

2. Sequence Election Requirement Applicable to All Groups

In addition, each Group detailed above reads on distinct sequences. Each sequence is distinct because they are unrelated sequences, and a further restriction is applied to each Group. For an selected Group drawn to amino acid sequences, the Applicants must elect a single amino acid sequence. For an elected Group drawn to nucleotide sequences, the Applicants must elect one nucleic acid sequence.

Examination will be restricted to only the elected sequence.

3. The inventions listed as Groups I-IX do not relate to a single

INTERNATIONAL SEARCH REPORT

International application No. .
PCT/US02/04197

general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical feature(s).

The claims as drawn are related to each other because of the product i.e. the isolated nucleic acid molecule of Claim 1. However, since the isolated nucleic acid molecule of Claim 1, as claimed, is known, the claims are no longer linked by a special technical feature, because, by definition, the special technical feature must distinguish over the prior art. Without the special technical feature the claims lack unity.